

# MPRA

Munich Personal RePEc Archive

## **Downturn LGD: A Spot Recovery Approach**

Li, Hui

13 January 2010

Online at <https://mpra.ub.uni-muenchen.de/20010/>  
MPRA Paper No. 20010, posted 14 Jan 2010 15:58 UTC

# Downturn LGD: A Spot Recovery Approach

Hui Li\*

January 13, 2010

## Abstract

Basel II suggests that banks estimate downturn loss given default (DLGD) in capital requirement calculation. There have been studies that focused on the dependence of default rates and loss given defaults through economic cycles. However, the models proposed are still not satisfactory. In this paper, we propose a new model framework based on our recent work of stochastic spot recovery for Gaussian copula. We also compare our model with the previous approaches.

## 1. Introduction

Evidence from historic data suggests that recovery rates on corporate defaults tend to go down when default rates go up in an economic downturn [1]. This phenomenon leads the BIS to suggest banks estimate downturn loss given default (DLGD) for capital requirement calculation [4, 5]. The main reason for this requirement is that the Vasicek model [20] used in the Basel Accord does not have systematic correlation between probability of default (PD) and loss given default (LGD), which would underestimate downturn risk.

There have been several attempts to model the dependence between PD and LGD, see for example [2, 3, 7, 8, 9, 10, 11, 16, 17, 18, 19, 21]. All these approaches model the period loss given default by assuming it is driven by a latent variable that is correlated with the latent variable driving default. This kind of approach has some drawbacks, as will be discussed in section 2 of the current paper. The key point is that the relationship between expected loss and probability of default may produce results not supported by economic evidence or intuition. Similar problems in CDO pricing with stochastic recovery have lead to the direct modeling of spot recovery (or recovery upon default) to avoid the problems [6, 14]. The purpose of this paper is to use our recently proposed stochastic spot recovery model for Gaussian copula to build a consistent downturn LGD model for Basel II capital calculation.

The paper is organized as follows. In section 2, we discuss the Tasche model [20] and other factor models [7] and show that they may have features not supported by economic evidence. In section 3, we present our stochastic spot recovery model using the two factor setup for a homogeneous portfolio discussed in [7]. We also derive the Gaussian copula

---

\* Email: [xhuili@yahoo.com](mailto:xhuili@yahoo.com). The opinions expressed in this article are the author's own and do not reflect in any way the views of his employer.

that correlates both default time and recovery rate. A special form of recovery distribution is presented to be used in capital calculation. In section 4, we derive the large homogeneous pool limit for the Tasche model, the Chabaane-Laurent-Salomon model and our spot recovery model in the single factor case. Then we show how VaR and expected shortfall can be calculated and define the downturn LGD for all these models. In section 5, we give numerical examples to compare the downturn LGD in these models. Section 6 concludes the paper.

## 2. Problems with current LGD factor models

The models proposed for downturn LGD are mostly factor models except that of Giese [10], where the conditional expected LGD was specified to have non-linear dependence structure on the conditional PD. There are two types of factor models. The Tasche model [19] assumes the same latent variable drives both default and loss given default so that the latent variable is actually driving the unconditional loss, see also our discussion in [13]. All other models assume a correlated latent variable drives the loss given default, where the difference is in the number of systematic or idiosyncratic factors. Frye [9] uses a single systematic factor with an independent idiosyncratic factor to drive the loss given default. Pykhtin [17] also uses a single systematic factor but with an idiosyncratic factor that is correlated with the idiosyncratic factor driving default. Barco [3] assumes two systematic factors but no idiosyncratic factor. Chabaane, Laurent and Salomon [7] discusses a more general factor correlation structure which is equivalent to two correlated systematic factors and two correlated idiosyncratic factors. The two types of models both have problem with the relationship between expected loss and probability of default, which has been touched on in our previous paper [13] and will be discussed below. We will see in section 3 the problem can be solved in a stochastic spot recovery model.

### 2.1. The Tasche Model

First we discuss the Tasche model [19] following our previous work [13]. Let  $L$  be the unconditional loss before time  $t$  as a percentage of the total exposure to an obligor. Then  $L$  will be zero with probability  $1 - p$  when the obligor is not in default before time  $t$ .  $L$  may take positive values with probability  $p$  when the obligor defaults before time  $t$ . Formally, the cumulative distribution function  $F_L$  of  $L$  has the following general form (see Tasche [19])

$$F_L(l) = P(L \leq l) = 1 - p + p \cdot F_D(l) \quad \text{for } l \in [0,1] \quad (1)$$

where  $F_D(l) = P(L \leq l | \tau \leq t)$  is the cumulative distribution of loss given default and  $\tau$  is the default time random variable. We will not make the assumption of hard default where obligor default is equivalent to loss greater than zero. So  $F_D(0) > 0$  is possible in the current framework. Define the generalized inverse or quantile function  $F_L^{-1}$  of  $F_L$  as

$$F_L^{-1}(y) = \min\{l \in [0,1] : F_L(l) \geq y\} \quad \text{for } y \in [0,1] \quad (2)$$

Assume default of an obligor is determined by the latent variable  $V = \sqrt{\rho}Z + \sqrt{1-\rho}\varepsilon$  through a default threshold  $v = \Phi^{-1}(p)$ , where  $Z$  and  $\varepsilon$  are independent standard normal random variables  $\sim N(0,1)$ ,  $Z$  is the systematic factor and  $\Phi(x)$  is the standard cumulative normal distribution function. Then we can model the dependence of loss and default by defining

$$L = F_L^{-1}(\Phi(-V)) \quad (3)$$

where the negative sign is meant to introduce a negative correlation between loss and asset value represented by the latent variable. Note that this definition will not change the distribution of  $L$ , which is still  $F_L$ .

Conditional on  $Z = z$ , the probability of default is

$$P(z) = P(V \leq \Phi^{-1}(p) | Z = z) = \Phi\left(\frac{\Phi^{-1}(p) - \sqrt{\rho}z}{\sqrt{1-\rho}}\right) \quad (4)$$

The conditional cumulative loss distribution is

$$\begin{aligned} P(L = F_L^{-1}(\Phi(-V)) \leq l | Z = z) &= \Phi\left(\frac{\Phi^{-1}(F_L(l)) + \sqrt{\rho}z}{\sqrt{1-\rho}}\right) \\ &= \Phi\left(\frac{-\Phi^{-1}(p(1 - F_D(l))) + \sqrt{\rho}z}{\sqrt{1-\rho}}\right) \\ &\equiv 1 - P(z) + P(z) \cdot P(L \leq l | \tau < t, Z = z) \end{aligned} \quad (5)$$

The last line in the above equation is just the definition of loss given default conditional on  $z$ . So

$$P(L \leq l | \tau < t, Z = z) = 1 - P(z)^{-1} \cdot \Phi\left(\frac{\Phi^{-1}(p(1 - F_D(l))) - \sqrt{\rho}z}{\sqrt{1-\rho}}\right) \quad (6)$$

The conditional expected loss is

$$\begin{aligned}
E(L|Z = z) &= \int_0^1 l \cdot d_l P(L \leq l | Z = z) \\
&= P(z) \left( 1 - \int_0^1 P(L \leq l | \tau < t, Z = z) \cdot dl \right) \\
&= \int_0^1 \Phi \left( \frac{\Phi^{-1}(p(1 - F_D(l))) - \sqrt{\rho}z}{\sqrt{1 - \rho}} \right) \cdot dl
\end{aligned} \tag{7}$$

Next we compare the change in conditional expected loss and the change in conditional probability of default induced by an infinitesimal change in marginal probability of default  $p$  :

$$\begin{aligned}
\frac{dE(L|Z = z)/dp}{dP(z)/dp} &= \int_0^1 \frac{d\Phi \left( \frac{\Phi^{-1}(p(1 - F_D(l))) - \sqrt{\rho}z}{\sqrt{1 - \rho}} \right) / dp}{d\Phi \left( \frac{\Phi^{-1}(p) - \sqrt{\rho}z}{\sqrt{1 - \rho}} \right) / dp} \cdot dl \\
&= \int_0^1 \exp \left( \frac{\Phi^{-1}(p)^2 - \Phi^{-1}(p(1 - F_D(l)))^2 - 2\sqrt{\rho}z(\Phi^{-1}(p) - \Phi^{-1}(p(1 - F_D(l))))}{2(1 - \rho)} \right. \\
&\quad \left. - \frac{\Phi^{-1}(p)^2 - \Phi^{-1}(p(1 - F_D(l)))^2}{2} \right) \cdot (1 - F_D(l)) \cdot dl
\end{aligned} \tag{8}$$

It is obvious that, when  $z \rightarrow -\infty$ , the above ratio will go to  $+\infty$ . This means that, when  $z$  is sufficient negative, conditional expected loss will increase at a much higher speed than that of conditional PD. This does not make sense since PD is equivalent to 100% loss and will always dominant expected loss such that the ratio should never exceed one. The other argument against the model is that, since default and recovery are driven by the same latent variable, the model is too restrictive and may not be able to calibrate to economic data.

## 2.2. Other LGD Factor Models

Chabaane, Laurent and Salomon [7] discussed the general factor structure for the underlying latent variables driving default and recovery under the assumption of a homogeneous credit portfolio. This general structure covers the models of Frye [9], Pykhtin [17] and Barco [3]. We will use their setup to discuss the problem with this type of models.

Again, we assume  $V = \sqrt{\rho}Z + \sqrt{1 - \rho}\varepsilon$  drives the default of an obligor. The latent variable driving LGD has the following form

$$W = \sqrt{\beta}(\eta Z + \sqrt{1-\eta^2} Z_r) + \sqrt{1-\beta}(\gamma \varepsilon + \sqrt{1-\gamma^2} \xi) \quad (9)$$

where  $Z$ ,  $Z_r$  are independent systematic factors and  $\varepsilon$ ,  $\xi$  are independent idiosyncratic factors. Loss given default is defined as  $L = F_D^{-1}(\Phi(-W))$ . Conditional on  $Z$  and  $Z_r$ , default and loss will be independent between obligors, although they are still correlated through the idiosyncratic factors for each obligor. The conditional cumulative loss given default distribution will be

$$\begin{aligned} & P(L = F_D^{-1}(\Phi(-W)) \leq l | \tau \leq t, Z = z, Z_r = z_r) \\ &= P\left(\gamma \varepsilon + \sqrt{1-\gamma^2} \xi \geq -\frac{\Phi^{-1}(F_D(l)) + \sqrt{\beta}(\eta z + \sqrt{1-\eta^2} z_r)}{\sqrt{1-\beta}} \mid \tau \leq t, Z = z, Z_r = z_r\right) \quad (10) \\ &= P(z)^{-1} \cdot \Phi_2\left(\frac{\Phi^{-1}(F_D(l)) + \sqrt{\beta}(\eta z + \sqrt{1-\eta^2} z_r)}{\sqrt{1-\beta}}, \frac{\Phi^{-1}(p) - \sqrt{\rho} z}{\sqrt{1-\rho}}; -\gamma\right) \end{aligned}$$

where  $\Phi_2(x, y; \rho)$  is the cumulative bivariate normal distribution with correlation  $\rho$ .

So

$$\begin{aligned} & P(L \leq l | Z = z, Z_r = z_r) \\ &= 1 - P(z) + \Phi_2\left(\frac{\Phi^{-1}(F_D(l)) + \sqrt{\beta}(\eta z + \sqrt{1-\eta^2} z_r)}{\sqrt{1-\beta}}, \frac{\Phi^{-1}(p) - \sqrt{\rho} z}{\sqrt{1-\rho}}; -\gamma\right) \quad (11) \end{aligned}$$

such that, after integration over  $z$  and  $z_r$ ,

$$P(L \leq l) = 1 - p + \Phi_2(\Phi^{-1}(F_D(l)), \Phi^{-1}(p); -K) \quad (12)$$

where  $K = \eta\sqrt{\rho\beta} + \gamma\sqrt{(1-\rho)(1-\beta)}$  is the correlation between  $V$  and  $W$ . We have used the following formula in the integration, see Appendix in [13],

$$\int_{-\infty}^{+\infty} \Phi_2(az + b, cz + d; \rho) \cdot \phi(z) dz = \Phi_2\left(\frac{b}{\sqrt{1+a^2}}, \frac{d}{\sqrt{1+c^2}}; \frac{ac + \rho}{\sqrt{(1+a^2)(1+c^2)}}\right) \quad (13)$$

So the marginal loss given default distribution is

$$F_D^M(l) = P(L \leq l | \tau \leq t) = p^{-1} \cdot \Phi_2(\Phi^{-1}(F_D(l)), \Phi^{-1}(p); -K) \quad (14)$$

Note that the marginal loss given default distribution  $F_D^M$  is different from  $F_D$  unless the correlation term in the above formula is zero. The correlation term is always negative, which make sense since increase in  $F_D$  means decrease in the expected LGD. However, when marginal probability of default  $p$  increases,  $F_D^M$  will always increase which leads to the decrease of the expected LGD. This is counter-intuitive and is not supported by any economic evidence. So this kind of models has an unwelcome side-effect. Besides, the distribution of marginal LGD and the expected LGD will depend on PD and correlation, which makes the model calibration more complicated.

### 3. Stochastic Recovery in the Default Time Copula Framework

The problems of the previous section can be solved through a stochastic spot recovery model in a default time copula framework. The term structure of default probability curve means that PD is always increasing with time. So the problem with the Tasche model is equivalent to an unlimited conditional spot LGD or negative conditional spot recovery, see [13]. The way to solve it would be to model the spot LGD or spot recovery directly to make sure it is in proper range. Meanwhile, the Tasche model preserves the marginal loss distribution, which is a problem for other factor models. So we hope this is also preserved in the spot recovery model, which is indeed the case [14]. Here we generalize our one-factor Gaussian model of spot recovery [14, 15] to two systematic factors with correlation between idiosyncratic variables. We will follow the factor structure of Chabaane, Laurent and Salomon [7] for a homogeneous credit portfolio. It is straight forward to extend the model to multi-factor or non-Gaussian copula cases.

In the default time copula framework of D. Li [12], the joint distribution of default times is determined by the marginal default time distributions (given by default probability curve) and the default time copula. In the Gaussian Copula setup, the latent variable  $V_i = \sqrt{\rho}Z + \sqrt{1-\rho}\varepsilon_i$  drives the default of obligor  $i$  of a homogeneous credit portfolio. The default event  $1_{\tau_i \leq t}$  can be characterized by  $V_i \leq v = \Phi^{-1}(p(t))$ , where  $\tau_i$  is the default time random variable,  $p(t)$  is the cumulative default probability of the obligor  $i$ . We define the default time random variable  $\tau_i$  as

$$\tau_i = p^{-1}(\Phi(V_i)) \quad (15)$$

We assume that the stochastic spot recovery is driven by the latent variable  $W_i = \sqrt{\beta}(\eta Z + \sqrt{1-\eta^2}Z_r) + \sqrt{1-\beta}(\gamma\varepsilon_i + \sqrt{1-\gamma^2}\xi_i)$  through a time-independent cumulative distribution function  $F_R(r)$ .

Conditional on  $\tau_i = t$  or  $V_i = \Phi^{-1}(p(t))$ ,  $W_i$  follows a normal distribution with mean  $K \cdot \Phi^{-1}(p(t))$  and standard deviation  $\sqrt{1-K^2}$ , where  $K = \eta\sqrt{\rho\beta} + \gamma\sqrt{(1-\rho)(1-\beta)}$ . To

ensure that  $F_R(r)$  is indeed the marginal cumulative distribution for the spot recovery upon default at time  $t$ , we define

$$R_i = F_R^{-1} \left( \Phi \left( \frac{W_i - K \cdot \Phi^{-1}(p(t))}{\sqrt{1 - K^2}} \right) \right) \quad (16)$$

Thus

$$P(R_i \leq r | \tau_i = t) = P \left( F_R^{-1} \left( \Phi \left( \frac{W_i - K \cdot \Phi^{-1}(p(t))}{\sqrt{1 - K^2}} \right) \right) \leq r | \tau_i = t \right) = F_R(r) \quad (17)$$

If we fix  $Z = z$  and  $Z_r = z_r$ , then  $\varepsilon_i = \frac{\Phi^{-1}(p(t)) - \sqrt{\rho}z}{\sqrt{1 - \rho}}$  and the conditional spot recovery distribution will be

$$\begin{aligned} & P(R_i \leq r | \tau_i = t, Z = z, Z_r = z_r) \\ &= P \left( F_R^{-1} \left( \Phi \left( \frac{W_i - K \cdot \Phi^{-1}(p(t))}{\sqrt{1 - K^2}} \right) \right) \leq r | \tau_i = t, Z = z, Z_r = z_r \right) \\ &= \Phi \left( \frac{-Dz - \sqrt{\beta(1 - \rho)(1 - \eta^2)}z_r + \sqrt{(1 - \rho)(1 - K^2)} \cdot \Phi^{-1}(F_R(r)) + D\sqrt{\rho} \cdot \Phi^{-1}(p(t))}{\sqrt{(1 - \rho)(1 - \beta)(1 - \gamma^2)}} \right) \end{aligned} \quad (18)$$

where  $D = \eta\sqrt{\beta(1 - \rho)} - \gamma\sqrt{\rho(1 - \beta)}$ . If  $D = 0$ , then  $W_i$  is linear in  $V_i$ . We may require  $D > 0$  such that, when  $z$  increases, the conditional cumulative distribution decreases and conditional expected recovery will increase. Conditional on the systematic factor  $Z$ , obligor defaults are independent and the conditional default probability for each obligor is given by

$$p(t, z) = p(\tau \leq t | Z = z) = \Phi \left( \frac{\Phi^{-1}(p(t)) - \sqrt{\rho}z}{\sqrt{1 - \rho}} \right) \quad (19)$$

Now we can derive the distribution for conditional period recovery rate defined as

$$\begin{aligned} & P(R_i \leq r | \tau_i \leq t, Z = z, Z_r = z_r) \\ &= \frac{1}{p(t, z)} \cdot \int_0^t \Phi \left( \frac{-Dz - \sqrt{\beta(1 - \rho)(1 - \eta^2)}z_r + \sqrt{(1 - \rho)(1 - K^2)} \cdot \Phi^{-1}(F_R(r)) + D\sqrt{\rho} \cdot \Phi^{-1}(p(s))}{\sqrt{(1 - \rho)(1 - \beta)(1 - \gamma^2)}} \right) \cdot dp(s, z) \\ &= \frac{1}{p(t, z)} \cdot \Phi_2 \left( \frac{-D\sqrt{1 - \rho}z - \sqrt{\beta(1 - \eta^2)}z_r + \sqrt{1 - K^2}\Phi^{-1}(F_R(r))}{\sqrt{(1 - \beta)(1 - \gamma^2) + D^2\rho}}, c(t, z); -\tilde{\rho} \right) \end{aligned} \quad (20)$$



where

$$c(t, z) = \frac{\Phi^{-1}(p(t)) - \sqrt{\rho}z}{\sqrt{1-\rho}} = \Phi^{-1}(p(t, z)) \text{ and } \tilde{\rho} = \frac{D\sqrt{\rho}}{\sqrt{(1-\beta)(1-\gamma^2) + D^2\rho}}$$

We also have

$$\begin{aligned} & P(1_{\{\tau_i \leq t\}} \cdot 1_{\{R_i \leq r\}} \mid Z = z, Z_r = z_r) \\ &= P(R_i \leq r \mid \tau_i \leq t, Z = z, Z_r = z_r) \cdot P(\tau_i \leq t \mid Z = z) \\ &= \Phi_2 \left( \frac{-D\sqrt{1-\rho}z - \sqrt{\beta(1-\eta^2)}z_r + \sqrt{1-K^2}\Phi^{-1}(F_R(r))}{\sqrt{(1-\beta)(1-\gamma^2) + D^2\rho}}, c(t, z); -\tilde{\rho} \right) \end{aligned} \quad (21)$$

The unconditional period recovery distribution can be calculated using equation (13) as follows

$$P(R_i \leq r \mid \tau_i \leq t) = \frac{1}{p(t)} \iint P(1_{\{\tau_i \leq t\}} \cdot 1_{\{R_i \leq r\}} \mid Z = z, Z_r = z_r) \cdot \phi(z)\phi(z_r) dz dz_r = F_R(r) \quad (22)$$

So the marginal distribution of period recovery rate is the same as the marginal distribution of spot recovery rate and is time-independent. Note that, in a dynamic model, the spot recovery distribution  $F_R(r)$  could be time dependent, then the integration in equation (20) would be more complicated.

Consider two obligors with correlated default and recovery rate, here we derive the copula of default time and recovery rate. Conditional on  $Z$  and  $Z_r$ , the default and recovery process are independent for the two obligors, and we have

$$\begin{aligned} & P(1_{\{\tau_1 \leq t_1\}} \cdot 1_{\{R_1 \leq r_1\}} \cdot 1_{\{\tau_2 \leq t_2\}} \cdot 1_{\{R_2 \leq r_2\}} \mid Z = z, Z_r = z_r) \\ &= \Phi_2 \left( \frac{-D_1\sqrt{1-\rho_1}z - \sqrt{\beta_1(1-\eta_1^2)}z_r + \sqrt{1-K_1^2}\Phi^{-1}(F_{R_1}(r_1))}{\sqrt{(1-\beta_1)(1-\gamma_1^2) + D_1^2\rho_1}}, c_1(t, z); -\tilde{\rho}_1 \right) \\ & \cdot \Phi_2 \left( \frac{-D_2\sqrt{1-\rho_2}z - \sqrt{\beta_2(1-\eta_2^2)}z_r + \sqrt{1-K_2^2}\Phi^{-1}(F_{R_2}(r_2))}{\sqrt{(1-\beta_2)(1-\gamma_2^2) + D_2^2\rho_2}}, c_2(t, z); -\tilde{\rho}_2 \right) \end{aligned} \quad (23)$$

Integrating over  $z$  and  $z_r$ , we will have the copula as

$$\begin{aligned}
C(p_1(t_1), F_{R_1}(r_1); p_2(t_2), F_{R_2}(r_2)) &= P(\tau_1 \leq t_1, R_1 \leq r_1; \tau_2 \leq t_2, R_2 \leq r_2) \\
&= \iint P(\mathbf{1}_{\{\tau_1 \leq t_1\}} \cdot \mathbf{1}_{\{R_1 \leq r_1\}} \cdot \mathbf{1}_{\{\tau_2 \leq t_2\}} \cdot \mathbf{1}_{\{R_2 \leq r_2\}} \mid Z = z, Z_r = z_r) \cdot \phi(z) \phi(z_r) dz dz_r \quad (24) \\
&= \Phi_4(\Phi^{-1}(p_1(t_1)), \Phi^{-1}(F_{R_1}(r_1)), \Phi^{-1}(p_2(t_2)), \Phi^{-1}(F_{R_2}(r_2)); \Sigma_\rho)
\end{aligned}$$

where  $\Phi_4$  is the 4-variable cumulative normal distribution and the correlation matrix is defined as

$$\Sigma_\rho = \begin{pmatrix} 1 & 0 & \sqrt{\rho_1 \rho_2} & \frac{D_1 \sqrt{\rho_1 (1-\rho_2)}}{\sqrt{1-K_2^2}} \\ 0 & 1 & \frac{D_2 \sqrt{\rho_2 (1-\rho_1)}}{\sqrt{1-K_1^2}} & \frac{D_1 D_2 \sqrt{\rho_1 \rho_2 (1-\rho_1)(1-\rho_2) + \sqrt{\beta_1 \beta_2 (1-\eta_1^2)(1-\eta_2^2)}}}{\sqrt{(1-K_1^2)(1-K_2^2)}} \\ \sqrt{\rho_1 \rho_2} & \frac{D_1 \sqrt{\rho_2 (1-\rho_1)}}{\sqrt{1-K_1^2}} & 1 & 0 \\ \frac{D_2 \sqrt{\rho_1 (1-\rho_2)}}{\sqrt{1-K_2^2}} & \frac{D_1 D_2 \sqrt{\rho_1 \rho_2 (1-\rho_1)(1-\rho_2) + \sqrt{\beta_1 \beta_2 (1-\eta_1^2)(1-\eta_2^2)}}}{\sqrt{(1-K_1^2)(1-K_2^2)}} & 0 & 1 \end{pmatrix}$$

This can be proven through the following result

$$\begin{aligned}
&\iint \Phi_2(a_1 z + b_1, c_1 z + d_1 + e_1 z_r; \rho_1) \cdot \Phi_2(a_2 z + b_2, c_2 z + d_2 + e_2 z_r; \rho_2) \cdot \phi(z) \phi(z_r) dz dz_r \\
&= \Phi_4 \left( \frac{b_1}{\sqrt{1+a_1^2}}, \frac{d_1}{\sqrt{1+c_1^2+e_1^2}}, \frac{b_2}{\sqrt{1+a_2^2}}, \frac{d_2}{\sqrt{1+c_2^2+e_2^2}}; \Sigma \right) \quad (25)
\end{aligned}$$

where

$$\Sigma = \begin{pmatrix} 1 & \frac{\rho_1 + a_1 c_1}{\sqrt{(1+a_1^2)(1+c_1^2+e_1^2)}} & \frac{a_1 a_2}{\sqrt{(1+a_1^2)(1+a_2^2)}} & \frac{a_1 c_2}{\sqrt{(1+a_1^2)(1+c_2^2+e_2^2)}} \\ \frac{\rho_1 + a_1 c_1}{\sqrt{(1+a_1^2)(1+c_1^2+e_1^2)}} & 1 & \frac{c_1 a_2}{\sqrt{(1+c_1^2+e_1^2)(1+a_2^2)}} & \frac{c_1 c_2 + e_1 e_2}{\sqrt{(1+c_1^2+e_1^2)(1+c_2^2+e_2^2)}} \\ \frac{a_1 a_2}{\sqrt{(1+a_1^2)(1+a_2^2)}} & \frac{c_1 a_2}{\sqrt{(1+c_1^2+e_1^2)(1+a_2^2)}} & 1 & \frac{\rho_2 + a_2 c_2}{\sqrt{(1+a_2^2)(1+c_2^2+e_2^2)}} \\ \frac{a_1 c_2}{\sqrt{(1+a_1^2)(1+c_2^2+e_2^2)}} & \frac{c_1 c_2 + e_1 e_2}{\sqrt{(1+c_1^2+e_1^2)(1+c_2^2+e_2^2)}} & \frac{\rho_2 + a_2 c_2}{\sqrt{(1+a_2^2)(1+c_2^2+e_2^2)}} & 1 \end{pmatrix}$$

If we define

$$\begin{cases} X_1 = \sqrt{\rho_1} \varepsilon + \sqrt{1-\rho_1} \varepsilon_1 - a_1 Z \\ Y_1 = \sqrt{\rho_1} \varepsilon + \sqrt{1-\rho_1} \varepsilon_2 - c_1 Z - e_1 Z_r \\ X_2 = \sqrt{\rho_2} \xi + \sqrt{1-\rho_2} \xi_1 - a_2 Z \\ Y_2 = \sqrt{\rho_2} \xi + \sqrt{1-\rho_2} \xi_2 - c_2 Z - e_2 Z_r \end{cases}$$

where  $\varepsilon, \varepsilon_1, \varepsilon_2, \xi, \xi_1, \xi_2, Z, Z_r$  are independent standard normal random variables, then

$$\begin{aligned} & \iint \Phi_2(a_1 z + b_1, c_1 z + d_1 + e_1 z_r; \rho_1) \cdot \Phi_2(a_2 z + b_2, c_2 z + d_2 + e_2 z_r; \rho_2) \cdot \phi(z) dz \cdot \phi(z_r) dz_r \\ & = E(X_1 \leq b_1, Y_1 \leq d_1, X_2 \leq b_2, Y_2 \leq d_2) \end{aligned}$$

which leads to the equation (25).

Equation (24) can be compared with the standard Gaussian copula of default times with fixed recovery


$$C(p_1(t_1), p_2(t_2)) = P(\tau_1 \leq t_1, \tau_2 \leq t_2) = \Phi_2(\Phi^{-1}(p_1(t_1)), \Phi^{-1}(p_2(t_2)); \sqrt{\rho_1 \rho_2}) \quad (26)$$

Note that, in equation (24), default and recovery of an obligor are not correlated, this is because recovery is always conditional on default. The copula for default and recovery is still Gaussian. However, the correlation matrix can not be generated by a simple one-factor model. Equation (24) can be easily extended to more than two obligors, multi-factors and other types of copulas.

For capital calculation, we need the conditional expected loss for each obligor before time  $t$

$$\begin{aligned} L_t^i(z, z_r) &= \int_0^1 (1-r) \cdot d_r P(1_{\{\tau_i \leq t\}} \cdot 1_{\{R_i \leq r\}} \mid Z = z, Z_r = z_r) \\ &= \int_0^1 P(1_{\{\tau_i \leq t\}} \cdot 1_{\{R_i \leq r\}} \mid Z = z, Z_r = z_r) \cdot dr \end{aligned} \quad (27)$$

For numeric purpose, we consider the recovery distribution discussed in [14], which is similar to the beta distribution as shown in the Figure below.



It has the following form

$$F_R(r) = P(R \leq r) = \Phi(a \cdot \Phi^{-1}(r) - \sqrt{1+a^2} \Phi^{-1}(r_0)) \quad (28)$$

or, for the density function,

$$f_R(r) = a \cdot \frac{\phi(a \cdot \Phi^{-1}(r) - \sqrt{1+a^2} \Phi^{-1}(r_0))}{\phi(\Phi^{-1}(r))} \quad (29)$$

where  $a \geq 0$  and  $0 \leq r_0 \leq 1$ . This distribution will simplify calculation for Gaussian Copula model. The expected recovery rate is  $r_0$  and the variance of recovery rate is

$$V(R) = \Phi_2\left(\Phi^{-1}(r_0), \Phi^{-1}(r_0); \frac{1}{1+a^2}\right) - r_0^2 \quad (30)$$

When  $a$  goes to zero, the variance goes to the maximum value  $r_0(1-r_0)$ , which corresponds to the case where  $R$  takes the extreme value 0 or 1. When  $a$  goes to infinity, the variance goes to zero and the distribution reduces to a constant recovery  $r_0$ .

The original spot recovery equation (16) can be written as

$$R = \Phi \left( \frac{W_i - K\Phi^{-1}(p(t))}{a\sqrt{1-K^2}} + \sqrt{1 + \frac{1}{a^2}} \Phi^{-1}(r_0) \right) \quad (31)$$

Then we have

$$\begin{aligned} & P(R \leq r | \tau = t, Z = z, Z_r = z_r) \\ &= \Phi \left( \frac{-Dz - \sqrt{\beta(1-\rho)(1-\eta^2)}z_r + \sqrt{(1-\rho)(1-K^2)} \cdot (a\Phi^{-1}(r) - \sqrt{1+a^2}\Phi^{-1}(r_0)) + D\sqrt{\rho} \cdot \Phi^{-1}(p(t))}{\sqrt{(1-\rho)(1-\beta)(1-\gamma^2)}} \right) \end{aligned} \quad (32)$$

The expected conditional spot recovery is

$$\begin{aligned} r(t, z, z_r) &= \int_0^1 r \cdot d_r P(R \leq r | \tau = t, Z = z, Z_r = z_r) \\ &= \Phi \left( \frac{Dz + \sqrt{\beta(1-\rho)(1-\eta^2)}z_r + \sqrt{(1-\rho)(1-K^2)} \cdot \sqrt{1+a^2}\Phi^{-1}(r_0) - D\sqrt{\rho} \cdot \Phi^{-1}(p(t))}{\sqrt{(1-\rho)(1-\beta)(1-\gamma^2) + a^2(1-\rho)(1-K^2)}} \right) \end{aligned} \quad (33)$$

The expected conditional loss up to time  $t$  is

$$L_t(z, z_r) = \int_0^t (1 - r(s, z, z_r)) \cdot dp(s, z) = \Phi_2(c(t, z), b(z, z_r); -\hat{\rho}) \quad (34)$$

where  $c(t, z)$  is defined in equation (20) and

$$\begin{aligned} b(z, z_r) &= -\frac{D\sqrt{1-\rho}z + \sqrt{\beta(1-\eta^2)}z_r + \sqrt{1-K^2}\sqrt{1+a^2}\Phi^{-1}(r_0)}{\sqrt{(1-\beta)(1-\gamma^2) + D^2\rho + a^2(1-K^2)}} \\ \hat{\rho} &= \frac{D\sqrt{\rho}}{\sqrt{(1-\beta)(1-\gamma^2) + D^2\rho + a^2(1-K^2)}} \end{aligned}$$

Conditional on  $Z$  and  $Z_r$ , the expected recovery rate will be time-dependent through  $p(t)$ .

## 4. Large Homogeneous Pool Limit and Downturn LGD

In the Basel II capital requirement calculation, the portfolio is normally assumed to be fully granular which corresponds to the large homogeneous pool (LHP) limit. We look at

the LHP limit for the Tasche model, the Chabaane-Laurent-Salomon model and our spot recovery model and compare them to the standard Vasicek model.

In all these models, conditional on the systematic factors, loss of each obligor is independent. So in the LHP limit, the portfolio loss equals to the expected loss of each obligor conditional on the systematic factors,  $L(Z)$  or  $L(Z, Z_r)$ .

In the Tasche model, the conditional expected loss is shown in equation (7). We will use the distribution in equation (28) as an example for calculation purpose. Since  $L = 1 - R$ , we have

$$F_D(l) = P(L \leq l) = P(R \geq 1 - l) = 1 - F_R(r) \quad (35)$$

where  $l = 1 - r$ . So the conditional expected loss is

$$L(z) = \int_0^1 \Phi \left( \frac{\Phi^{-1}(p(1 - F_D(l))) - \sqrt{\rho}z}{\sqrt{1 - \rho}} \right) \cdot dl = \int_0^1 \Phi \left( \frac{\Phi^{-1}(pF_R(r)) - \sqrt{\rho}z}{\sqrt{1 - \rho}} \right) \cdot dr \quad (36)$$

The portfolio loss in the LHP limit is  $L_p = L(Z)$ . The portfolio loss distribution can be calculated as

$$F_{L_p}(l) = P(L_p \leq l) = P(L(Z) \leq l) = \Phi(-L^{-1}(l)) \quad (37)$$

where the negative sign is because  $L(z)$  is a monotonically decreasing function of  $z$ . Equivalently, we have  $l = L(-\Phi^{-1}(F_{L_p}(l)))$ . This gives an easy way to calculate VaR as

$$VaR(\alpha) = F_{L_p}^{-1}(\alpha) = L(-\Phi^{-1}(\alpha)) \quad (38)$$

where  $\alpha$  is the confidence level. Specifically, for the Tasche model, we have

$$\begin{aligned} VaR(\alpha) &= \int_0^1 \Phi \left( \frac{\Phi^{-1}(pF_R(r)) + \sqrt{\rho}\Phi^{-1}(\alpha)}{\sqrt{1 - \rho}} \right) \cdot dr \\ &= \int_0^1 \Phi \left( \frac{\Phi^{-1}(p\Phi(a\Phi^{-1}(r) - \sqrt{1 + a^2}\Phi^{-1}(r_0))) + \sqrt{\rho}\Phi^{-1}(\alpha)}{\sqrt{1 - \rho}} \right) \cdot dr \end{aligned} \quad (39)$$

The expected shortfall can be calculated as

$$\begin{aligned}
ES(\alpha) &= E[L | L > VaR(\alpha)] = \frac{1}{1-\alpha} \cdot \int_{-\infty}^{-\Phi^{-1}(\alpha)} L(z) d\Phi(z) \\
&= \frac{1}{1-\alpha} \cdot \int_0^1 \Phi_2\left(\Phi^{-1}(pF_R(r)), \Phi^{-1}(1-\alpha); \sqrt{\rho}\right) \cdot dr \\
&= \frac{1}{1-\alpha} \cdot \int_0^1 \Phi_2\left(\Phi^{-1}(p\Phi(a\Phi^{-1}(r) - \sqrt{1+a^2}\Phi^{-1}(r_0))), \Phi^{-1}(1-\alpha); \sqrt{\rho}\right) \cdot dr
\end{aligned} \tag{40}$$

For the recovery distribution (28), the VaR and expected shortfall do not have analytical solution and numerical integration or Monte Carlo method has to be used for calculation.

Next, we look at the Chabaane-Laurent-Salomon model as discussed in section 2.2. For the two factor model, loss is no longer a monotonic function and calculation is more complicated. In the special case  $\gamma = 0$ , Hillebrand [11] proposed an estimation method and it was used in Barco [3] for the two systematic factor case. Here we will confine to the special case of a single systematic factor when  $\eta = 1$ , which is the Pykhtin case. The conditional cumulative LGD distribution is

$$P(L \leq l, \tau \leq t | Z = z) = \Phi_2\left(\frac{\Phi^{-1}(F_D(l)) + \sqrt{\beta}z}{\sqrt{1-\beta}}, \frac{\Phi^{-1}(p) - \sqrt{\rho}z}{\sqrt{1-\rho}}; -\gamma\right) \tag{41}$$

So the conditional expected loss is

$$\begin{aligned}
L(z) &= \int_0^1 l \cdot d_l \Phi_2\left(\frac{\Phi^{-1}(F_D(l)) + \sqrt{\beta}z}{\sqrt{1-\beta}}, \frac{\Phi^{-1}(p) - \sqrt{\rho}z}{\sqrt{1-\rho}}; -\gamma\right) \\
&= P(z) - \int_0^1 \Phi_2\left(\frac{\Phi^{-1}(F_D(l)) + \sqrt{\beta}z}{\sqrt{1-\beta}}, \frac{\Phi^{-1}(p) - \sqrt{\rho}z}{\sqrt{1-\rho}}; -\gamma\right) \cdot dl \\
&= \int_0^1 \Phi_2\left(\frac{\Phi^{-1}(F_R(r)) - \sqrt{\beta}z}{\sqrt{1-\beta}}, \frac{\Phi^{-1}(p) - \sqrt{\rho}z}{\sqrt{1-\rho}}; \gamma\right) \cdot dr
\end{aligned} \tag{42}$$

So VaR will be

$$VaR(\alpha) = \int_0^1 \Phi_2\left(\frac{\Phi^{-1}(F_R(r)) + \sqrt{\beta}\Phi^{-1}(\alpha)}{\sqrt{1-\beta}}, \frac{\Phi^{-1}(p) + \sqrt{\rho}\Phi^{-1}(\alpha)}{\sqrt{1-\rho}}; \gamma\right) \cdot dr \tag{43}$$

And expected shortfall is

$$ES(\alpha) = \frac{1}{1-\alpha} \cdot \int_0^1 \Phi_3\left(\Phi^{-1}(p), \Phi^{-1}(F_R(r)), \Phi^{-1}(1-\alpha); \Sigma_C\right) \cdot dr \tag{44}$$

where we have used equation (33) in [14],  $\Phi_3(x, y, z; \Sigma)$  is the 3-variable cumulative normal distribution and the correlation matrix is

$$\Sigma_C = \begin{pmatrix} 1 & K & \sqrt{\rho} \\ K & 1 & \sqrt{\beta} \\ \sqrt{\rho} & \sqrt{\beta} & 1 \end{pmatrix}$$

In the special case of the recovery distribution in equation (28), we have

$$VaR(\alpha) = \Phi_2 \left( \frac{-\sqrt{1+a^2}\Phi^{-1}(r_0) + \sqrt{\beta}\Phi^{-1}(\alpha)}{\sqrt{1-\beta+a^2}}, \frac{\Phi^{-1}(p) + \sqrt{\rho}\Phi^{-1}(\alpha)}{\sqrt{1-\rho}}; \frac{\gamma\sqrt{1-\beta}}{\sqrt{1-\beta+a^2}} \right) \quad (45)$$

and

$$ES(\alpha) = \frac{1}{1-\alpha} \cdot \Phi_3(\Phi^{-1}(p), \Phi^{-1}(1-r_0), \Phi^{-1}(1-\alpha); \tilde{\Sigma}_C) \quad (46)$$

where

$$\tilde{\Sigma}_C = \begin{pmatrix} 1 & \frac{K}{\sqrt{1+a^2}} & \sqrt{\rho} \\ \frac{K}{\sqrt{1+a^2}} & 1 & \frac{\sqrt{\beta}}{\sqrt{1+a^2}} \\ \sqrt{\rho} & \frac{\sqrt{\beta}}{\sqrt{1+a^2}} & 1 \end{pmatrix}$$

The correlation matrix can be easily derived by looking at pair-wise correlation through equation (13).

For our new model, again we assume  $\eta = 1$  and the results will be similar to the above except correlations. The conditional cumulative LGD distribution from equation (21) is

$$P(R \leq r, \tau \leq t | Z = z) = \Phi_2 \left( \frac{-D\sqrt{1-\rho}z + \sqrt{1-K^2}\Phi^{-1}(F_R(r))}{\sqrt{(1-\beta)(1-\gamma^2) + D^2\rho}}, \frac{\Phi^{-1}(p(t)) - \sqrt{\rho}z}{\sqrt{1-\rho}}; \frac{-D\sqrt{\rho}}{\sqrt{(1-\beta)(1-\gamma^2) + D^2\rho}} \right) \quad (47)$$

So the conditional expected loss is



$$L(z) = \int_0^1 \Phi_2 \left( \frac{-D\sqrt{1-\rho}z + \sqrt{1-K^2}\Phi^{-1}(F_R(r))}{\sqrt{(1-\beta)(1-\gamma^2) + D^2\rho}}, \frac{\Phi^{-1}(p(t)) - \sqrt{\rho}z}{\sqrt{1-\rho}}, \frac{-D\sqrt{\rho}}{\sqrt{(1-\beta)(1-\gamma^2) + D^2\rho}} \right) \cdot dr \quad (48)$$

So VaR will be

$$VaR(\alpha) = \int_0^1 \Phi_2 \left( \frac{D\sqrt{1-\rho}\Phi^{-1}(\alpha) + \sqrt{1-K^2}\Phi^{-1}(F_R(r))}{\sqrt{(1-\beta)(1-\gamma^2) + D^2\rho}}, \frac{\Phi^{-1}(p(t)) + \sqrt{\rho}\Phi^{-1}(\alpha)}{\sqrt{1-\rho}}, \frac{-D\sqrt{\rho}}{\sqrt{(1-\beta)(1-\gamma^2) + D^2\rho}} \right) \cdot dr \quad (49)$$

And expected shortfall is

$$ES(\alpha) = \frac{1}{1-\alpha} \cdot \int_0^1 \Phi_3(\Phi^{-1}(p), \Phi^{-1}(F_R(r)), \Phi^{-1}(1-\alpha); \Sigma_L) \cdot dr \quad (50)$$

where

$$\Sigma_L = \begin{pmatrix} 1 & 0 & \sqrt{\rho} \\ 0 & 1 & \frac{D\sqrt{1-\rho}}{\sqrt{1-K^2}} \\ \sqrt{\rho} & \frac{D\sqrt{1-\rho}}{\sqrt{1-K^2}} & 1 \end{pmatrix}$$

Note that the zero entry in the correlation matrix means there is no correlation between default and LGD of an obligor, same as what we saw in equation (24).

In the special case of the recovery distribution in equation (28), we have

$$VaR(\alpha) = \Phi_2 \left( \frac{-\sqrt{1-K^2}\sqrt{1+a^2}\Phi^{-1}(r_0) + D\sqrt{1-\rho}\Phi^{-1}(\alpha)}{\sqrt{(1-\beta)(1-\gamma^2) + D^2\rho + a^2(1-K^2)}}, \frac{\Phi^{-1}(p) + \sqrt{\rho}\Phi^{-1}(\alpha)}{\sqrt{1-\rho}}, \frac{-D\sqrt{\rho}}{\sqrt{(1-\beta)(1-\gamma^2) + D^2\rho + a^2(1-K^2)}} \right) \quad (51)$$

and

$$ES(\alpha) = \frac{1}{1-\alpha} \cdot \Phi_3(\Phi^{-1}(p), \Phi^{-1}(1-r_0), \Phi^{-1}(1-\alpha); \tilde{\Sigma}_L) \quad (52)$$

where

$$\tilde{\Sigma}_L = \begin{pmatrix} 1 & 0 & \frac{\sqrt{\rho}}{D\sqrt{1-\rho}} \\ 0 & 1 & \frac{D\sqrt{1-\rho}}{\sqrt{1+a^2}\sqrt{1-K^2}} \\ \sqrt{\rho} & \frac{D\sqrt{1-\rho}}{\sqrt{1+a^2}\sqrt{1-K^2}} & 1 \end{pmatrix}$$

In the limit  $a \rightarrow \infty$ , the recovery distribution converges to the constant case, which is just the original Basel II formulation with no correlation between default and LGD:

$$VaR_{Vasicek}(\alpha) = LGD \cdot \Phi\left(\frac{\Phi^{-1}(p) + \sqrt{\rho}\Phi^{-1}(\alpha)}{\sqrt{1-\rho}}\right) \quad (53)$$

and

$$ES_{Vasicek}(\alpha) = LGD \cdot \frac{\Phi_2\left(\Phi^{-1}(p), \Phi^{-1}(1-\alpha); \sqrt{\rho}\right)}{1-\alpha} \quad (54)$$

where  $LGD = 1 - r_0$  is the expected loss given default of each obligor. This limit can also be obtained if  $K = 0$ , which is equivalent to  $\beta = 0$  and  $\gamma = 0$ .

The downturn LGD (DLGD) for a general LGD model is defined as (see [3])

$$DLGD(\alpha) = \frac{VaR(\alpha)}{\Phi\left(\frac{\Phi^{-1}(p) + \sqrt{\rho}\Phi^{-1}(\alpha)}{\sqrt{1-\rho}}\right)} \quad (55)$$

which will be 1 for the Vasicek model, and may be greater than 1 for correlated models with more tail risk. We will study this phenomenon in the next section.

## 5. Numeric Examples

We present some numerical examples here to compare downturn LGD in our model with those of other models. The confidence level is set at  $\alpha = 99.9\%$ . Below is a table showing the ratio between downturn LGD and expected  $LGD = 1 - r_0$  under various parameter combinations (any parameter change is colored in yellow). The ratio is equivalent to the ratio between VaR of the correlated model and VaR of the Vasicek model.

$$ratio = \frac{DLGD(\alpha)}{LGD} = \frac{VaR(\alpha)}{VaR_{Vasicek}(\alpha)}$$

$r_0$	$a$	$p$	$\rho$	$\beta$	$\gamma$	Tasche	Chabaane	Ours
40%	10	1%	15%	15%	25%	101.9%	110.0%	104.0%
	1					112.3%	154.4%	127.0%
	0.1					116.1%	163.3%	136.7%
40%	10	5%	15%	15%	25%	102.6%	109.2%	104.4%
	1					116.7%	151.9%	129.3%
	0.1					121.7%	161.8%	139.5%
40%	10	1%	50%	15%	25%	103.2%	108.9%	101.2%
	1					120.9%	150.9%	108.5%
	0.1					127.4%	161.1%	111.9%
40%	10	1%	15%	50%	25%	101.9%	115.4%	109.0%
	1					112.3%	165.1%	153.4%
	0.1					116.1%	166.7%	164.1%
40%	10	1%	15%	15%	50%	101.9%	112.4%	103.0%
	1					112.3%	160.8%	120.8%
	0.1					116.1%	166.1%	128.7%
70%	10	1%	15%	15%	25%	103.5%	119.0%	107.3%
	1					127.5%	245.4%	155.8%
	0.1					139.6%	293.5%	180.7%
40%	10	1%	50%	15%	50%	103.2%	110.2%	99.4%
	1					120.9%	156.0%	95.4%
	0.1					127.4%	164.6%	93.6%

From the table we can see the following features:

1. The ratio generally increases with recovery volatility (decreasing with  $a$ ) ;
2. The ratio generally increases with default probability ( $p$ ) for the Tasche model and our model, but decreases for the Chabaane-Laurent-Salomon model, which is related to the problem discussed in section 2.2;
3. The ratio generally increases with default correlation ( $\rho$ ) for the Tasche model, but decreases for the Chabaane-Laurent-Salomon model and our model;
4. The ratio generally increases with  $\beta$ , however the Tasche model does not depend on  $\beta$  ;
5. The ratio increases with  $\gamma$  for the Chabaane-Laurent-Salomon model, but decreases with  $\gamma$  for our model. The Tasche model does not depend on  $\gamma$  ;
6. The ratio generally increases with the mean recovery  $r_0$ , but the actual mean recovery for the Chabaane-Laurent-Salomon model may be smaller, which leads to a higher ratio;
7. The ratio is less than 100% for our model in case  $D < 0$  which leads to negative correlation between default and LGD and is against economic evidence, in this case, the ratio is decreasing with recovery volatility (increasing with  $a$ ) ;
8. In general, the Chabaane-Laurent-Salomon model has the highest ratio, the Tasche model has lower ratio, and our model has more flexible behavior;

## 6. Conclusion

In this paper, we present a new model framework for the quantification of downturn LGD in the Basel II capital requirement. We show the problems with previous approaches which are avoided in our new model of stochastic spot recovery in a default time copula framework. We also give the large homogeneous pool limit and derive analytic formula for VaR and expected shortfall given a specific form of recovery distribution. The downturn LGD in the new model is compared with previous models with numerical examples.

Further research is required to connect the model with economic data to verify the soundness of the model and to make robust estimation of model parameters.

## Acknowledgement

The author is grateful to Zhiyong Duan at JPMorgan Chase for valuable suggestions.

## References

- [1] E. Altman (2006): Default Recovery Rates and LGD in the Credit Risk Model and Practice: An Updated Review of the Literature and Empirical Evidence. *defaultrisk.com*.
- [2] L. Andersen and J. Sidenius (2004): Extension to the Gaussian Copula: Random Recovery and Random Factor Loading. *Journal of Credit Risk* **1**(1), pp. 29-70.
- [3] M. Barco (2007): Going Downturn. *Risk* **20**(8), pp. 70-75.
- [4] Basel Committee on Banking Supervision (2004): Background Note on LGD Quantification. *Bank of International Settlements*.
- [5] Basel Committee on Banking Supervision (2005): Guidance on Paragraph 468 of the Framework Document. *Bank of International Settlements*.
- [6] N. Bennani and J. Maetz (2009): A Spot Recovery Rate Extension of the Gaussian Copula. *defaultrisk.com*.
- [7] A.Chabaane, J-P. Laurent and J. Salomon (2004): Double Impact: Credit Risk Assessment and Collateral Value. *Revue Finance*, **25**, pp. 157-178.
- [8] K. Düllmann and M. Trapp (2004): Systematic risk in recovery rates – an empirical analysis of US corporate credit exposures. Deutsche Bundesbank discussion paper 2.

- [9] J. Frye (2000): Collateral Damage. *Risk* **13**(4), pp. 91-94.
- [10] G. Giese (2005): The Impact of PD/LGD correlation on Credit Risk Capital. *Risk*, **18**(4), pp. 78-84.
- [11] M. Hillebrand (2006): Modeling and Estimating Dependent Loss Given Default. *Risk* **19**(9), pp. 120-125.
- [12] D. Li (2000): On Default Correlation: A Copula Function Approach. Working Paper, Nr. 99-07, RiskMetrics.
- [13] H. Li (2009): On Models of Stochastic Recovery for Base Correlation. <http://mpira.ub.uni-muenchen.de/17894>
- [14] H. Li (2009): Extension of Spot Recovery Model for Gaussian Copula. [defaultrisk.com](http://defaultrisk.com).
- [15] H. Li (2010): Double Impact on CVA for CDS: Wrong-Way Risk with Stochastic Recovery. [defaultrisk.com](http://defaultrisk.com)
- [16] P. Miu and B. Ozdemir (2006): Basel Requirement of Downturn LGD: Modeling and Estimating PD and LGD Correlations. *Journal of Credit Risk*, **2**(2), pp. 43-68.
- [17] M. Pykhtin (2003): Unexpected Recovery Risk. *Risk* **16**(8), pp. 74-78.
- [18] D. Rösch and H. Scheule (2005): A Multi-Factor Approach for Systematic Default and Recovery Risk. *Journal of Fixed Income*, **15**(2), pp. 63-75.
- [19] D. Tasche (2004): The single risk factor approach to capital charges in case of correlated loss given default rates. Working Paper, Deutsche Bundesbank.
- [20] O. Vasicek (2002): Loan Portfolio Value. *Risk* **15**(12), pp. 160-162.
- [21] J. Witzany (2009): Loss, Default, and Loss Given Default Modeling. Working Paper, IES.