



Munich Personal RePEc Archive

Testing day-of-the-week persistence and seasonality in Spanish Electricity Energy prices

Yaya, OlaOluwa S. and Vo, Xuan Vinh

Economic and Financial Statistics Unit, Department of Statistics,
University of Ibadan, Ibadan, Nigeria, Institute of Business
Research, University of Economics, Ho Chi Minh City, Viet Nam

15 December 2021

Online at <https://mpra.ub.uni-muenchen.de/112652/>
MPRA Paper No. 112652, posted 06 Apr 2022 13:49 UTC

Testing day-of-the-week persistence and seasonality in Spanish Electricity Energy prices

OlaOluwa S. Yaya¹ and Xuan V. V²

ABSTRACT

Day-of-the-week persistence and seasonality of electricity prices in Spain, spanning 01/01/2006 to 04/11/2021, are investigated by employing updated fractional persistence frameworks in nonlinear settings. The results show marginal higher persistence in electricity prices during the working days (Tuesday, Wednesday, Thursday and Friday), compared to weekend days. In all cases, electricity prices are mean-reverting with long-range dependence properties. Results also show that the monthly electricity price series contain no seasonal effect.

Keywords: *Electricity prices; Fractional persistence; Unit root; Day-of-the-week effect; seasonality; Spain.*

JEL Classifications: C22; Q47.

I. Introduction

In this paper, we investigate the day-of-the-week price persistence and seasonality in Spanish energy prices using an updated fractional persistence framework. We employ two recent nonlinear approaches, i.e. nonlinear I(d) frameworks recently proposed. The first is the Cuestas and Gil-Alana (2016) [CG2016] nonlinear I(d) method which is based on Chebyshev polynomial in time, while the second is the Gil-Alana and Yaya (2021) [GY2021] method that relies on Fourier-form nonlinearity, a mimic of Enders and Lee (2012), in the fractional persistence framework. These two nonlinear functions allow for the smooth modelling of structural breaks in the time series. Bai and Perron (2003) multiple break test considers a maximum of five instantaneous breaks to be detected and in reality, breaks could be more than five in a given long time series, so, this test is restrictive, and in reality, some breaks are

¹Economic and Financial Statistics Unit, Department of Statistics & Institute of Business Research, University of Economics Ho Chi Minh City, Ho Chi Minh City, Vietnam. Email: os.yaya@ui.edu.ng

²Institute of Business Research & CFVG, University of Economics Ho Chi Minh City, Ho Chi Minh City, Vietnam

Email address: vinhvx@ueh.edu.vn

smoothly changing with time. Enders and Lee (2012) have found that a model with trigonometric function could have more power than the Bai-Perron break test in this sense in modelling as many multiple breaks in a smooth fashion.

Fractional persistence allows for a given time series (i.e. energy price in this case) to be checked for the possibility of mean reversion or non-mean reversion. Another subset of the mean reversion case is long memory where a certain real value d lies in the interval i.e., $0 < d < 0.5$, where future values closely depend on the past lagged values since the autocorrelation function of the time series decay exponentially slowly, unlike the case of Autoregressive Moving Average (ARMA) process. Thus, this is the case of stationary mean reversion. Generally, mean reversion implies d lying in the range $0 < d < 1$, including both stationary and nonstationary ranges. Thus, once the external shock is triggered to the time series process, there is the tendency for the series to revert to its mean level after a short time. The case of non-mean reversion is $(1 \leq d < 2)$, where it is unlikely for the series to revert to its mean level, even with as strong government and regulator policies, and if at all, it will take a very long time. These are the appealing properties of the fractional persistence framework, which makes it distinguishable among classical unit root tests in the literature. Meanwhile, the Dickey-Fuller-like unit root test has been found to lack powers in the presence of fractional unit root alternatives (Lee and Schmidt, 1996).

Energy pricing and its distribution, as it affects households and businesses is one of the major global discussions. The present work is gingered by the alarmingly rising electricity prices in Spain since July 2021 due to the global gas crisis. Findings here will therefore guide regulators and the government on the price dynamics of electricity in Spain.

II. Nonlinear Fractional $I(d)$ frameworks

Following Granger and Joyeux (1980) and Hosking (1981), we define fractional integration in time series as,

$$(1-L)^d x_t = \tilde{\varepsilon}_t \quad (1)$$

where L is the lag operator, i.e., $L^q y_t = y_{t-q}$, d is the fractional integration parameter operating on the differencing operator $(1-L)^d$ on the time series y_t to produce a stable/stationary series ε_t that has the properties of white noise. In some cases of highly autocorrelated series, one can consider the possibility of Autoregressive [AR(1)] error disturbance, i.e.

$$\tilde{\varepsilon}_t = \theta_1 \tilde{\varepsilon}_{t-1} + v_t \quad (2)$$

where θ_1 is the AR(1) parameter for $\tilde{\varepsilon}_{t-1}$ which is the first lag of $\tilde{\varepsilon}_t$, and v_t is some noise process. Much recently, the applicability of the operator d (1) has gone beyond unit root testing of time series to testing persistence of series in terms of mean or non-mean reversion. By extending the above to using the deterministic term of a linear trend with constant, α and intercept, β , one obtains the Robinson (1994) fractional persistence modelling framework,

$$y_t = \alpha + \beta t + x_t, \quad (1-L)^d x_t = \tilde{\varepsilon}_t, \quad t = 1, 2, \dots, \quad (3)$$

which is estimated using the Ordinary Least Square (OLS) estimation method, i.e.,

$$\tilde{\varepsilon}_t = y_t^* - \hat{\alpha}_0 1_t^* + \hat{\beta} t_t^*, \quad (4)$$

and $y_t^* = (1-L)^{d_0} y_t$; $1_t^* = (1-L)^{d_0} 1_t$; $t_t^* = (1-L)^{d_0} t_t$, with the complex Lagrange Multiplier test statistic,

$$\hat{R} = \frac{T}{\hat{\sigma}^4} \hat{a}' \hat{A}^{-1} \hat{a}, \quad (5)$$

where T is the sample size, and

$$\hat{\mathbf{a}} = \frac{-2\pi}{T} \sum_j^* \psi(\lambda_j) \mathbf{g}_u(\lambda_j; \hat{\tau})^{-1} \mathbf{I}(\lambda_j); \quad \hat{\sigma}^2 = \sigma^2(\hat{\tau}) = \frac{2\pi}{T} \sum_{j=1}^{T-1} \mathbf{g}_u(\lambda_j; \hat{\tau})^{-1} \mathbf{I}(\lambda_j),$$

$$\hat{\mathbf{A}} = \frac{2}{T} \left\{ \sum_j^* \psi(\lambda_j) \psi(\lambda_j)' - \sum_j^* \psi(\lambda_j) \hat{\xi}(\lambda_j)' \left[\sum_j^* \hat{\xi}(\lambda_j) \hat{\xi}(\lambda_j)' \right]^{-1} \sum_j^* \hat{\xi}(\lambda_j) \psi(\lambda_j)' \right\};$$

$$\psi(\lambda_j) = \log \left| 2 \sin \frac{\lambda_j}{2} \right|; \quad \hat{\xi}(\lambda_j) = \frac{\partial}{\partial \tau} \log g_\varepsilon(\lambda_j; \hat{\tau}), \quad \lambda_j = 2\pi j/T,$$

and * implies that the sums are taken over all frequencies that are bounded in the spectrum, with periodogram $\mathbf{I}(\lambda_j)$ for $\tilde{\varepsilon}_t$ and $\hat{\tau} = \arg \min_{\tau \in T^*} \sigma^2(\tau)$.

The linear I(d) framework of Robinson (1994) is extended to the nonlinear I(d) framework in Cuestas and Gil-Alana (2016) by using the function $f(t)$ which depends on unknown parameters θ_i ($i=1,2,\dots,m$) in the I(d) framework,

$$\begin{aligned} (1-L)^d y_t &= \alpha + f(t) + \tilde{\varepsilon}_t \\ &\cong \theta_0 + \sum_{i=1}^m \theta_i \mathbf{P}_{i,T}(t) + \tilde{\varepsilon}_t, \quad t=1,2,\dots, i=1,\dots,m \end{aligned} \quad (6)$$

where θ_0 is the constant and θ_i are the parameters of the m^{th} order expansion of the Chebyshev polynomial

$$\mathbf{P}_{i,T}(t) = \sqrt{2} \cos \left[i\pi(t-0.5)/T \right], \quad t=1,2,\dots, i=1,\dots,m \quad (7)$$

and $\pi = 3.142$. Then, in the absence of nonlinearity, the I(d) model in (6) is expressed with only a constant, if $m=1$, the model becomes a linear I(d) model of Robinson (1994) with a linear trend. For $m > 1$, the model becomes nonlinear such that the higher the order of m , the higher the nonlinearity. We can re-express the model in (7), for a truncation order of $m=3$. For simplicity of OLS estimation, (y7) in (6) is expressed as,

$$\tilde{\varepsilon}_t = y_t^* - \alpha 1_t^* - \cos 1^* - \cos 2^* - \cos 3^* \quad (8)$$

where $y_t^* = (1-L)^{d_0} y_t$, $1_t^* = (1-L)^{d_0} 1_t$, $\cos 1^* = (1-L)^{d_0} \sqrt{2} \cos[\pi(t-0.5)/T]$,
 $\cos 2^* = (1-L)^{d_0} \sqrt{2} \cos[2\pi(t-0.5)/T]$ and $\cos 3^* = (1-L)^{d_0} \sqrt{2} \cos[3\pi(t-0.5)/T]$.

As a follow-up to Cuestas and Gil-Alana (2016) method, Gil-Alana and Yaya (2021) considered using flexible Fourier fractional (FFF) form $f(t) = \sum_k^n \lambda_k \sin(2\pi j_k t/T) + \sum_k^n \gamma_k \cos(2\pi j_k t/T)$ in the I(d) structure in (6) to mimic the nonlinearity induced by function $f(t)$, where n is the optimal FFF value, $n = j_k$; k a particular frequency which could be fractional or unit; λ_k and γ_k determine the amplitude and displacement of the sine/cosine in the Fourier function. Thus, the nonlinear I(d) is,

$$(1-L)^d y_t = \alpha + \left\{ \sum_k^n \lambda_k \sin(2\pi j_k t/T) + \sum_k^n \gamma_k \cos(2\pi j_k t/T) \right\} + \tilde{\varepsilon}_t \quad (9)$$

Empirically, (9) can be re-expressed as,

$$\tilde{\varepsilon}_t = y_t^* - \alpha 1_t^* - \sum_k^n \lambda_k \sin_{k,t}^* - \sum_k^n \gamma_k \cos_{k,t}^*, \quad (10)$$

where $\sin_{k,t}^* = (1-L)^{d_0} \sin(2\pi j_k t/T)$, $\cos_{k,t}^* = (1-L)^{d_0} \cos(2\pi j_k t/T)$ with y_t^* and 1_t^* defined in (7) above.

III. Data and Results

Daily historical spot prices of electricity for Spain, spanning 01/01/2006 to 04/11/2021, covering a total of 5788 data points, are obtained from the website of Iberian Energy Market Operator/ Operador del Mercado Ibérico de la Energía-Polo Portugués (OMIP) at <https://www.omip.pt/en/electricity>. Plots of daily and monthly electricity prices are given in Figure 1. For daily prices, there are fluctuations over time and occasional price swings that are pronounced to both up and down prices. Between 2006 and June 2021, prices hover between

50 and 150 euro/MWh while prices started increasing astronomically in the middle of 2021³. The average monthly pricing is superimposed on the daily price, and the essence of this is to check for possible monthly seasonal movement. By looking at the plot, the monthly seasonality is not observed.

Figure 1: Plot of electricity prices in Spain based on daily energy auctions




Table 1 presents the results of persistence using nonlinear smooth trend functions by employing the Chebyshev polynomial in time and Fourier function. In the case of the Chebyshev polynomial, the nonlinear trigonometric parameters, \cos_1 , \cos_2 and \cos_3 are not significant, while in the case of Fourier function, the trigonometric parameter \sin_k is only significant with the optimal Fourier frequency $k = 0.46$. For both the Chebyshev polynomial and Fourier function, fractional persistence values are 0.6739 and 0.6702, respectively, under white noise disturbance and each estimate with an upper CI bound of about 0.69. To check for the possible interference of unchecked serial correlations in the model residuals which could bias our results, we allow for an AR(1) error disturbance instead of the white noise error

³ <https://www.euronews.com/2021/10/28/why-europe-s-energy-prices-are-soaring-and-could-get-much-worse>

disturbance, with the specification given in (2). The results for both nonlinear I(d) are then re-presented in the lower panel of Table 1. In this case, cos2 parameter in the Chebyshev function is found to be significant, while \sin_k parameter in the Fourier function is also significant. For both nonlinear functions, the AR(1) parameter estimates are low, around 0.20, but this is significant at 5% level. The persistence values estimated for both nonlinear I(d) models are found to be lower compared to the corresponding values under the white noise error disturbances. These values are found to be 0.5601 and 0.5548, respectively, which is quite around 0.5 implying mean reversion with long-range dependence characteristics in the electricity prices.

Table 1: Results based on nonlinear I(d) frameworks

In bold are significant parameters at a 5% level of significance for one-sided t-test, while values of t-statistic are in parenthesis.

a. White noise error disturbance			
Chebyshev polynomial in time		Fourier function	
<i>d</i>	0.6739 (0.6519, 0.6960)	<i>d</i>	0.6702 (0.6484, 0.6920)
<i>c</i>	4.3549 (0.202)	<i>c</i>	55.6407 (2.91)
cos1	1.9496 (0.159)	\sin_k	-71.3371 (-2.68)
cos 2	10.7374 (1.24)	\cos_k	-20.7772 (-1.21)
cos 3	3.5145 (0.522)	<i>k</i>	0.46
b. AR(1) error disturbance			
Chebyshev polynomial in time		Fourier function	
<i>d</i>	0.5601 (0.5327, 0.5875)	<i>d</i>	0.5548 (0.5236, 0.5822)
<i>c</i>	5.4170 (0.500)	<i>c</i>	35.5051 (11.2)
cos1	-1.1613 (-0.182)	\sin_k	-49.4753 (-3.56)

cos 2	11.7063 (2.39)	cos _k	-10.6573 (-1.26)
cos 3	-2.3742 (-0.598)	k	0.50
AR(1)	0.2084 (11.0)	AR(1)	0.2118 (39.7)

The analysis is extended by obtaining the day-of-the-week series persistence using the two nonlinear I(d) frameworks, and the results, as presented in Table 2 are only reported for the case of AR(1) error disturbance. In the upper panel of the results table for the case of Chebyshev polynomial I(d), it is found that none of the nonlinear parameters for the days-of-the-week series (i.e. Monday – Friday) is significant while for Wednesday, Saturday and Sunday series, nonlinear parameters are significant. The AR(1) parameters are significant for Wednesday, Saturday and Sunday series. The persistence estimates for the seven series are 0.65 (Monday), 0.73 (Tuesday), 0.78 (Wednesday), Friday (0.77), -0.30 (Saturday) and -0.39 (Sunday). By looking at the results in the lower panel of the table for the Fourier function, with optimal Fourier frequency k recorded in the last column of the table, the significance of Fourier form parameters (sine and cosine) are observed. Thus, the results in Table 2 confirm the nonlinear nature of day-of-the-week in Spanish electricity prices. The persistence results, though, all in mean reverting ranges, are in the interval 0.62-0.77. The persistence on Monday, Saturday and Sunday are 0.64, 0.69 and 0.64, respectively while others (Tuesday - Friday) are in the range of 0.71-0.77.

Table 2: Results of persistence for each day of the week with Chebyshev polynomial in time

In bold are significant parameters at a 5% level of significance for one-sided t-test, while values of t-statistic are in parenthesis.

Chebyshev polynomial in time						
Day	d	AR(1)	c	cos 1	cos 2	cos 3
Monday	0.6563 (0.5716, 0.7410)	0.0180 (0.299)	18.8074 (0.884)	-1.4340 (-0.123)	8.2163 (0.996)	-1.5559 (-0.243)

Tuesday	0.7307 (0.6535, 0.8079)	-1.24 (0.217)	24.4815 (0.867)	-2.7505 (-0.180)	8.4563 (0.833)	-2.5285 (-0.331)
Wednesday	0.7833 (0.7114, 0.8552)	-0.1832 (-3.77)	25.6113 (0.710)	-5.3899 (-0.280)	8.2595 (0.684)	-3.3157 (-0.373)
Thursday	---	---	---	---	---	---
Friday	0.7784 (0.6990, 0.8578)	-0.0560 (-1.07)	28.4877 (0.456)	-2.9122 (-0.144)	8.6224 (0.670)	-3.2657 (-0.346)
Saturday	-0.3093 (-0.3644, -0.2542)	0.9919 (0.158.0)	6.0728 (1.01)	-2.5571 (-0.444)	13.5251 (2.45)	-1.0626 (-0.195)
Sunday	-0.3911 (-0.4474, -0.3348)	0.9872 (124.0)	2.1566 (0.853)	-2.8419 (-0.853)	8.1349 (2.29)	-1.9550 (-0.581)
Fourier function						
Day	d	AR(1)	c	\sin_k	\cos_k	k
Monday	0.6414 (0.5567, 0.7260)	0.0254 (0.421)	96.5035 (2.87)	-90.5852 (-2.30)	-62.3713 (-2.28)	0.34
Tuesday	0.7165 (0.6395, 0.7935)	-0.0574 (-1.09)	74.6100 (2.57)	-86.3709 (-2.35)	-36.3699 (-1.59)	0.42
Wednesday	0.7687 (0.6964, 0.8410)	-0.1760 (-3.60)	115.468 (2.59)	-115.820 (-2.13)	-79.8170 (-2.20)	0.36
Thursday	0.7713 (0.6966, 0.8459)	-0.1235 (-2.49)	133.444 (2.68)	-128.972 (-2.17)	-93.7717 (-2.27)	0.35
Friday	0.7649 (0.6857, 0.8441)	-0.0484 (-0.922)	77.1158 (2.19)	-95.1085 (-2.23)	-34.8309 (-1.24)	0.44
Saturday	0.6902 (0.6119, 0.7684)	-0.0339 (-0.627)	69.6715 (2.09)	-75.1435 (-1.78)	-54.8357 (-2.00)	0.37
Sunday	0.6282 (0.5506, 0.7058)	-0.0668 (-1.21)	30.1621 (1.75)	-43.4384 (-2.08)	-4.5923 (-0.336)	0.51

To probe further for possible monthly seasonality, the monthly dataset is estimated by employing a seasonal AR model in the nonlinear I(d) frameworks. Thus, we re-specify (2) as,

$$\tilde{\varepsilon}_t = \theta_{12} \tilde{\varepsilon}_{t-12} + v_t \quad (11)$$

where θ_{12} is the seasonal AR parameter in the case of monthly time series, and $\tilde{\varepsilon}_{t-12}$ is the lag 12 of the resulting fractionally differenced series. Thus, θ_{12} determines the monthly seasonal autoregression between the previous 12th observations (i.e $\tilde{\varepsilon}_{t-12}$) and the current observations, $\tilde{\varepsilon}_t$. The model in (11) is then analyzed with (8) and (10) to obtain results for the Chebyshev polynomial and Fourier function, respectively. The results of the Chebyshev polynomial

reported in the upper panel of the results table indicated non-significance of seasonal AR parameter with a very low coefficient (0.0816). Meanwhile, the fractional persistence estimate is computed as 0.9323 which is quite higher than that obtained for the case of daily pricing. Similar results are obtained in the case of the Fourier function presented in the last panel of Table 3, where the non-significance of the seasonal AR parameter with nonlinearity detected as \cos_k is found to be significant at the 5% level.

Table 3: Results based on Seasonal AR(1) models

a. Nonlinear Chebyshev polynomial with Seasonal AR			
d	0.9323 (-0.5395, 2.4041)	1.24	0.216
c	-0.4151	-0.000	1.000
cos 1	-0.0632	-0.0922	0.927
cos 2	-0.0028	-0.0041	0.997
cos 3	0.1038	0.149	0.882
Seasonal AR	0.0816	0.166	0.868
b. Nonlinear/Fourier function with Seasonal AR			
d	0.9118 (0.7824, 1.0412)	13.8	0.000
c	34296.5	2.50	0.013
\sin_k	0.0582	0.050	0.960
\cos_k	-34238.0	-2.42	0.016
k	0.41		
Seasonal AR	0.0923	0.723	0.470

IV. Concluding remarks

Energy pricing and its distribution, as it affects households and businesses is one of the major global discussions. The present work is gingered by the alarmingly rising electricity prices in Spain since July 2021 due to the global gas crisis, thus the finding in the work guide regulators and the government to know the price dynamics of electricity in Spain. We employed updated fractional persistence frameworks in nonlinear settings in investigating day-of-the-week affect persistence and seasonality in the pricing from 01.01.2006 to 04.11.2021. Results obtained

indicated mean reversion in prices across all days of the week and there is a fair increase in persistence during the working days (Tuesday, Wednesday, Thursday and Friday) compared to weekend days. No evidence of seasonality is found in the monthly price series. It is hoped that the results obtained here will guide in making energy usage policy in Spain.

Acknowledgements:

Comments of the Editor and reviewers are gratefully acknowledged.

References

- Cuestas, J. C. & Gil-Alana, L. A. (2016). A nonlinear approach with long range dependence based on Chebyshev polynomials. *Studies in Nonlinear Dynamics and Econometrics*, 20(1), 57-94.
- Enders, W. and Lee, J. (2012). The flexible Fourier form and Dickey-Fuller-type unit root tests. *Economic Letters*, 117, pp. 196–199.
- Gil-Alana, L. A. and Yaya, O. S. (2021). Testing Fractional Unit Roots with Non-linear Smooth Break Approximations using Fourier functions. *Journal of Applied Statistics*, 48 (13-15), 2542-2559.
- Granger, C. W. J. & Joyeux, R. (1980). An introduction to long memory time series and fractional differencing. *Journal of Time Series Analysis*, 1(1), 15-29.
- Hosking, J. R. M. (1981). Fractional Differencing. *Biometrika*, 68(1), 168-176.
- Lee, D. and Schmidt, P. (1996). On the power of the KPSS test of stationary against fractionally integrated alternatives. *Journal of Econometrics*, 73: 285–302.
- Robinson, P.M. (1994). Efficient Tests of Nonstationary Hypotheses. *Journal of the American Statistical Association* 89: 1420–1437.