



Munich Personal RePEc Archive

Acyclicity of improvements in finite game forms

Kukushkin, Nikolai S.

Russian Academy of Sciences, Dorodnicyn Computing Center

27 November 2008

Online at <https://mpra.ub.uni-muenchen.de/11802/>
MPRA Paper No. 11802, posted 28 Nov 2008 01:16 UTC

Acyclicity of improvements in finite game forms*

Nikolai S. Kukushkin[†]

November 27, 2008

Abstract

Game forms are studied where the acyclicity, in a stronger or weaker sense, of (coalition or individual) improvements is ensured in all derivative games. In every game form generated by an “ordered voting” procedure, individual improvements converge to Nash equilibria if the players restrict themselves to “minimal” strategy changes. A complete description of game forms where all coalition improvement paths lead to strong equilibria is obtained: they are either dictatorial, or voting (or rather lobbying) about two outcomes. The restriction to minimal strategy changes ensures the convergence of coalition improvements to strong equilibria in every game form generated by a “voting by veto” procedure. *JEL Classification Number: C 72.*

Key words: Improvement dynamics; Game form; Perfect information game; Potential game; Voting by veto

*Financial support from a Presidential Grant for the State Support of the Leading Scientific Schools (NSH-2982.2008.1), the Russian Foundation for Basic Research (project 08-07-00158), the Russian Foundation for Humanities (project 08-02-00347), and the Spanish Ministry of Education and Science (project SEJ 2007-67135) is acknowledged. I have benefitted from discussions with Vladimir Gurvich, Hervé Moulin, and Endre Boros, especially during my visit to U.S. universities. I thank Hervé Moulin and Francisco Marhuenda, whose effort and generosity made the trip possible.

[†]Russian Academy of Sciences, Dorodnicyn Computing Center, 40, Vavilova, Moscow 119333 Russia
E-mail: ququns@inbox.ru

1 Introduction

A.-A. Cournot considered best response dynamics long before the expression “game theory” came into use. Such processes were studied in various contexts since then (Topkis, 1979, 1998; Moulin, 1984; Vives, 1990; Milgrom and Roberts, 1991; Kandori and Rob, 1995). Monderer and Shapley (1996) started a similar approach to *better* reply dynamics.

This paper continues the search for natural classes of strategic games where the acyclicity, in a stronger or weaker sense, of (coalition or individual) improvements is ensured (Rosenthal, 1973; Germeier and Vatel, 1974; Sela, 1992; Monderer and Shapley, 1996; Milchtaich, 1996; Holzman and Law-yone, 1997; Konishi et al., 1997; Kukushkin, 1999, 2000, 2002ab, 2004ab, 2006, 2007bc; Friedman and Mezzetti, 2001; Kukushkin et al., 2005). Unlike most of the previous literature, we only consider *game forms*, i.e., we put no restrictions on the preferences of the players, only on strategic interactions.

The topic is somewhat related to the study of consistent, or solvable, game forms (Gurvich, 1975, 1988; Moulin, 1976; Peleg, 1978; Abdou 1995, 1998; Abdou and Keiding, 2003; Boros et al., 2007); however, the acyclicity of improvements is a much stronger (and rarer) property than just the existence of an equilibrium.

In the case of two players, a kind of complete description of game forms where all individual improvement paths in all derivative games lead to Nash equilibria was obtained by Boros et al. (2008ab); for more than two players, there is no clear prospect for that. Weaker notions of acyclicity of individual improvements result in wider classes of game forms, also without clear prospects for a characterization. The most interesting and important class of game forms with acyclic individual improvements is that of games with perfect information (Kukushkin, 2002a); the results of that paper are somewhat extended here. Similar properties of “ordered voting game forms” (Proposition 3.4 and Theorem 3.6) are also established; such game forms have been considered before (e.g., Moulin, 1980, Kukushkin, 1995, or Mariotti, 2000), but improvement dynamics in them seem to have never been studied.

Concerning coalition improvements, a complete description of game forms where all such improvement paths in all derivative games lead to strong equilibria *is* obtained; not surprisingly, there are not so many of them (Theorem 4.2). A slight weakening of the requirement widens the class significantly. It is shown that “voting by veto” procedures (Mueller, 1978; Peleg, 1978) generate game forms where the convergence of coalition improvements to strong equilibria is ensured if the players restrict themselves to “minimal” strategy changes (Theorem 4.10).

In Section 2 the basic definitions concerning improvement dynamics in finite strategic games are given; the notion of a game form is introduced and examples of game forms ensuring the

acyclicity of improvements are provided. Section 3 contains some general results about game forms with acyclic individual improvements; the class of “ordered voting game forms” is defined. Subsection 3.2 is about games with perfect information; Subsection 3.3, about exact potential, a cardinal analog of acyclicity. Section 4 contains the characterization of game forms with acyclic coalition improvements; in Subsection 4.2, voting by veto procedures are defined and their interesting properties proven.

2 Basic Notions

2.1 Improvement paths in strategic games

Our basic model is a *finite strategic game with ordinal preferences*. It is defined by a finite set of players N (we denote $n = \#N$), and finite strategy sets X_i and *ordinal utility functions* $u_i: X_N \rightarrow \mathbb{R}$, where $X_N = \prod_{i \in N} X_i$, for all $i \in N$. We denote $\mathcal{N} = 2^N \setminus \{\emptyset\}$ (the set of potential coalitions) and $X_I = \prod_{i \in I} X_i$ for each $I \in \mathcal{N}$; instead of $X_{N \setminus \{i\}}$ and $X_{N \setminus I}$, we write X_{-i} and X_{-I} , respectively. If $n = 2$, then $-i$ refers to the partner of player i .

Remark. Whenever $v_i: \mathbb{R} \rightarrow \mathbb{R}$ is strictly increasing, the functions u_i and $v_i \circ u_i$ represent the same ordering. Therefore, any meaningful definition, condition, statement, etc., involving ordinal utility functions must be invariant to strictly increasing transformations. Auxiliary constructions, however, may well use numeric values.

With every strategic game, a number of improvement relations on X_N are associated ($i \in N$, $I \in \mathcal{N}$, $y_N, x_N \in X_N$):

$$y_N \triangleright^{\text{Ind}_i} x_N \Leftrightarrow [y_{-i} = x_{-i} \ \& \ u_i(y_N) > u_i(x_N)]; \quad (2.1a)$$

$$y_N \triangleright^{\text{Ind}} x_N \Leftrightarrow \exists i \in N [y_N \triangleright^{\text{Ind}_i} x_N] \quad (2.1b)$$

(*individual improvement relation*);

$$y_N \triangleright^{\text{sCo}_I} x_N \Leftrightarrow [y_{-I} = x_{-I} \ \& \ \forall i \in I [u_i(y_N) > u_i(x_N)]]; \quad (2.2a)$$

$$y_N \triangleright^{\text{sCo}} x_N \Leftrightarrow \exists I \in \mathcal{N} [y_N \triangleright^{\text{sCo}_I} x_N] \quad (2.2b)$$

(*strong coalition improvement relation*);

$$y_N \triangleright^{\text{wCo}_I} x_N \Leftrightarrow [y_{-I} = x_{-I} \ \& \ \forall i \in I [u_i(y_N) \geq u_i(x_N)] \ \& \ \exists i \in I [u_i(y_N) > u_i(x_N)]]; \quad (2.3a)$$

$$y_N \triangleright^{\text{wCo}} x_N \Leftrightarrow \exists I \in \mathcal{N} [y_N \triangleright^{\text{wCo}_I} x_N] \quad (2.3b)$$

(*weak coalition improvement relation*).

Defining the *best response correspondence* $\mathcal{R}_i: X_{-i} \rightarrow 2^{X_i}$ for each $i \in N$ in the usual way,

$$\mathcal{R}_i(x_{-i}) = \underset{x_i \in X_i}{\text{Argmax}} u_i(x_i, x_{-i})$$

for every $x_{-i} \in X_{-i}$, we may introduce one more relation:

$$y_N \triangleright^{\text{BR}}_i x_N \Leftrightarrow [y_{-i} = x_{-i} \ \& \ x_i \notin \mathcal{R}_i(x_{-i}) \ni y_i]; \quad (2.4a)$$

$$y_N \triangleright^{\text{BR}} x_N \Leftrightarrow \exists i \in N [y_N \triangleright^{\text{BR}}_i x_N] \quad (2.4b)$$

(*best response improvement* relation).

It is often convenient to speak of just “an improvement relation” \triangleright without specifying which of the above-defined relations is meant. A *maximizer* of an improvement relation \triangleright , i.e., a strategy profile $x_N \in X_N$ such that $y_N \triangleright x_N$ holds for no $y_N \in X_N$, is an equilibrium: a Nash equilibrium if \triangleright is $\triangleright^{\text{Ind}}$; a (“very”) strong equilibrium if \triangleright is $\triangleright^{\text{sCo}}$ ($\triangleright^{\text{wCo}}$). Every Nash equilibrium is a maximizer of $\triangleright^{\text{BR}}$. If $\mathcal{R}_i(x_{-i}) \neq \emptyset$ for all $i \in N$ and $x_{-i} \in X_{-i}$, then the converse statement is also true; in a finite game, the condition holds automatically.

Following Kukushkin (2004a), we consider an arbitrary binary relation \triangleright on a finite set X . An *improvement path* (for \triangleright) is a (finite or infinite) sequence $\{x^k\}_{k=0,1,\dots}$ such that $x^{k+1} \triangleright x^k$ whenever $k \geq 0$ and x^{k+1} is defined. A *finite improvement cycle* is an improvement path $x_N^0, x_N^1, \dots, x_N^m = x_N^0$ ($m > 0$); a relation is *acyclic* if it admits no finite improvement cycle. On a finite set, that property is equivalent to the impossibility of an infinite improvement path; therefore, every improvement path, if continued whenever possible, reaches a maximizer (equilibrium) in a finite number of steps.

It is easy to see that a binary relation \triangleright on a finite set X is acyclic if and only if it admits a *numeric potential*, i.e., a function $P: X \rightarrow \mathbb{R}$ such that

$$\forall y, x \in X [y \triangleright x \Rightarrow P(y) > P(x)]; \quad (2.5a)$$

the property is also equivalent to the existence of an *order potential*, i.e., an irreflexive and transitive binary relation \succ on X such that

$$\forall y, x \in X [y \triangleright x \Rightarrow y \succ x]. \quad (2.5b)$$

The relation \triangleright is *weakly acyclic* if every $x \in X$ is connected to a maximizer of \triangleright with an improvement path, i.e., there is a finite improvement path $\{x^0, \dots, x^m\}$ ($m \geq 0$) such that $x^0 = x$ and x^m is a maximizer. The weak acyclicity does not exclude the possibility that an improvement process may continue indefinitely without reaching an equilibrium; however, this is improbable under reasonable assumptions (Kalai and Schmeidler, 1977; Milchtaich, 1996; Friedman and Mezzetti, 2001).

Clearly, acyclicity implies weak acyclicity, which, in turn, implies the existence of a maximizer of \triangleright . Neither statement can be reversed.

An essential feature of the improvement relations in a strategic game defined by (2.1)–(2.4) is their *disjunctive structure*, reflected in (2.1b), etc. It allows us to introduce an intermediate class of properties. We consider an abstract relation \triangleright with a disjunctive structure, i.e., assume that there are a finite set M (in strategic games, $M = N$ for individual improvements and $M = \mathcal{N}$ for coalition improvements) and binary relations \triangleright_i on X for each $i \in M$ such that $y \triangleright x \iff \exists i \in M [y \triangleright_i x]$ for all $y, x \in X$. We say that \triangleright is *acyclic under restrictions* if there are binary relations $\triangleright\triangleright$ and $\triangleright\triangleright_i$ on X such that, for all $i \in M$ and $y, x \in X$, there holds

$$y \triangleright\triangleright_i x \Rightarrow y \triangleright_i x; \quad (2.6a)$$

$$\exists y \in X [y \triangleright_i x] \Rightarrow \exists z \in X [z \triangleright\triangleright_i x]; \quad (2.6b)$$

$$y \triangleright\triangleright x \iff \exists i \in M [y \triangleright\triangleright_i x]; \quad (2.6c)$$

$$\triangleright\triangleright \text{ is acyclic.} \quad (2.6d)$$

If \triangleright is acyclic, then all conditions (2.6) are satisfied by \triangleright itself as $\triangleright\triangleright$, i.e., \triangleright is acyclic under restrictions. In any case, the conditions (2.6a) and (2.6b) imply that every improvement path of $\triangleright\triangleright$ is an improvement path of \triangleright and both relations have the same maximizers. It follows immediately that a relation acyclic under restrictions is weakly acyclic. When \triangleright is acyclic under restrictions, an order potential of $\triangleright\triangleright$ may be called a *restricted order potential* of \triangleright [cf. Proposition 6.4 of Kukushkin (2004a)].

Restricted acyclicity means that it is possible to impose restrictions on each player's (or coalition's) strategy changes so that whenever an improvement is possible, an admissible improvement is possible as well, and the convergence to an equilibrium is ensured. When an improvement relation is only weakly acyclic, an agreement between the players as to who is allowed to improve at each stage may be needed. (Explicit cooperation can be replaced with a stochastic choice of the player or coalition to move at each step.) The example in Section 7.7 of Kukushkin (2004a) clarifies the difference between the weak acyclicity and acyclicity under restrictions.

The (weak or restricted) acyclicity of the individual improvement relation $\triangleright^{\text{Ind}}$ (2.1) in a finite strategic game Γ is called the *(weak or restricted) finite individual improvement property* ((*weak or restricted*) *FIP*) of Γ . Similarly, the *(weak or restricted) finite coalition improvement property* ((*weak or restricted*) *FCP*) refers to the strong coalition improvement relation $\triangleright^{\text{sCo}}$ defined by (2.2); the *(weak or restricted) FCP⁺*, to the weak coalition improvement relation $\triangleright^{\text{wCo}}$ defined by (2.3); the *(weak or restricted) FBRP*, to the best response improvement relation $\triangleright^{\text{BR}}$ defined

by (2.4). It is easy to see that the following implications hold:

$$\begin{array}{ccccccc}
\text{FCP}^+ & \Rightarrow & \text{FCP} & \Rightarrow & \text{FIP} & \Rightarrow & \text{FBRP} \\
\Downarrow & & \Downarrow & & \Downarrow & & \Downarrow \\
\text{restricted FCP}^+ & \Rightarrow & \text{restricted FCP} & \Rightarrow & \text{restricted FIP} & \Leftarrow & \text{restricted FBRP} \\
\Downarrow & & \Downarrow & & \Downarrow & & \Downarrow \\
\text{weak FCP}^+ & & \text{weak FCP} & & \text{weak FIP} & \Leftarrow & \text{weak FBRP}
\end{array}$$

Actually, the FBRP *is* a restricted FIP.

These properties admit the same dynamic interpretation as the (weak or restricted) acyclicity of an abstract binary relation. The FCP(+), FIP, or FBRP ensure that all appropriate adaptive dynamics converge to an equilibrium in a finite number of steps. The weak FCP(+), weak FIP, or weak FBRP ensure the convergence to an appropriate equilibrium in a finite number of steps with probability one under reasonable assumptions. The properties are also conducive to the convergence of more sophisticated scenarios (Young, 1993; Kandori and Rob, 1995; Friedman and Mezzetti, 2001).

Remark. According to Proposition 6.4 of Kukushkin (2004a), weak and restricted FBRP are equivalent for two person games; unfortunately, neither property seems natural for *game forms*, which are the subject of this paper. The equivalence does not hold w.r.t. the FIP or FCP(+).

2.2 Game forms

A *game form* G is defined by a finite set of players N , a finite strategy set X_i for each $i \in N$, a finite set of outcomes A and a mapping $g: X_N \rightarrow A$, where $X_N = \prod_{i \in N} X_i$ is the set of strategy profiles. For notational simplicity, we assume $X_i \cap X_j = \emptyset$ whenever $i \neq j$; the assumption is obviously innocuous. We denote $\mathcal{X} = \bigcup_{i \in N} X_i$.

Once preferences of the players over the outcomes are specified (and we always assume this to be done with ordinal utilities $v_i: A \rightarrow \mathbb{R}$), a *derivative game* $G(v_N)$ (where v_N denotes a list $\langle v_i \rangle_{i \in N}$) emerges, in which the set of players is N , the strategy sets are X_i 's and utilities are $u_i(x_N) = v_i(g(x_N))$.

If every derivative game $G(v_N)$ possesses a Nash (strong) equilibrium, G is called *Nash (strong) consistent*. We say that G has the FIP, FCP or FCP(+) if so does every derivative game $G(v_N)$. We also use the expression G is an FIP, or FCP(+) *game form*. If G has the FIP (FCP), then G is Nash (strongly) consistent; the converse statements are wrong.

Remark. *FBRP game forms* could be defined quite similarly; however, there is no example of an FBRP game form without the FIP. Moreover, Corollary 2 from Kukushkin (2007a) shows that the FIP and FBRP are equivalent as properties of *two person* game forms.

Example 2.1. Let us consider four game forms with two players:

$$\text{a. } \begin{bmatrix} a & a & a \\ b & b & b \\ c & c & c \end{bmatrix} \quad \text{b. } \begin{bmatrix} a & a & a \\ b & c & c \\ b & d & e \end{bmatrix} \quad \text{c. } \begin{bmatrix} a & a & a \\ a & b & b \\ a & b & c \end{bmatrix} \quad \text{d. } \begin{bmatrix} a & a & c \\ a & b & b \\ c & b & c \end{bmatrix}.$$

It is easily seen that each of them has the FIP, but only the first has FCP (and FCP⁺ at that).

Example 2.2. Let us consider two game forms with two players:

$$\text{a. } \begin{bmatrix} a & a & a \\ a & \underline{b} & \underline{c} \\ a & \underline{d} & \underline{e} \end{bmatrix} \quad \text{b. } \begin{bmatrix} a & a & a \\ a & \underline{b} & \underline{a} \\ a & \underline{a} & \underline{b} \end{bmatrix}.$$

It is easily checked that both game forms are Nash consistent; actually, the northwestern corner is a Nash equilibrium for all utilities. Meanwhile, the underlined strategy profiles form an improvement cycle for appropriate utilities in either game form, hence neither has the FIP. However, there is an important difference between them.

Suppose that the utilities of the players in the first game form satisfy these inequalities: $v_1(b) > v_1(e) > v_1(d) > v_1(c) > v_1(a)$ and $v_2(c) > v_2(d) > v_2(b) > v_2(e) > v_2(a)$. The northwestern corner is a unique Nash equilibrium; an agreement to choose it is self-policing in the usual sense. On the other hand, an agreement *not* to choose equilibrium strategies is self-policing as well: if I believe that my partner honors the agreement, I have no incentive to cheat. Moreover, each player would prefer the second agreement, which ensures the choice of one of the underlined outcomes, to the first, notwithstanding the fact that the resulting outcome remains unpredictable. The “irrelevance of equilibria” of this kind was discussed by Kreps (1990, pp. 416-417). Example 1.3 in Kukushkin (2002b) demonstrates a similar problem concerning strong equilibria.

One could argue that the second agreement is just to choose a mixed equilibrium; note, however, that our players have ordinal preferences, hence they need not be able to compare probability distributions on the set of outcomes. More technically speaking, a mixed equilibrium whose support consists of the underlined strategy profiles exists for every pair of utility functions representing the same preferences, but its probability distribution is not invariant under monotonic transformations.

No such ugly thing may happen when a game has the FIP or FCP⁽⁺⁾; actually, the weak FIP or weak FCP⁽⁺⁾ is sufficient (Theorems 3.1–3.3 of Kukushkin, 2002b). The second game form has the weak FIP, hence is immune to this pathology at least.

Remark. No game form is known such that every derivative game has the weak FIP or weak FCP⁽⁺⁾, but not necessarily a restricted FIP or FCP⁽⁺⁾.

The notion of restricted acyclicity admits a strengthening when applied to game forms. An *admissible change (by coalition $I \in \mathcal{N}$) relation* is a binary relation \vdash_I on X_N such that $x_{-I} = y_{-I}$ whenever $x_N \vdash_I y_N$. When $I = \{i\}$, we use the notation \vdash_i . Such a relation is *liberal* if

$$\forall x_N, y_N \left[[y_{-I} = x_{-I} \ \& \ g(y_N) \neq g(x_N)] \Rightarrow \exists z_N [x_N \vdash_I z_N \ \& \ g(z_N) = g(y_N)] \right], \quad (2.7)$$

i.e., if every change of outcome available to a coalition (or a player) can be done in an admissible way.

A game form G has an *almost unrestricted FIP* if there is a list of admissible change relations $\langle \vdash_i \rangle_{i \in \mathcal{N}}$ such that every \vdash_i is liberal and, in every derivative game $G(v_N)$, the admissible improvement relation \triangleright defined by (2.6c) with $M = N$ and

$$y_N \triangleright_i x_N \Leftrightarrow [x_N \vdash_i y_N \ \& \ y_N \triangleright^{\text{Ind}}_i x_N] \quad (2.8)$$

is acyclic. A game form G has an *almost unrestricted FCP* if there is a list of admissible change relations $\langle \vdash_I \rangle_{I \in \mathcal{N}}$ such that every \vdash_I is liberal and, in every derivative game $G(v_N)$, the admissible improvement relation \triangleright defined by (2.6c) with $M = \mathcal{N}$ and

$$y_N \triangleright_I x_N \Leftrightarrow [x_N \vdash_I y_N \ \& \ y_N \triangleright^{\text{sCo}}_I x_N] \quad (2.9)$$

is acyclic. Naturally, an almost unrestricted FIP (FCP) of a game form ensures the existence of a (strong) Nash equilibrium in every derivative game, but it is much more than that.

Remark. The game form in Example 2.2b has an almost unrestricted FCP although it seems not to be covered by Theorems 3.6 or 3.13 or 4.10 below.

The notion admits a more concrete interpretation. We may assume that there are costs associated with every change of strategy and that a change is admissible if its cost is minimal among all changes leading to the same outcome. To be more formal, we define a *quasidistance* on X_I as a mapping $\delta: X_I \times X_I \rightarrow \mathbb{N} \cup \{+\infty\}$ such that $\delta(x_I, y_I) \leq \delta(x_I, z_I) + \delta(z_I, y_I)$ for all $x_I, y_I, z_I \in X_I$. When an almost unrestricted FIP is concerned, only singleton I are needed, naturally. Given a family of quasidistances on X_I parameterized with $x_{-I} \in X_{-I}$, we define an admissible change relation by

$$x_N \vdash_I y_N \Leftrightarrow [y_{-I} = x_{-I} \ \& \ \nexists z_I \in X_I [g(z_I, x_{-I}) = g(y_N) \ \& \ \delta_{x_{-I}}(x_I, z_I) < \delta_{x_{-I}}(x_I, y_I)]] \quad (2.10)$$

Since X_I is finite, (2.7) holds, i.e., the relation is liberal. Therefore, a list of quasidistances on all X_I (X_i) generates an admissible improvement relation \triangleright , based on (2.10), (2.9) or (2.8), and (2.6c) with $M = \mathcal{N}$ or $M = N$, in every derivative game $G(v_N)$.

Proposition 2.3. *A game form G has an almost unrestricted FCP (FIP) if and only if there is a family of quasidistances $\delta_{x_{-I}}$ ($\delta_{x_{-i}}$) on each X_I (X_i) such that the admissible improvement relation \triangleright generated in every derivative game $G(v_N)$ is acyclic.*

Proof. Sufficiency is tautological. Let G have an almost unrestricted FCP with an admissible improvement relation \triangleright . For each $I \in \mathcal{N}$, we define \vdash_I^* as the transitive closure of \vdash_I , and $\delta_{x_{-I}}(x_I, y_I) = 0$ if $(x_I, x_{-I}) \vdash_I^*(y_I, x_{-I})$ while $\delta_{x_{-I}}(x_I, y_I) = +\infty$ otherwise. Clearly, every $\delta_{x_{-I}}$ is a quasidistance. Now (2.10) with this family $\delta_{x_{-I}}$ defines \vdash_I^* ; in every derivative game, (2.9) defines the admissible improvement relation \triangleright_I^* , which is the transitive closure of \triangleright_I ; therefore, \triangleright^* is acyclic too. The case of an almost unrestricted FIP is treated in the same way. \square

In the following theorems, we always define admissible changes through quasidistances, i.e., “costs,” which look reasonable in each case.

A *fragment* G' of G is a game form with the same set of players N , nonempty subsets $\emptyset \neq X'_i \subseteq X_i$ for all $i \in N$, and the restriction of g to X'_N as g' . If G has the FIP (FCP), then so does every fragment of G ; (strong) Nash consistency, or even a restricted FIP (FCP), need not be “inherited” in this sense.

3 FIP Game Forms

3.1 Simple and ordered voting game forms

A game form G is *separable* if it is possible to define a “pointer” mapping $p: \mathcal{X} \rightarrow A$ such that

$$g(x_N) \in \{p(x_i)\}_{i \in N} \quad (3.1)$$

for every $x_N \in X_N$. In principle, the same g may be described by (3.1) with different mappings p ; moreover, some $p(x_i)$ may be arbitrary. An interpretation should be clear: choosing a strategy x_i , player i as if expresses a wish to see $p(x_i)$ chosen; the mapping g determines whose desire will be fulfilled at each strategy profile.

Remark. The term is due to Vladimir Gurvich (a seminar presentation). Boros et al. (2008b) suggest “assignable game forms”; however, the inevitable association with the “assignment problem” appears undesirable.

Hypothesis 3.1. *Every FIP game form is separable.*

The converse implication is obviously wrong. Every game form in Examples 2.1 and 2.2 is separable. For $n = 2$, the statement is proven in Boros et al. (2008b).

The simplest separable game forms are *dictatorial* ones, where there is a player $i \in N$ such that $g(x_N) = p(x_i)$ for all $x_N \in X_N$; see Example 2.1a. Unless $g(X_N)$ is a singleton, there cannot be more than one dictator; $p(x_j)$ for $j \neq i$ may be arbitrary.

A game form is *simple* if it is separable and there is a linear order on \mathcal{X} such that

$$g(x_N) = p(\min_{i \in N} x_i) \quad (3.2)$$

for every $x_N \in X_N$. (Again, the same g may be described by (3.2) with different orders on \mathcal{X} .) Besides a desirable outcome, every strategy specifies a priority of the desire; then the choice with the highest priority is implemented. If G is dictatorial, we may define $x_i < x_j$ for every strategy of the dictator i and all $j \neq i$. Every fragment of a simple game form is simple as well.

Remark. Boros et al. (2008ab) call such game forms “totally reducible”; however, a lone ad-jective seems preferable.

Given a game form, we call $x_i \in X_i$ *simple* if $\#g(x_i, X_{-i}) = 1$.

Proposition 3.2. *A game form G is simple if and only if there is a simple strategy in every fragment of G .*

Proof. If G is simple and G' is a fragment of G , we pick the minimal strategy in $\bigcup_{i \in N} X'_i$; (3.2) immediately implies that it is simple in G' . Conversely, we pick a simple strategy in \mathcal{X} and declare it the least in \mathcal{X} ; then we forget it and define a linear order on \mathcal{X} by induction. \square

Theorem 3.3. *Every simple game form has the FIP.*

Proof. Suppose to the contrary that $x_N^0, x_N^1, \dots, x_N^m = x_N^0$ ($m > 0$) is an individual improvement cycle in a derivative game $G(v_N)$. Without restricting generality, we may assume that there is no shorter improvement cycle. For each $i \in N$, we define $X'_i = \{x_i^0, x_i^1, \dots, x_i^{m-1}\}$. By Proposition 3.2, the fragment contains a simple strategy; let it be x_i^k . Without restricting generality, we may assume that $x_N^k \triangleright_{\text{Ind}_i} x_N^{k-1}$, hence $x_N^{k+1} \triangleright_{\text{Ind}_j} x_N^k$ with $j \neq i$. Since $x_i^k = x_i^{k+1}$ and x_i^k is simple, we have $g(x_N^k) = g(x_N^{k+1})$, which contradicts the supposed inequality $u_j(x_N^{k+1}) > u_j(x_N^k)$. \square

All game forms in Example 2.1 except the last one are simple. Theorem 5 of Boros et al. (2008a) shows that every two person game form which has the FIP but is not simple must contain that matrix. When $n > 2$, even that much cannot be asserted, see Example 4.8 below.

“Dually” to dictatorial game forms, priorities of the strategies may be determined by the outcomes pointed to. Let A be linearly ordered and

$$g(x_N) = \min_{i \in N} p(x_i) \quad (3.3)$$

for each $x_N \in X_N$; Example 2.1c presents such a game form. A possible interpretation: each player chooses a maximal “level of cooperation” she is ready to accept; then the highest level acceptable to everybody becomes the norm. The term *unanimity game forms* may be appropriate.

Proposition 3.4. *Every unanimity game form has the FIP.*

Proof. The game form is simple: $y_i > x_j$ whenever $p(y_i) > p(x_j)$; the strategies with the same $p(x_i)$ are ordered arbitrarily. Now Theorem 3.3 applies. \square

Generalizing the notion, we may assume that an outcome is socially acceptable if a certain fraction of players finds it so; the median is most usual in political sciences. Moulin (1980) considered such game forms, with the addition of “fixed votes” to the players’ choices, and established some nice properties of them when the preferences are single-peaked. Here we abandon the anonymity requirement and allow arbitrary preferences. Thus, the order on the set of outcomes becomes an element of the decision making procedure, unrelated to the players’ preferences.

An *ordered voting game form* is defined by the following construction. A is linearly ordered and $X_i = A$ for each $i \in N$. Strictly speaking, our assumption $X_i \cap X_j = \emptyset$ is thus violated, but it does not matter here. For each $i \in N$, a “weight” $\mu_i \geq 0$ is given; for each $a \in A$, a number $\lambda_a > 0$. We assume that λ_a decreases in a . Given $x_N \in X_N$ and $a \in A$, we denote $N^-(a, x_N) = \{i \in N \mid x_i < a\}$, $\varkappa^-(a, x_N) = \sum_{i \in N^-(a, x_N)} \mu_i$, and define

$$g(x_N) = \max\{a \in A \mid \varkappa^-(a, x_N) < \lambda_a\}.$$

The interpretation is that each player chooses a “personal cap” x_i supported by her weight μ_i ; λ_a is the minimal total weight against a that makes it ineligible.

If all $\mu_i = 1$ and $\lambda_a = 1$, we have a unanimity game form, the FIP of which was established by Proposition 3.4; if all $\mu_i = 1$ and $\lambda_a = n$, we have $g(x_N) = \max_{i \in N} x_i$, i.e., Proposition 3.4 is still applicable after the order on A is reversed. Generally, there is no FIP.

Example 3.5. Let us consider an ordered voting game form G with $N = \{1, 2\}$, $A = \{a, b, c\}$ ($a > b > c$), both $\mu_i = 1$, $\lambda_c = \lambda_b = 2$, and $\lambda_a = 1$ (median voting scheme with a fixed vote at b). The game form is described by the following matrix:

$$\begin{array}{ccc} \underline{a} & b & \underline{b} \\ b & b & b \\ \underline{b} & b & \underline{c} \end{array}$$

The underlined outcomes form an improvement cycle in $G(v_N)$ with appropriate utilities; therefore, G does not have the FIP (although it is Nash consistent).

Somewhat unnatural behavior may be observed in the cycle: in the “clockwise” movement, player 2 switches from a to b by choosing c , and from c to b by choosing a .

Given $a, b \in A$, we define the *order distance* between them as

$$d(a, b) = \#\{x \in A \mid \min\{a, b\} \leq x < \max\{a, b\}\}; \quad (3.4)$$

then we define an admissible change relation by (2.10) with d as $\delta_{x_{-i}}$ for every $i \in N$ and $x_{-i} \in X_{-i}$. In other words, a change of strategy is admissible if the same change of outcome could not be produced by a shorter movement of x_i .

Theorem 3.6. *Every ordered voting game form has an almost unrestricted FIP with the quasidistance on each X_i defined by (3.4) [independently of x_{-i}].*

Proof. We denote $M = \{\mu_i\}_{i \in N} \subset \mathbb{R}$ and $N(m) = \{i \in N \mid \mu_i \geq m\}$ for each $m \in M$. Given $x_N \in X_N$ and $m \in M$, we define $\gamma_m^-(x_N) = \max\{a \in A \mid \varkappa^-(a, x_N) + m < \lambda_a\}$ and $\gamma_m^+(x_N) = \max\{a \in A \mid \varkappa^-(a, x_N) - m < \lambda_a\}$; clearly, $\gamma_m^-(x_N) \leq g(x_N) \leq \gamma_m^+(x_N)$. Then we define $C_m(x_N) = \{a \in A \mid \gamma_m^-(x_N) \leq a \leq \gamma_m^+(x_N)\}$, $B_m(x_N) = \{i \in N(m) \mid x_i \in C_m(x_N)\}$, and $P_m(x_N) = \sum_{i \in B_m(x_N)} v_i(x_i)$. A binary relation on X_N is defined as a lexicography:

$$y_N \succeq_m x_N \iff \left[C_m(y_N) \subset C_m(x_N) \text{ or } [C_m(y_N) = C_m(x_N) \& (B_m(y_N) \supset B_m(x_N) \text{ or } [B_m(y_N) = B_m(x_N) \& P_m(y_N) \geq P_m(x_N)])] \right];$$

$$y_N \succ_m x_N \iff \left[C_m(y_N) \subset C_m(x_N) \text{ or } [C_m(y_N) = C_m(x_N) \& (B_m(y_N) \supset B_m(x_N) \text{ or } [B_m(y_N) = B_m(x_N) \& P_m(y_N) > P_m(x_N)])] \right].$$

Clearly,

$$y_N \succ_m x_N \iff [y_N \succeq_m x_N \& x_N \not\succeq_m y_N].$$

Finally, a lexicographic aggregate is formed of \succeq_m ($m \in M$):

$$y_N \succ x_N \iff \exists m \in M [y_N \succ_m x_N \& \forall m' > m [y_N \succeq_{m'} x_N]].$$

Obviously, \succ is a strict order; we'll show that it is a potential, in the sense of (2.5b), for admissible improvements. Let $i \in N$ and $x_N, y_N \in X_N$ be such that $y_{-i} = x_{-i}$; we denote $a = g(x_N)$ and $b = g(y_N)$ and assume $b \neq a$. For every $c \in A$, we denote $c+1$ the next point in A , uniquely defined by $c+1 > c$ and $d(c, c+1) = 1$.

Step 3.6.1. $b \in C_{\mu_i}(x_N)$.

Proof. Since $\varkappa^-(b, y_N) < \lambda_b$ while $\varkappa^-(b, y_N) \geq \varkappa^-(b, x_N) - \mu_i$, we have $\varkappa^-(b, x_N) < \lambda_b + \mu_i$, hence $b \geq \gamma_{\mu_i}^-(x_N)$. For each $c > \gamma_{\mu_i}^+(x_N)$, we have $\varkappa^-(c, y_N) \geq \varkappa^-(c, x_N) - \mu_i \geq \lambda_c$, hence $c \neq b$; therefore, $b \leq \gamma_{\mu_i}^+(x_N)$. \square

Step 3.6.2. $x_N \vdash_i y_N \iff y_i = b$.

Proof. If $x_i < a$, then $\varkappa^-(a, y_N) \leq \varkappa^-(a, x_N) < \lambda_a$, hence $b > a$. If $x_i > a$, then $\varkappa^-(a+1, y_N) \geq \varkappa^-(a+1, x_N) \geq \lambda_{a+1}$, hence $b < a$.

Let $b > a$; then $x_i \leq a$. If $y_i < b$, then $\varkappa^-(b, y_N) = \varkappa^-(b, x_N) \geq \lambda_b$: a contradiction. Obviously $\min_{y_i \geq b_i} d(x_i, y_i)$ is attained when $y_i = b$.

Let $b < a$; then $x_i \geq a$. If $y_i > b$, then $\varkappa^-(b+1, y_N) = \varkappa^-(b+1, x_N) < \lambda_{b+1}$: a contradiction. Obviously $\min_{y_i \leq b_i} d(x_i, y_i)$ is attained when $y_i = b$. \square

In the following, we assume $y_i = b$.

Step 3.6.3. $C_m(y_N) \subseteq C_m(x_N)$ for each $m \geq \mu_i$.

Proof. Let $b > a$; then $\varkappa^-(c, y_N) \leq \varkappa^-(c, x_N)$ for each $c \leq b$, while $\varkappa^-(c, y_N) = \varkappa^-(c, x_N)$ for each $c > b$. Since $\gamma_m^-(x_N) \leq a < b \leq \gamma_m^+(x_N)$, we have $\gamma_m^-(y_N) \geq \gamma_m^-(x_N)$ and $\gamma_m^+(y_N) = \gamma_m^+(x_N)$.

The case of $b < a$ is treated dually. \square

Step 3.6.4. $y_N \succeq_m x_N$ for each $m > \mu_i$.

Proof. Indeed, $C_m(y_N) \subseteq C_m(x_N)$ by Step 3.6.3. If the inclusion is strict, we even have $y_N \succ_m x_N$. Otherwise, $B_m(y_N) = B_m(x_N)$ and $P_m(y_N) = P_m(x_N)$ since $y_{N(m)} = x_{N(m)}$. \square

Step 3.6.5. If $v_i(b) > v_i(a)$, then $y_N \succ_{\mu_i} x_N$.

Proof. Again, $C_{\mu_i}(y_N) \subseteq C_{\mu_i}(x_N)$ by Step 3.6.3. Since $i \in B_{\mu_i}(y_N)$, we only have to consider the case of $C_{\mu_i}(y_N) = C_{\mu_i}(x_N)$ and $B_{\mu_i}(y_N) = B_{\mu_i}(x_N)$, i.e., $i \in B_{\mu_i}(x_N)$. If $x_i = a$, we have $P_{\mu_i}(y_N) - P_{\mu_i}(x_N) = v_i(b) - v_i(a) > 0$. Suppose $x_i < a$; then $\gamma_m^-(x_N) \leq x_i < a$ and $b = y_i > a$. On the other hand, $\varkappa^-(a, y_N) \geq \varkappa^-(a, x_N) - \mu_i$, hence $\varkappa^-(a, y_N) + \mu_i < \lambda_a$, hence $\gamma_{\mu_i}^-(y_N) \geq a > \gamma_{\mu_i}^-(x_N)$, hence $C_{\mu_i}(y_N) \subset C_{\mu_i}(x_N)$, contradicting our assumption. The case of $x_i > a$ is treated dually. \square

In the light of Steps 3.6.4 and 3.6.5, the proof of the theorem is accomplished. \square

Theorem 3.6 becomes wrong if strategy sets $X_i \subset A$ are allowed: the assumption $X_2 = \{a, c\}$ in Example 3.5 would make the changes made by player 2 along the cycle unique.

Remark. In the absence of fixed votes, i.e., when λ_a is the same for all $a \in A$, an ordered voting game form is obviously separable: $p(x_i) = x_i$. The statement seems to be wrong generally although I have not studied the question carefully.

3.2 Games with perfect information

The most important examples of game forms with FIP or almost unrestricted FIP are provided by the normal form of games with perfect information. Following Kukushkin (2002a), we reproduce the familiar concepts in a fashion most convenient for our purposes.

A *perfect information game form (PIGF)* is a game form with arbitrary (finite) sets N and A , and strategies and the mapping g generated by a construction as follows. A *game tree* K is a finite partially ordered set satisfying these two conditions. (a) For every $\alpha \in K$, the set $\{\beta \in K \mid \beta \leq \alpha\}$ is a chain. (b) There exists the minimum α_0 of K (the *origin*), $\alpha_0 \leq \alpha$ for every $\alpha \in K$. The existence of the meet (greatest common lower bound) $\alpha \wedge \beta$ for every $\alpha, \beta \in K$ easily follows. We call $\beta \in K$ an *immediate successor* of $\alpha \in K$ if $\alpha < \beta$ while $\alpha < \beta' < \beta$ is impossible; the set of all immediate successors of $\alpha \in K$ is denoted X_α . Imagining an arc from every $\alpha \in K$ to every $\beta \in X_\alpha$ turns K into a tree in a geometrical sense.

The set of maximizers of the order on K is denoted T (*terminal nodes*); the set $K \setminus T$, D (*decision nodes*). There is an *ownership mapping* $\nu: D \rightarrow N$; player i moves at nodes from $D_i = \nu^{-1}(i)$. We denote $X_i = \prod_{\alpha \in D_i} X_\alpha$ for $i \in N$ and $X_N = \prod_{i \in N} X_i$. We identify X_N with $\prod_{\alpha \in D} X_\alpha$. For every $x_N \in X_N$ and $\alpha \in D$, we denote $\pi(\alpha, x_N) \subseteq K$ the intersection of all subsets K' of K satisfying these two conditions: $\alpha \in K'$ and $[\beta \in K' \Rightarrow x_\beta \in K']$; note that $\pi(\alpha, x_N)$ is a chain: a play of the game starting at α . The unique element of $T \cap \pi(\alpha, x_N)$ is denoted $\tau(\alpha, x_N)$: the result of playing x_N starting at α .

Finally, there is a mapping $\gamma: T \rightarrow A$ and $g(x_N) = \gamma(\tau(\alpha_0, x_N))$. If γ is a bijection, G is called *free*; in this case, we may just assume $A = T$ (as was done in Kukushkin, 2002a). For each $\alpha \in D$, we denote $F(\alpha) = \gamma(\{\beta \in T \mid \beta > \alpha\})$, the set of outcomes feasible if the play passes through α .

To avoid pathologies, we always assume that $\#X_\alpha > 1$ and $\#F(\alpha) > 1$ for every $\alpha \in D$ (i.e., there is no “meaningless” decision node), while $X_\alpha \cap D_i = \emptyset$ for all $i \in N$ and $\alpha \in D_i$ (i.e., no player can make two consecutive choices). The restrictions are innocuous and allow more compact formulations of Theorem 3.8, Hypothesis 3.10, and Proposition 3.19 below.

Proposition 3.7. *A game form is simple if and only if it can be represented as a perfect information game form where D is a chain in the tree order, i.e., there exists a play of the game containing all decision nodes.*

Proof. Straightforward. □

Theorem 3.8 (Theorem 1 of Kukushkin, 2002a). *If G is a free perfect information game form, then G has the FIP if and only if each D_i is a chain in the tree order, i.e., for each player there exists a play of the game containing all his decision nodes.*

The sufficiency part, naturally, holds for every PIGF, which cannot be said about the necessity.

Example 3.9. Let us consider a three-person perfect information game form with five decision nodes, six terminal nodes, and five outcomes. We assume $D_1 = \{\alpha_0, \alpha_2, \alpha_3\}$, $D_2 = \{\beta_2\}$, $D_3 = \{\beta_3\}$ and $A = \{a, b, c, d, e\}$; γ is described by putting outcomes at terminal nodes.

$$\begin{array}{ccccccccc}
 a & \xleftarrow{1} & \alpha_2 & \xleftarrow{2} & \beta_2 & \xleftarrow{1} & \alpha_0 & \xrightarrow{1} & \beta_3 & \xrightarrow{3} & \alpha_3 & \xrightarrow{1} & e \\
 & & \downarrow 1 & & \downarrow 2 & & & & \downarrow 3 & & \downarrow 1 & & \\
 & & b & & c & & & & c & & d & &
 \end{array}$$

The FIP is not difficult to check.

To extend Theorem 3.8 to the general case, we introduce the following requirements. First, the decision nodes of each player form a subsemilattice of the game tree, i.e., if a player moves at two incomparable nodes, he must move at their meet too:

$$\alpha, \alpha' \in D_i \Rightarrow \alpha \wedge \alpha' \in D_i. \quad (3.5a)$$

The second condition needs some auxiliary notations. For $i \in N$ and $\alpha, \beta \in D_i$, we denote:

$$\begin{aligned}
 F^*(\alpha, \beta) &= \gamma(\{t \in T \mid \alpha < t \wedge \beta < t \wedge t \wedge \beta \notin D_i\}); \\
 F^{**}(\alpha, \beta) &= \begin{cases} F^*(\alpha, \beta), & \text{if } \#F(\beta) > 2; \\ F^*(\alpha, \beta) \setminus F(\beta), & \text{if } \#F(\beta) = 2. \end{cases}
 \end{aligned}$$

Now the condition is

$$\forall i \in N \forall \alpha \in D_i \forall \beta', \beta'' \in D_i \setminus \{\alpha\} [\beta' \wedge \beta'' = \alpha \ \& \ c' \in F^{**}(\alpha, \beta') \ \& \ c'' \in F^{**}(\alpha, \beta'') \Rightarrow c' = c'']. \quad (3.5b)$$

If D_i is a chain, then (3.5a) holds trivially while (3.5b) holds by default. Generally, conditions (3.5) mean that each D_i contains a subchain D_i^* such that if player i makes the play to turn towards a decision node from $D_i \setminus D_i^*$ then all other players have a rather little say in determining the outcome.

Hypothesis 3.10. *A perfect information game form has the FIP if and only if it satisfies assumptions (3.5).*

The necessity part is easily derived from the proof of Theorem 1 in Kukushkin (2002a).


Proposition 3.11. *Every perfect information game form satisfying assumptions (3.5) is separable.*

Proof. Given $i \in N$ and $x_i \in X_i$, we define $\alpha(x_i)$ as the greatest in the tree order (i.e., the furthest from the origin) decision node of player i that can be reached when player i chooses x_i . It is well defined because of (3.5a). Now if x_i prescribes to player i to choose a terminal node t at $\alpha(x_i)$, i.e., if $x_{\alpha(x_i)} = t \in T$, we define $p(x_i) = \gamma(t)$; otherwise, $p(x_i) \in A$ is arbitrary.

Checking (3.1) is easy. Let $\tau(\alpha_0, x_N) = t$, hence $g(x_N) = \gamma(t)$. We denote i the player who made the last move in the play ending at t , and β the decision node where that last move was made. Obviously, $\beta = \alpha(x_i)$ and $\gamma(t) = p(x_i)$. \square

By Proposition 3.7, every two-person perfect information game form satisfying assumptions (3.5) is simple. The converse to Proposition 3.11 is wrong: the proof only needs (3.5a). On the other hand, Proposition 3.11 becomes wrong if assumptions (3.5) are dropped altogether.

Example 3.12. Let us consider a free two-person perfect information game form with three decision nodes; we assume $D_1 = \{\alpha_0\}$ and $D_2 = \{\beta', \beta''\}$.



If it were a separable game form, we could, without restricting generality, assume $p(\beta') = a$ and $p(\beta'') = d$. Now if $p(b, c) \neq b$, then (3.1) is violated for $x_N = \langle \beta', (b, c) \rangle$; if $p(b, c) \neq c$, then (3.1) is violated for $x_N = \langle \beta'', (b, c) \rangle$.

Given a perfect information game form, we define the distance between two strategies of the same player $i \in N$ as

$$d(x_i, y_i) = \#\{\alpha \in D_i \mid y_\alpha \neq x_\alpha\}. \quad (3.6)$$

Then we define admissible change relations for each $i \in N$ by (2.10) with the quasidistances $\delta_{x_{-i}}(x_i, y_i) = d(x_i, y_i)$ for every $x_{-i} \in X_{-i}$. In other words, a change of the strategy of a player is admissible if the number of nodes involved is minimal, i.e., there was no change at irrelevant nodes.

Theorem 3.13 (Theorem 3 of Kukushkin, 2002a). *Every perfect information game form has an almost unrestricted FIP with admissible change relations defined by the “costs” (3.6).*

Remark. A modification of the proof from Kukushkin (2002a) allows us to derive a corollary (Proposition 3.18 in the next subsection), which may be of some interest.

Proof. Let $x_N \in X_N$ and $t \in T$; we say that t is blocked by player i at x_N if there exist $\alpha \in D_i$ and $\beta \in X_\alpha$ such that $t \in \pi(\beta, x_N) \setminus \pi(\alpha, x_N)$ (it follows immediately that $\tau(\beta, x_N) = t$ and $x_\alpha \neq \beta$). Let $t \neq \tau(\alpha_0, x_N)$; then the set $\{\alpha < t \mid t \notin \pi(\alpha, x_N)\}$ is a nonempty chain, so we may pick its maximum α and denote $i = \nu(\alpha)$. Obviously, t is blocked by i at x_N and t cannot be blocked at x_N by any other player. We thus obtain a partitioning of $T \setminus \{\tau(\alpha_0, x_N)\}$ into subsets $B_i(x_N)$ of terminal nodes blocked by each particular player $i \in N$ at x_N .

It is important to note that $\#B_i(x_N)$ does not depend on x_N ; actually, $\#B_i(x_N) = [\sum_{\alpha \in D_i} \#X_\alpha] - \#D_i$ because choosing any alternative from X_α for $\alpha \in D_i$, player i does not choose each of $\#X_\alpha - 1$ others thereby blocking the same number of terminal nodes (exactly which terminal nodes are blocked depends on the choices at successive nodes).

Defining $P(x_N) = -\sum_{i \in N} \sum_{t \in B_i(x_N)} v_i \circ \gamma(t)$, let us show that P is a numeric potential of \gg . Assuming $y_N \gg_i x_N$, we have to prove that $P(y_N) > P(x_N)$.

We denote $t = \tau(\alpha_0, x_N)$ and $t' = \tau(\alpha_0, y_N)$; then $v_i \circ \gamma(t') > v_i \circ \gamma(t)$ and $t \wedge t' \in D_i$. Let us show first $B_j(x_N) \subseteq B_j(y_N)$ for each $j \neq i$. Assuming $t'' \in B_j(x_N)$, we, by the definition of blocking, have $\alpha \in D_j$ and $\beta \in X_\alpha$ such that $t'' \in \pi(\beta, x_N) \setminus \pi(\alpha, x_N)$. Since $j \neq i$, $y_\alpha = x_\alpha$, hence $t'' \notin \pi(\alpha, y_N)$ and $t'' \neq t'$; moreover, $t'' \wedge t' \leq \alpha$. Now $t'' \neq \tau(\beta, y_N)$ could only be possible if there were $\beta' \in \pi(\beta, x_N)$ such that $y_{\beta'} \neq x_{\beta'}$, hence $\beta' \in D_i$; but then the replacement of $x_{\beta'}$ with $y_{\beta'}$ would be a superfluous change incompatible with the minimization of “costs” (3.6). More formally, considering z_i which coincides with x_i at β' and with y_i at all other nodes from D_i , we immediately see that $g(z_i, x_{-i}) = g(y_i, x_{-i}) = g(y_N)$ while $\delta_{x_{-i}}(x_i, z_i) < \delta_{x_{-i}}(x_i, y_i)$. Therefore, $\tau(\beta, y_N) = t''$, hence $t'' \in B_j(y_N)$.

Now $B_j(x_N) \subseteq B_j(y_N)$ implies $B_j(x_N) = B_j(y_N)$ for all $j \neq i$. It follows immediately that $B_i(y_N) = (B_i(x_N) \setminus \{t'\}) \cup \{t\}$, hence

$$P(y_N) - P(x_N) = v_i \circ \gamma(t) - v_i \circ \gamma(t') = u_i(y_N) - u_i(x_N). \quad (3.7)$$

Thus, we have $P(y_N) > P(x_N)$. □

An extension of the notion of a PIGF is met in the literature quite often, see, e.g., Boros and Gurvich (2003) and references therein. Suppose there is a directed graph, its nonterminal nodes are partitioned among the players, and one of them is fixed as the origin. Each player is free to choose an arc leading from each of her decision nodes (“stationary strategies”); once

all strategies are chosen, a play of the game is uniquely defined: after a finite number of steps, either a terminal node is reached, or a cycle starts repeating itself ad infinitum. Treating cycles as additional outcomes, we obtain a game form.

If the players were allowed to condition their choices on history, we would return to a game on a (finite or infinite) tree. The stationarity requirement is effectively equivalent to imperfect information: when making a decision at a node, the player does not know how the play has come there. It is well known that a game with imperfect information, even on a finite tree, need not possess a Nash equilibrium (unless mixed strategies are invoked). In *this* class of games, however, the lack of information only concerns the past, which does not affect future possibilities. Therefore, it seems natural to expect no big difference with the standard perfect information model.

Indeed, if the graph is acyclic, i.e., the underlying tree is finite, an equilibrium can be obtained by the standard backward recursion. If cycles are possible, the underlying tree becomes infinite, and there is nowhere to derive an equilibrium from. An intermediate case emerges when the graph contains cycles, but every cycle is worse for each player than any terminal node (Boros and Gurvich, 2003); then no infinite play can result from any improvement.

Here we briefly consider the possibility to extend Theorem 3.13 to a game on an acyclic graph. The result, Theorem 3.14 below, is distinctly weaker; to be more precise, a restricted FIP is established, but an almost unrestricted FIP is not (so far).

The definition of a *positional acyclic game form* is most conveniently given in the same style as in the beginning of this subsection. Instead of a game *tree*, we consider a finite partially ordered set K satisfying condition (b) from the definition of a PIGF: the existence of the origin α_0 . The partition of K into *decision nodes* D and *terminal nodes* T is the same; the *ownership mapping* $\nu: D \rightarrow N$ is the same; the definitions of *immediate successors* X_α of $\alpha \in K$, as well as *strategies* $X_i = \prod_{\alpha \in D_i} X_\alpha$ and *strategy profiles* $X_N = \prod_{i \in N} X_i = \prod_{\alpha \in D} X_\alpha$ are the same. Since condition (a) from the definition of a PIGF is dropped, we may always assume that γ is a bijection, i.e., $T = A$. A play of the game starting at α , $\pi(\alpha, x_N) \subseteq K$, is defined in the same way, and still is a chain; the result of playing x_N starting at α , $\tau(\alpha, x_N)$, is still uniquely defined. Finally, the assumption $X_\alpha \cap D_i = \emptyset$ for all $i \in N$ and $\alpha \in D_i$ no longer looks innocuous and is not made.

We start with the definition of a strengthened version of restricted FIP. Given a family of “costs” $\delta_{x_{-i}}$ on each X_i , we define the *cost-efficient individual improvement relation* on X_N by

$$y_N \triangleright_i x_N \iff [y_N \triangleright_i^{\text{Ind}} x_N \ \& \ \nexists z_i \in X_i [\delta_{x_{-i}}(x_i, z_i) < \delta_{x_{-i}}(x_i, y_i) \ \& \ u_i(z_i, x_{-i}) \geq u_i(y_N)]] \quad (3.8a)$$

and

$$y_N \triangleright x_N \Leftrightarrow \exists i \in N [y_N \triangleright_i x_N]. \quad (3.8b)$$

A game form has an *FCIP* if there are “costs” $\delta_{x_{-i}}$ on each X_i such that the cost-efficient individual improvement relation (3.8) is acyclic in every derivative game.

Theorem 3.14. *Every positional acyclic game form has an FCIP with the “costs” (3.6).*

Proof. Given $x_N \in X_N$ and $\alpha \in D$, we define $v_\alpha(x_N) = v_{\nu(\alpha)}(\tau(\alpha, x_N))$: what the player who moves at α would get if the play passes through the node. Then we define a lexicographic order on X_N :

$$y_N \succ x_N \Leftrightarrow [\exists \alpha \in K [v_\alpha(y_N) > v_\alpha(x_N)] \& \forall \alpha \in K (v_\alpha(y_N) < v_\alpha(x_N) \Rightarrow \exists \beta \in K [\beta > \alpha \& v_\beta(y_N) > v_\beta(x_N)])]. \quad (3.9)$$

Clearly, \succ is a strict order.

Supposing that $x_N, y_N \in X_N$ and $y_N \triangleright_i x_N$, we show that $y_N \succ x_N$. We denote $B = \{\beta \in K \mid y_\beta \neq x_\beta\} \neq \emptyset$. Exactly as in the proof of Theorem 3.13, we have $B \subseteq D_i \cap \pi(\alpha_0, y_N)$, hence there exists $\beta^* = \max B$.

Let us show that $v_{\beta^*}(y_N) > v_{\beta^*}(x_N)$. Supposing the contrary, we define $z_N \in X_N$ by $z_{\beta^*} = x_{\beta^*}$ and $z_\alpha = y_\alpha$ for all $\alpha \neq \beta^*$. Clearly, $u_i(z_N) = v_{\beta^*}(z_N) = v_{\beta^*}(x_N) \geq v_{\beta^*}(y_N) = u_i(y_N)$. Since $d(x_i, z_i) < d(x_i, y_i)$, we have a contradiction with the assumption $y_N \triangleright_i x_N$.

Finally, let $\alpha \in K$ and $v_\alpha(y_N) \neq v_\alpha(x_N)$, hence $B \cap \pi(\alpha, y_N) \neq \emptyset$. Picking $\beta \in B \cap \pi(\alpha, y_N)$, we have $\beta^* \geq \beta \geq \alpha$. Since $v_{\beta^*}(y_N) > v_{\beta^*}(x_N)$, (3.9) holds. \square

It is impossible to derive an almost unrestricted FIP from the above argument: the inequality $v_{\beta^*}(y_N) > v_{\beta^*}(x_N)$ need not hold without the “cost-efficiency” of the improvement. Whether the conditions of Theorem 3.14 imply the property itself remains an open question.

Proposition 3.15. *A positional acyclic game form has an almost unrestricted FIP with admissible change relations defined by the “costs” (3.6) if each D_i is an anti-chain, i.e., all $\beta, \beta' \in D_i$ are incomparable in the order on K .*

Proof. Let $y_N \triangleright_i x_N$. In the notation from the proof of Theorem 3.14, we have $B = \{\beta^*\}$, hence $v_{\beta^*}(y_N) = u_i(y_N) > u_i(x_N) = v_{\beta^*}(x_N)$, hence $y_N \succ x_N$. \square

It is funny to contrast Proposition 3.15 with Theorem 3.8.

Proposition 3.16. *A positional acyclic game form has the FIP if each D_i is a singleton.*

Proof. Retaining the notation from the proof of Theorem 3.14 and arguing in the same way, it is easy to show that $y_N \succ x_N$ whenever $y_N \triangleright^{\text{Ind}} x_N$. \square

The converse to Proposition 3.16 is wrong. There is no plausible hypothesis about necessary and sufficient conditions for FIP in this class of game forms.

3.3 Cardinal utilities and potentials

Although our main subject are games with ordinal preferences and we do not consider mixed extensions, it seems impossible not to say a few words about cardinal utilities. Actually, Monderer and Shapley (1996) paid most attention to that case. Moreover, they assumed that the utilities of all players are measured in the same scale; we make the same assumption here.

Let Γ be a game with such “co-cardinal” utilities. Monderer and Shapley (1996) defined an *exact potential* of Γ as a function $P: X_N \rightarrow \mathbb{R}$ such that

$$u_i(y_N) - u_i(x_N) = P(y_N) - P(x_N) \quad (3.10)$$

whenever $i \in N$, $y_N, x_N \in X_N$, and $y_{-i} = x_{-i}$. An *ordinal potential* of Γ is a function $P: X_N \rightarrow \mathbb{R}$ such that

$$\text{sign}(u_i(y_N) - u_i(x_N)) = \text{sign}(P(y_N) - P(x_N)) \quad (3.11)$$

whenever $i \in N$, $y_N, x_N \in X_N$, and $y_{-i} = x_{-i}$. The latter notion is indeed ordinal, i.e., invariant under strictly increasing transformations of utilities. Voorneveld and Norde (1996) showed that the existence of an ordinal potential is equivalent to the absence of “weak improvement cycles.”

Obviously, (3.10) implies (3.11), of which it is the most natural cardinal analogue.

Theorem 3.17. *For every game form G , the following statements are equivalent.*

1. *Every derivative game $G(v_N)$ admits an exact potential.*
2. *Every derivative game $G(v_N)$ admits an ordinal potential.*
3. *G is dictatorial.*

Proof. The implications [Statement 3 \Rightarrow Statement 1 \Rightarrow Statement 2] are straightforward. Let Statement 2 hold.

Step 3.17.1. *If $i, j \in N$, $i \neq j$, $x_N, y_N, z_N \in X_N$, $x_{-i} = y_{-i}$, $y_{-j} = z_{-j}$, and $g(x_N) \neq g(y_N)$, then $g(y_N) = g(z_N)$.*

Proof. Supposing the contrary, we denote $y'_N = (z_j, x_{-j})$. Then we pick a constant as v_j , and define $v_i(g(x_N)) = 1$ and $v_i(g(y_N)) = 0$. If $g(y'_N) = g(z_N)$, we obtain a weak improvement cycle in $G(v_N)$: $v_i(g(x_N)) > v_i(g(y_N))$, $v_j(g(y_N)) = v_j(g(z_N))$, $v_i(g(z_N)) = v_i(g(y'_N))$, and $v_j(g(y'_N)) = v_j(g(x_N))$; clearly, this is incompatible with (3.11). Otherwise, if $g(x_N) = g(z_N)$, we set $v_i(g(y'_N)) = 0$ and again obtain the same cycle: $v_i(g(x_N)) > v_i(g(y_N))$, $v_j(g(y_N)) = v_j(g(z_N))$, $v_i(g(z_N)) > v_i(g(y'_N))$, and $v_j(g(y'_N)) = v_j(g(x_N))$. Finally, if $g(x_N) \neq g(z_N) \neq g(y'_N)$, we set $v_i(g(y'_N)) = v_i(g(z_N))$ [a numeric value cannot be specified because both $g(y'_N) = g(x_N)$ and $g(y'_N) = g(y_N)$ are possible] and again obtain the same cycle. \square

Step 3.17.2. *If $i \in N$, $x_N, y_N \in X_N$, $x_{-i} = y_{-i}$, and $g(x_N) \neq g(y_N)$, then player i is a dictator.*

Proof. Given $j \in N \setminus \{i\}$, Step 3.17.1 immediately implies that $g(y_i, z_j, x_{-ij}) = g(y_N)$ and $g(x_i, z_j, x_{-ij}) = g(x_N)$ for all $z_j \in X_j$. For every $z_N \in X_N$ such that $z_{-i} = x_{-i} = y_{-i}$, we have either $g(z_N) \neq g(x_N)$ or $g(z_N) \neq g(y_N)$, hence the previous argument applies and player j cannot change the outcome. Therefore, if $x'_{-j} = x_{-j}$ and $y'_{-j} = y_{-j}$, then the assumptions of the lemma hold for x'_N and y'_N as well, hence no player k can change the outcome deviating from x_k at x'_{-k} . Iterating the reasoning, we see that player i is a dictator. \square

If $\#g(X_N) = 1$, then every player is a dictator. If $g(x_N) \neq g(y_N)$, then there is a strategic path $x_N^0, x_N^1, \dots, x_N^n$ such that $x_N^0 = x_N$ and $x_N^n = y_N$. Clearly, we must have $g(x_N^k) \neq g(x_N^{k+1})$ for some k , hence Step 3.17.2 applies. \square

We call G a *nearly potential game form* if there is a liberal admissible change relation \vdash_i on X_N for each $i \in N$, and a function $P: X_N \rightarrow \mathbb{R}$ for every derivative game $G(v_N)$ such that (3.10) holds whenever $i \in N$, $y_N, x_N \in X_N$, $x_N \vdash_i y_N$, and $g(y_N) \neq g(x_N)$. We call G an *almost potential game form* if there is a function $P: X_N \rightarrow \mathbb{R}$ for every derivative game $G(v_N)$ such that (3.10) holds whenever $i \in N$, $y_N, x_N \in X_N$, $x_{-i} = y_{-i}$, and $g(y_N) \neq g(x_N)$. Clearly, every almost potential game form has the FIP while every nearly potential game form has an almost unrestricted FIP.

Proposition 3.18. *Every perfect information game form is a nearly potential one. If ν is injective, i.e., each player has at most one decision node, then it is an almost potential game form.*

Proof. Defining the admissible change relations by the “costs” (3.6) and potential P as in the proof of Theorem 3.13, we refer to (3.7). If each D_i is a singleton, then every change of strategy producing a change of outcome is admissible. \square

Proposition 3.19. *If a free perfect information game form is an almost potential one, then ν is injective.*

Proof. The necessity of (3.5a) for the FIP is easy to see regardless of Hypothesis 3.10, so if ν is not injective, we may assume that $\nu(\alpha) = \nu(\alpha') = i$ while $\alpha < \alpha'$. By the non-degeneracy assumptions, there must be $\beta \in D_j$ such that $\alpha < \beta < \alpha'$ and $j \neq i$. Since G is free, there must be a fragment of the type

$$\begin{array}{ccc} a & b & b \\ a & c & d \end{array}$$

where player i chooses columns and player j rows, and different letters denote different outcomes. Individual changes of strategies starting at the northwestern corner produce this cycle of outcomes: $a \rightarrow b \rightarrow c \rightarrow d \rightarrow b \rightarrow a$. Therefore, (3.10) would imply that $v_i(b) - v_i(a) + v_j(c) - v_j(b) + v_i(d) - v_i(c) + v_j(b) - v_j(d) + v_i(a) - v_i(b) = 0$, hence $v_i(d) - v_i(c) = v_j(d) - v_j(c)$. Clearly, the equality does not hold for all utility functions. \square

The assumption that the game form is *free* cannot be dropped. Example 4.8 below disproves putative hypotheses like “every nearly (or almost) potential game form can be represented as a perfect information game form”; on the other hand, such examples seem rare, so a plausible hypothesis may be obtainable.

4 FCP Game Forms

4.1 Characterization

Given $I \in \mathcal{N} \setminus \{N\}$, a *superfragment* of G is a game form with two players, “ I ” and “ $-I$,” nonempty subsets of X_I and X_{-I} as strategy sets, respectively, and the appropriate restriction of g . The FCP⁽⁺⁾ of G obviously implies the same property of every superfragment; no such assertion holds for the FIP.

Following Boros et al. (2008a), we call a game form G *totally tight* if every 2×2 superfragment of G contains a simple strategy; in other words, if the condition

$$\{g(x'_I, x'_{-I}), g(x''_I, x''_{-I})\} \cap \{g(x'_I, x''_{-I}), g(x''_I, x'_{-I})\} \neq \emptyset \quad (4.1)$$

holds for each $I \in \mathcal{N}$, $x'_I, x''_I \in X_I$ and $x'_{-I}, x''_{-I} \in X_{-I}$.

Proposition 4.1. *Let G be a game form such that $g(X_N) = \{q, a\}$. Then G is totally tight if and only if there is an ordering \succeq_I on X_I for every $I \in \mathcal{N}$ such that*

$$\forall I \in \mathcal{N} \forall x_N, x'_N \in X_N [g(x'_N) = q \ \& \ x'_I \succeq_I x_I \ \& \ x'_{-I} = x_{-I}] \Rightarrow g(x_N) = q. \quad (4.2)$$

An interpretation should be clear: all strategies of a player or a coalition are (weakly) ordered according to how much effort is spent in trying to switch from a *status quo ante* outcome q to the unique available alternative; whenever the effort proves insufficient, less effort cannot be sufficient. The term “a binary lobbying game form” may be appropriate.

Proof. Necessity. Let (4.1) hold everywhere. For each $I \in \mathcal{N}$, we define a binary relation on X_I :

$$y_I \succeq_I x_I \iff \forall x_{-I} \in X_{-I} [g(y_I, x_{-I}) = q \Rightarrow g(x_I, x_{-I}) = q]; \quad (4.3)$$

the relation is obviously a preorder. Incomparability of y_I and x_I would immediately imply a violation of (4.1); therefore, \succeq_I is an ordering for every $I \in \mathcal{N}$. The “monotonicity” requirement (4.2) immediately follows from (4.3).

Sufficiency. Suppose that (4.2) holds, but (4.1) is violated by a superfragment. Without restricting generality, $x'_I \succeq_I x''_I$ and $x''_{-I} \succeq_{-I} x'_{-I}$; therefore, $g(x'_I, x'_{-I}) = q$. The supposed negation of (4.1) implies that $g(x'_I, x'_{-I}) = g(x''_I, x''_{-I}) = a$, but then $g(x''_I, x''_{-I}) = a$ by (4.2). \square

Theorem 4.2. *A game form G has the FCP if and only if G is either dictatorial, or totally tight with $\#g(X_N) \leq 2$.*

Remark. So far, there is no explanation for the obvious similarity with Arrow’s impossibility theorem.

Proof. Sufficiency. A dictatorial game form poses no problem. Let G be totally tight with $g(X_N) = \{a, q\}$ and $x_N^0, x_N^1, \dots, x_N^m = x_N^0$ be a coalition improvement cycle in a derivative game $G(v_N)$. Without restricting generality, $g(x_N^0) = a$, hence $g(x_N^{2k}) = a$, $g(x_N^{2k+1}) = q$, and m is even. We denote $I = \{i \in N \mid v_i(a) > v_i(q)\}$, $J = \{i \in N \mid v_i(q) > v_i(a)\}$, and $N^0 = \{i \in N \mid v_i(a) = v_i(q)\}$. Each player $i \in N^0$ chooses the same strategy at each x_N^k ; without restricting generality, we may assume $N = I \cup J$. Obviously, $x_I^{2k+1} = x_I^{2k}$ and $x_J^{2k+2} = x_J^{2k+1}$ for all k . By Proposition 4.1, we have $x_I^{2k+2} \succ_I x_I^{2k+1} = x_I^{2k}$ for each k , which contradicts the assumption $x_N^m = x_N^0$.

Necessity. Let G have the FCP. We start with an auxiliary statement.

Step 4.2.1. $\#g(X'_N) \leq 2$ for every 2×2 superfragment of G .

Proof. Suppose the contrary: there is a superfragment of G of the type $\begin{smallmatrix} a & b \\ d & c \end{smallmatrix}$ with $a \neq b \neq c \neq a$. Let I choose rows and $-I$ columns. We consider a utility vector v_N such that $v_i(a) > v_i(c) > v_i(b)$ for each $i \in I$ and $v_i(b) > v_i(a) > v_i(c)$ for each $i \notin I$. A coalition improvement cycle in $G(v_N)$ is obvious: $a \rightarrow b \rightarrow c \rightarrow a$. \square

Suppose that G is not dictatorial. Then for each $i \in N$ there is $x_i \in X_i$ such that $\#g(x_i, X_{-i}) \geq 2$. Moreover, at least two players are not “dummies,” i.e., there is $x_{-i} \in X_{-i}$ such that $\#g(X_i, x_{-i}) \geq 2$. Whenever $i \in N$, $x_i, x'_i \in X_i$, $x_{-i}, x'_{-i} \in X_{-i}$ and $g(x'_i, x_{-i}) \neq g(x_i, x_{-i}) \neq g(x_i, x'_{-i})$, there holds $g(x'_i, x_{-i}) = g(x_i, x'_{-i})$ by Step 4.2.1. It follows immediately that $\#g(x_i, X_{-i}) \leq 2$ for every non-dummy $i \in N$ and $x_i \in X_i$.

If $\#g(X_N) > 2$, there must exist $i \in N$, $x_i, x'_i \in X_i$ and $a, b, c, d \in A$ such that $a \neq b \neq c \neq a$, $b \neq d \neq c$, $g(x_i, X_{-i}) = \{a, b\}$, and $g(x'_i, X_{-i}) = \{c, d\}$. We pick $x_{-i} \in X_{-i}$ such that $g(x_i, x_{-i}) = b$. If $g(x'_i, x_{-i}) = a$, we pick $x'_{-i} \in X_{-i}$ such that $g(x'_i, x'_{-i}) = c$ and obtain a contradiction with Step 4.2.1. If $g(x'_i, x_{-i}) = c$, we pick $x'_{-i} \in X_{-i}$ such that $g(x'_i, x'_{-i}) = a$ and again obtain a contradiction with Step 4.2.1.

Thus, either G is dictatorial or $\#g(X_N) = 2$. In the latter case, the necessity of (4.1) is shown exactly as in Moulin (1976). \square

Proposition 4.3. *Every FCP game form is separable.*

Proof. If G is dictatorial, there is nothing to prove. If $A = \{q, a\}$, we define $p(x_i) = q$ for all strategies of one player, $p(x_j) = a$ for another, and arbitrarily for all others. \square

Theorem 4.4. *A game form G has the FCP⁺ if and only if it has the FCP and either $\#g(X_N) = 1$ or $\#N \leq 2$.*

Proof. Sufficiency. Let $\#N = 2$. Given a utility vector v_N , we take v_N^* with the property

$$v_i^*(a) > v_i^*(b) \iff [v_i(a) > v_i(b) \text{ or } [v_i(a) = v_i(b) \ \& \ v_{-i}(a) > v_{-i}(b)]]$$

for each $i \in N$ and $a, b \in A$. Clearly, a weak coalition improvement path in $G(v_N)$ is a coalition improvement path in $G(v_N^*)$, hence G has the FCP⁺ as well as the FCP.

Necessity. Let $\#g(X_N) > 1$; then there are $i \in N$, $x_i, x'_i \in X_i$, and $x_{-i} \in X_{-i}$ such that $a = g(x_i, x_{-i}) \neq g(x'_i, x_{-i}) = b$. Supposing $\#N > 2$, we can pick $j, k \in N$ such that $j \neq i \neq k \neq j$. Whenever $v_i(a) = v_i(b)$, $v_j(a) > v_j(b)$, and $v_k(a) < v_k(b)$, we have a weak coalition improvement cycle where player i switches between x_i and x'_i back and forth, while the other players choose x_{-i} . \square

4.2 Voting by veto

The title of this subsection refers to a class of voting procedures; each of them defines a game form. There are finite sets of players N and of outcomes, or alternatives, A . The players may have arbitrary preferences over the outcomes. A voting by veto procedure specifies positive

integer numbers λ_a and μ_i for each $a \in A$ and $i \in N$: μ_i is the number of black balls given to player i ; λ_a is the “veto-resistance” of outcome a . Each player allocates his/her black balls among the outcomes; if the number of balls allocated to $a \in A$ equals or exceeds λ_a , the outcome is vetoed. To ensure the existence of non-vetoed outcomes, we impose the restriction

$$\sum_{a \in A} \lambda_a \geq \sum_{i \in N} \mu_i + 1. \quad (4.4)$$

A linear order is assumed on A ; if several outcomes are not vetoed, the highest of them is selected.

To describe strategies formally, we assume there is a finite set D of black balls with an *ownership mapping* $\nu: D \rightarrow N$. Balls from $D_i = \nu^{-1}(i)$ are given to player i ; we assume $\#D_i = \mu_i$. A strategy of player i is a mapping $x_i: D_i \rightarrow A$; a strategy profile can be understood as a mapping $x_N: D \rightarrow A$. For each $I \in \mathcal{N}$, we denote $D_I = \bigcup_{i \in I} D_i$; every strategy of the coalition can be understood as a mapping $x_I: D_I \rightarrow A$. Given $a \in A$, $I \in \mathcal{N}$, and $x_I \in X_I$, we denote $\varkappa(a, x_I) = \#x_I^{-1}(a)$, the number of balls cast to a by I under x_I . An outcome a is *vetoed* at a strategy profile x_N if $\varkappa(a, x_N) \geq \lambda_a$ and *over-vetoed* if the inequality is strict. $g(x_N)$ is the highest non-vetoed outcome.

For brevity, a game form generated by a voting by veto procedure is called a *VV game form*. Every ordered voting game form from Subsection 3.1 with integer μ 's and λ 's can be represented as the fragment of a VV game form: each player i can choose $a \in A \setminus \{\min A\}$ and put μ_i black balls against a and each outcome above a . Since N and A are finite, it seems plausible that integer μ 's and λ 's are sufficient to generate all ordered voting game forms. Non-integer μ 's and λ 's, apparently, would generate a broader class of VV game forms; however, such voting procedures may seem too exotic, and are outside the realm of finite game forms anyway. Generally, fragments of VV game forms do not possess any nice property: the underlined 2×2 fragment in Example 4.6 is not even Nash consistent.

Remark. If there is $b \in A$ such that $\sum_{a > b} \lambda_a \geq \sum_{i \in N} \mu_i + 1$, then $b \notin g(X_N)$. In a sense, the elimination of such outcomes would not change the game form; however, their presence creates no difficulties either.

Proposition 4.5. *Every VV game form G with $\#g(X_N) \leq 2$ has the FCP.*

Proof. Without restricting generality, we assume $g(X_N) = \{q, a\}$ with $q > a$; then $\lambda_q + \lambda_a > \sum_{i \in N} \mu_i$. For each $I \in \mathcal{N}$ and $x_I, x'_I \in X_I$, we define $x'_I \succeq_I x_I \iff \varkappa(q, x'_I) \geq \varkappa(q, x_I)$. It is clear now that G satisfies (4.2), hence the sufficiency part of Theorem 4.2 applies. \square

Example 4.6. Let $A = \{a, b, c\}$, $a > b > c$, $N = \{1, 2\}$, and every μ and λ be equal to 1. The game form is described by the following matrix:

$$\begin{array}{ccc} \underline{b} & \underline{c} & b \\ \underline{c} & \underline{a} & a \\ b & a & a \end{array}$$

Assuming $v_1(a) = v_1(b) = 1$, $v_1(c) = 0$, $v_2(a) = v_2(b) = 0$, $v_2(c) = 1$, we see that the underlined outcomes form an improvement cycle in $G(v_N)$; therefore, G does not have even the FIP (although it is strongly consistent).

Example 4.6 carries a subtler message as well. We might distinguish between “outcomes” from A and “voting proposals”: since $v_i(a) = v_i(b)$ for both i in the example, we may assume that a and b are just different proposals implying the same outcome, hence $\#A = 2$. Therefore, Proposition 4.5 would not survive this generalization of the notion of voting by veto.

Theorem 4.7. *Let G be a game form with $N = \{1, 2\}$ and $A = \{q, a\}$. Then the following statements are equivalent.*

1. G has the FCP.
2. G is simple.
3. G can be represented as the fragment of a VV game form.

Proof. Suppose Statement 1 holds, hence Theorem 4.2 applies. If G is dictatorial, then both Statements 2 and 3 are obvious. Otherwise, we define an ordering \succeq_i on each X_i by (4.3). We pick x_i^- and x_i^+ (for each $i \in N$) among, respectively, the least and the greatest strategies in X_i w.r.t. \succeq_i . Let us show that \mathcal{X} contains a simple strategy. Pick $i \in N$; if $g(x_i^-, x_{-i}^+) = q$, then $g(x_i^-, x_{-i}) = q$ for all $x_{-i} \in X_{-i}$, and we are home; if $g(x_i^-, x_{-i}^+) = a$, then $g(x_i, x_{-i}^+) = a$ for all $x_i \in X_i$, and we are home again. Thus, G is simple by Proposition 3.2.

To start the derivation of Statement 3, we define $r_i(x_i)$ for $i \in N$ and $x_i \in X_i$ as the rank of x_i w.r.t. \succ_i , hence $r_i(x'_i) = r_i(x_i) \iff x'_i \sim_i x_i$, in particular, $r_i(x_i) = 0 \iff x_i \sim_i x_i^-$. We denote $m_i = r_i(x_i^+)$. The definition (4.3) of \succeq_i implies that there is an increasing mapping $\gamma: [0, m_1] \times [0, m_2] \rightarrow \{0, 1\}$ such that $g(x_N) = a$ if and only if $\gamma(r_1(x_1), r_2(x_2)) = 1$.

Step 4.7.1. *If $i \in N$, $\xi_i \in [0, m_i[$, $\xi_{-i} \in [0, m_{-i}[$, $\gamma(\xi_i, \xi_{-i}) = q$, and $\gamma(\xi_i, \xi_{-i} + 1) = a$, then $\gamma(\xi_i + 1, \xi_{-i}) = a$. If $i \in N$, $\xi_i \in]0, m_i]$, $\xi_{-i} \in [0, m_{-i}[$, $\gamma(\xi_i, \xi_{-i}) = q$, and $\gamma(\xi_i, \xi_{-i} + 1) = a$, then $\gamma(\xi_i - 1, \xi_{-i} + 1) = q$.*

Proof. By (4.3), there must be $\eta_{-i} \in [0, m_{-i}]$ such that $\gamma(\xi_i, \eta_{-i}) = q$ while $\gamma(\xi_i + 1, \eta_{-i}) = a$. Neither $\eta_{-i} < \xi_{-i}$, nor $\eta_{-i} \geq \xi_{-i} + 1$ are compatible with the monotonicity of γ , hence $\eta_{-i} = \xi_{-i}$ and we are home. The proof of the second implication is dual. \square

It is immediately clear from (4.3) that no $\xi_i \in]0, m_i[$ is simple. Denoting $\eta_{-i}(\xi_i) = \min\{\xi_{-i} \mid \gamma(\xi_i, \xi_{-i}) = a\}$, we obtain from Step 4.7.1 that $\xi_i + \eta_{-i}(\xi_i)$ does not depend on ξ_i , nor on i ; we denote the constant λ_q . Clearly, $g(x_N) = a \iff \sum_{i \in N} r_i(x_i) \geq \lambda_q$.

Now we consider a VV game form \bar{G} with the same N , $A = \{q, a, b\}$, $q > a > b$, $\mu_i = m_i + \#X_i$, λ_q just defined, $\lambda_a = \sum_{i \in N} \mu_i + 1$ and an arbitrary λ_b . To represent G as a fragment of \bar{G} , we order each strategy set, i.e., pick an arbitrary bijection $o_i: X_i \rightarrow \{1, \dots, \#X_i\}$; now each strategy $x_i \in X_i$ is interpreted as putting $r_i(x_i)$ balls at q , $o_i(x_i)$ balls at a , and all the rest at b . Clearly, a is never vetoed, hence b is never chosen and a is chosen if and only if q is vetoed. Therefore, g coincides with the restriction of \bar{g} to X_N .

Given Statement 2, we notice that (3.2) implies (4.1); then Proposition 4.1 and Theorem 4.2 imply Statement 1. The implication Statement 3 \Rightarrow Statement 1 immediately follows from Proposition 4.5. \square

Example 4.8. Let $N = \{1, 2, 3\}$, player 1 choose rows, player 2 columns, and player 3 matrices.

$$\begin{bmatrix} q & a \\ q & q \end{bmatrix} \quad \begin{bmatrix} a & a \\ q & a \end{bmatrix}$$

Condition (4.1) obviously holds, but there is no simple strategy. Thus, the implication Statement 1 \Rightarrow Statement 2 from Theorem 4.7 does not hold when $n > 2$ even if $\#A = 2$. The invalidity of the converse implication when $\#A > 2$ even if $n = 2$ is shown by Example 4.6.

It is not difficult to check that the game form is an almost potential one; however, it cannot be represented as a perfect information game form.

Example 4.9. Let $N = \{1, 2, 3\}$, player 1 choose rows, player 2 columns, and player 3 matrices:

$$\begin{bmatrix} q & a & a \\ q & q & q \\ q & q & q \end{bmatrix} \quad \begin{bmatrix} q & a & a \\ q & q & a \\ q & q & a \end{bmatrix} \quad \begin{bmatrix} q & a & a \\ q & a & a \\ q & q & a \end{bmatrix} \quad \begin{bmatrix} a & a & a \\ a & a & a \\ q & q & a \end{bmatrix}.$$

Conditions (4.1) are easy to check, hence the game form is totally tight. Suppose it can be represented as the fragment of a VV game form. Since the roles of a and q are perfectly symmetric, we may, without restricting generality, assume $q > a$; therefore, a is selected when and only when q is vetoed. We denote x_i^s the number of black balls cast at q by player i ($i \in \{1, 2\}$) when using s -th strategy, counting upwards for player 1 and from the left to right for player 2.

The rightmost matrix shows that $x_1^2 - x_1^1 > x_2^2 - x_2^1$; the leftmost, that $x_1^3 - x_1^2 > x_2^3 - x_2^2$; therefore, $x_1^3 - x_1^1 > x_2^3 - x_2^1$. On the other hand, each of the middle matrices shows that $x_2^3 - x_2^1 > x_1^3 - x_1^1$.

Thus, the implication Statement 1 \Rightarrow Statement 3 from Theorem 4.7 does not hold when $n > 2$ even if $\#A = 2$.

Theorem 4.10. *Every VV game form has an almost unrestricted FCP.*

Proof. For every $x_N, y_N \in X_N$, $a \in A$, $t \in D$, and $I \in \mathcal{N}$, we define:

$$\varepsilon(a, x_N) = \max\{\varkappa(a, x_N) - \lambda_a, 0\}; \quad E(x_N) = \sum_{a \in A} \varepsilon(a, x_N)$$

(over-vetoing at a and total over-vetoing);

$$\beta(a, x_N) = \max\{\lambda_a - \varkappa(a, x_N), 0\}$$

(“empty slots” at a);

$$C(x_N) = \{b \in A \mid \sum_{a>b} \beta(a, x_N) < 2 \ \& \ \varepsilon(b, x_N) = 0\} \ni g(x_N); \quad B(x_N) = x_N^{-1}(C(x_N))$$

(the sets of outcomes that are “close enough” to being selected and of the balls cast there);

$$\delta(x_N, y_N; t) = \begin{cases} 0, & \text{if } y_N(t) = x_N(t); \\ +\infty, & \text{if } y_N(t) \neq x_N(t) \ \& \ \varepsilon(y_N(t), y_N) > 0; \\ d(y_N(t), x_N(t)) + 1, & \text{otherwise [with } d \text{ defined by (3.4)];} \end{cases} \quad (4.5a)$$

$$\delta_{x_{-I}}(x_I, y_I) = \sum_{t \in D_I} \delta(x_N, (y_I, x_{-I}); t). \quad (4.5b)$$

For every $I \in \mathcal{N}$, the family of quasidistances on X_I defined in (4.5) generates an admissible change relation \vdash_I , hence an admissible improvement relation \gg in every derivative game. The notion of admissibility combines the ideas from the proofs of Theorems 3.6 and 3.13: the players shift as few balls as possible and move them as short a distance as possible; first of all, however, they avoid over-vetoing at any cost.

Given a list of utilities $\langle v_i \rangle_{i \in N}$, we define

$$P(x_N) = - \sum_{t \in B(x_N)} v_{\nu(t)}(x_N(t));$$

$$y_N \succ x_N \Leftrightarrow [\forall a \in A [\varepsilon(a, x_N) \geq \varepsilon(a, y_N)] \ \& \ (E(x_N) > E(y_N) \ \text{or} \ B(x_N) \subset B(y_N) \ \text{or} \ [B(x_N) = B(y_N) \ \& \ P(x_N) < P(y_N)])]. \quad (4.6)$$

The rest of the proof consists in showing that \succ is an order potential for the admissible improvement relation \triangleright . It seems worthwhile to describe it informally: we monitor the over-vetoing at each outcome, what balls are cast at $C(x_N)$ and who cast them; one strategy profile “dominates” another if either there is a strict decrease in over-vetoing, or more balls come under observation, or the balls are the same but the players try to veto worse (for them) outcomes.

Let $x_N, y_N \in X_N$, $I \in \mathcal{N}$, $y_{-I} = x_{-I}$, $a = g(x_N)$, $b = g(y_N)$, and $D^* = \{t \in D \mid x_N(t) \neq y_N(t)\}$. We have to show that $y_N \succ x_N$ in every derivative game where $y_N \triangleright_I x_N$.

By the definition of g , two things are necessary and sufficient for b to be selected at y_N : b must not be vetoed; everything above b must be vetoed. Formally:

$$\beta(b, y_N) > 0 = \varepsilon(b, y_N); \quad (4.7a)$$

$$\forall c > b [\beta(c, y_N) = 0]. \quad (4.7b)$$

We start with the demonstration of the possibility to switch from a to b without any increase in over-vetoing. If $\beta(b, x_N) = 0$, then $\varepsilon(b, x_N) + 1$ balls must be taken from b to ensure (4.7a); Inequality (4.4) ensures that the total number of “empty slots,” $\sum_{c \in A} \beta(c, x_N)$, is no less than that. Therefore, the members of I can make (4.7a) fulfilled without creating over-vetoing. (Since coalition I was able to make b selected, $\#x_I^{-1}(b)$ must be large enough.) Similarly, no over-vetoing could help ensuring (4.7b). Since y_I is the least cost way to switch from a to b , there must be $\delta_{x_{-I}}(x_I, y_I) < +\infty$. Therefore, there was no increase in over-vetoing at any outcome $c \in A$ when x_N was replaced with y_N , i.e.

$$\forall c \in A [\varepsilon(c, y_N) \leq \varepsilon(c, x_N)]. \quad (4.8)$$

Secondly, if $\varepsilon(b, x_N) > 0$, we have $E(y_N) < E(x_N)$, hence $y_N \succ x_N$ and we are home. The same conclusion is reached if $\varepsilon(c, y_N) < \varepsilon(c, x_N)$ for any $c \in A$. In the following, we assume that

$$\varepsilon(b, x_N) = 0; \quad (4.9a)$$

$$\forall c \in A [\varepsilon(c, y_N) = \varepsilon(c, x_N)]. \quad (4.9b)$$

Thirdly, (4.7) immediately implies $b \in C(y_N)$. The final argument depends on whether $b > a$ or $a > b$.

A. Let $b > a$. By (4.9a), shifting just one ball from b is enough to get that outcome selected; therefore, $D^* = \{t^*\}$ and $x_N(t^*) = b$. The minimality of $\delta(x_N, y_N; t^*)$ implies that the ball goes to the nearest empty slot, which is a , i.e., $y_N(t^*) = a$. It is clear now that $B(y_N) = B(x_N)$ and $P(y_N) - P(x_N) = v_{\nu(t^*)}(b) - v_{\nu(t^*)}(a) > 0$. Therefore, $y_N \succ x_N$ by the last disjunctive term in (4.6).

B. Let $b < a$. We again consider two alternatives. If $b \notin C(x_N)$, then every $c \in C(x_N)$ is above b , hence $C(x_N) \subset C(y_N)$, hence $B(x_N) \subset B(y_N)$ because at least one ball was added at $a \in C(y_N)$. Therefore, $y_N \succ x_N$ by the second disjunctive term in (4.6).

Now let $b \in C(x_N)$; then $\beta(a, x_N) = 1$ and $\beta(c, x_N) = 0$ for every c such that $b < c < a$. Therefore, $D^* = \{t^*\}$, $y_N(t^*) = a$, and $x_N(t^*) \leq b$. If $x_N(t^*) = b$, then $B(y_N) = B(x_N)$ and $P(y_N) - P(x_N) > 0$, hence $y_N \succ x_N$, exactly as in the case of $b > a$. This alternative must hold if $\beta(b, x_N) = 0$. Finally, let $x_N(t^*) = c < b$, hence $\beta(b, x_N) > 0$, hence $c \notin C(x_N)$. We see that $t^* \in B(y_N) \setminus B(x_N)$, hence $y_N \succ x_N$ exactly as in the previous paragraph. \square

Example 4.6 above shows that the adjective “almost unrestricted” in Theorem 4.10 cannot be dropped; actually, both improvements by player 1 there create over-vetoing.

In principle, the notion of voting by veto can be extended by considering other rules for the selection of a single non-vetoed outcome at every strategy profile. Assuming that (4.4) holds as an equality, the existence of a strong equilibrium can be shown for any g (Moulin, 1983). Under a strict inequality, even Nash consistency is not ensured.

Example 4.11. Let $A = \{a, b, c, d\}$, $N = \{1, 2\}$, every μ and λ be equal to 1. It is convenient to assume that each player just chooses an outcome, x_i , to veto. Let g be defined as follows: If $\{x_1, x_2\} \subseteq \{c, d\}$, then $g(x_N) = a$ if $x_1 = x_2$ and $g(x_N) = b$ if $x_1 \neq x_2$; otherwise, $g(x_N) = d$ if d is not vetoed whereas $g(x_N) = c$ if d is vetoed. The rule produces this matrix:

d	d	d	c
d	d	d	c
d	d	a	b
c	c	b	a

Player 1 cannot ensure $g(x_N) \in \{a, c\}$; player 2 cannot ensure $g(x_N) \in \{b, d\}$. Therefore, the game form is not Nash consistent (Gurvich, 1975, 1988; Moulin, 1976).

It remains unclear to what extent Theorem 4.10 could be generalized to other mappings g . So far, an almost unrestricted FCP was only established for a rather peculiar tie-breaking procedure suggested by Gol’berg and Gurvich (1986), which is only defined when (4.4) holds as an equality. Example 2.1d is generated by that procedure.

References

Abdou, J., 1995. Nash and strongly consistent two-player game forms. *International Journal of Game Theory* 24, 345–356.

- Abdou, J., 1998. Tight and effectively rectangular game forms: A Nash solvable class. *Games and Economic Behavior* 23, 1–11.
- Abdou, J., and H. Keiding, 2003. On necessary and sufficient conditions for solvability of game forms. *Mathematical Social Sciences* 46, 243–260.
- Boros, E., and V. Gurvich, 2003. On Nash-solvability in pure stationary strategies of finite games with perfect information which may have cycles. *Mathematical Social Sciences* 46, 207–241.
- Boros, E., V. Gurvich, K. Makino, and W. Shao, 2007. Nash-solvable bidirected cyclic game forms. RUTCOR Research Report RRR 30-2007. Available at http://rutcor.rutgers.edu/pub/rrr/reports2007/30_2007.ps
- Boros, E., V. Gurvich, K. Makino, and D. Papp, 2008a. On acyclic, or totally tight, two-person game forms. RUTCOR Research Report RRR 3-2008. Available at http://rutcor.rutgers.edu/pub/rrr/reports2008/03_2008.pdf
- Boros, E., V. Gurvich, K. Makino, and D. Papp, 2008b. Acyclic, or totally tight, two-person game forms; characterization and main properties. RUTCOR Research Report RRR 6-2008. Available at <http://rutcor.rutgers.edu/pub/rrr/reports2008/paper.pdf>
- Friedman, J.W., and C. Mezzetti, 2001. Learning in games by random sampling. *Journal of Economic Theory* 98, 55–84.
- Germeier, Yu.B., and I.A. Vatel', 1974. On games with a hierarchical vector of interests. *Izvestiya Akademii Nauk SSSR, Tekhnicheskaya Kibernetika*, 3, 54–69 (in Russian; English translation in *Engineering Cybernetics*, V.12).
- Gol'berg, A.I., and V.A. Gurvich, 1986. Collective choice based on the veto principle. VINITI Manuscript #3182–V86 (in Russian).
- Gurvich, V.A., 1975. Solvability of extensive games in pure strategies, *Zhurnal Vychislitel'noi matematiki i matematicheskoi fiziki* 15(2), 358–371 (in Russian; English translation in *USSR Computational Mathematics and Mathematical Physics* 15(2), 74–87).
- Gurvich, V.A., 1988. Equilibrium in pure strategies. *Doklady Akademii Nauk SSSR* 303, 789–793 (in Russian; English translation in *Soviet Mathematics. Doklady* 38, 597–602).
- Holzman, R., and N. Law-yone, 1997. Strong equilibrium in congestion games. *Games and Economic Behavior* 21, 85–101.
- Kalai, E. and D. Schmeidler, 1977. An admissible set occurring in various bargaining situations. *Journal of Economic Theory* 14, 402–411.

Kandori, M., and R. Rob, 1995. Evolution of equilibria in the long run: A general theory and applications. *Journal of Economic Theory* 65, 383–414.

Konishi, H., M. Le Breton, and S. Weber, 1997. Equilibria in a model with partial rivalry. *Journal of Economic Theory* 72, 225–237.

Kreps, D.M., 1990. *A Course in Microeconomic Theory*. Princeton University Press, Princeton.

Kukushkin, N.S., 1995. Two-person game forms guaranteeing the stability against commitment and delaying tactics. *International Journal of Game Theory* 24, 37–48.

Kukushkin, N.S., 1999. Potential games: A purely ordinal approach. *Economics Letters* 64, 279–283.

Kukushkin, N.S., 2000. Potentials for Binary Relations and Systems of Reactions. Russian Academy of Sciences, Computing Center, Moscow

Kukushkin, N.S., 2002a. Perfect information and potential games. *Games and Economic Behavior* 38, 306–317.

Kukushkin, N.S., 2002b. Unfocussed Nash Equilibria. Russian Academy of Sciences, Dorodnicyn Computing Center, Moscow. A revised version available at <http://www.ccas.ru/mmes/mmeda/ququ/Unfocussed.pdf>

Kukushkin, N.S., 2004a. Best response dynamics in finite games with additive aggregation. *Games and Economic Behavior* 48, 94–110.

Kukushkin, N.S., 2004b. Acyclicity of Improvements in Games with Common Intermediate Objectives. Russian Academy of Sciences, Dorodnicyn Computing Center, Moscow. A revised version available at <http://www.ccas.ru/mmes/mmeda/ququ/Common3.pdf>

Kukushkin, N.S., 2006. Acyclicity and Aggregation in Strategic Games. Russian Academy of Sciences, Dorodnicyn Computing Center, Moscow. A version available at <http://www.ccas.ru/mmes/mmeda/ququ/IntAA.pdf>

Kukushkin, N.S., 2007a. Shapley’s “2 by 2” theorem for game forms. *Economics Bulletin* 3, No. 33, 1–5. Available at <http://economicsbulletin.vanderbilt.edu/2007/volume3/EB-07C70017A.pdf>; with a typo corrected, at <http://www.ccas.ru/mmes/mmeda/ququ/Shapley.pdf>

Kukushkin, N.S., 2007b. Congestion games revisited. *International Journal of Game Theory* 36, 57–83.

Kukushkin, N.S., 2007c. Aggregation and Acyclicity in Strategic Games. Russian Academy of Sciences, Dorodnicyn Computing Center, Moscow. A version available at <http://www.ccas.ru/mmes/mmeda/ququ/AgrAcl.pdf>

- Kukushkin, N.S., S. Takahashi, and T. Yamamori, 2005. Improvement dynamics in games with strategic complementarities. *International Journal of Game Theory* 33, 229–238.
- Mariotti, M., 2000. Maximum games, dominance solvability, and coordination. *Games and Economic Behavior* 31, 97–105.
- Milchtaich, I., 1996. Congestion games with player-specific payoff functions. *Games and Economic Behavior* 13, 111–124.
- Milgrom, P., and J. Roberts, 1991. Adaptive and sophisticated learning in normal form games. *Games and Economic Behavior* 3, 82–100.
- Monderer, D., and L.S. Shapley, 1996. Potential games. *Games and Economic Behavior* 14, 124–143.
- Moulin, H., 1976. Prolongements des jeux à deux joueurs de somme nulle. *Bulletin de la Société Mathématique de France, Supplémentaire Mémoire No. 45*.
- Moulin, H., 1980. On strategy-proofness and single-peakedness. *Public Choice* 35, 437–455.
- Moulin, H., 1983. *The Strategy of Social Choice*. North-Holland Publ., Amsterdam.
- Moulin, H., 1984. Dominance solvability and Cournot stability. *Mathematical Social Sciences* 7, 83–102.
- Mueller, D.C., 1978. Voting by veto. *Journal of Public Economics* 10, 57–75.
- Peleg, B., 1978. Consistent voting systems. *Econometrica* 46, 153–161.
- Rosenthal, R.W., 1973. A class of games possessing pure-strategy Nash equilibria. *International Journal of Game Theory* 2, 65–67.
- Sela, A., 1992. *Learning Processes in Games*. M.Sc. Thesis. The Technion, Haifa (in Hebrew).
- Topkis, D.M., 1979. Equilibrium points in nonzero-sum n-person submodular games. *SIAM Journal on Control and Optimization* 17, 773–787.
- Topkis, D.M., 1998. *Supermodularity and Complementarity*. Princeton University Press, Princeton.
- Vives, X., 1990. Nash equilibrium with strategic complementarities. *Journal of Mathematical Economics* 19, 305–321.
- Young, H.P., 1993. The evolution of conventions. *Econometrica* 61, 57–84.