



Munich Personal RePEc Archive

Martingales, Detrending Data, and the Efficient Market Hypothesis

McCauley, Joseph L. and Bassler, Kevin E. and Gunaratne,
Gemunu H.

University of Houston, NUI Galway

February 2007

Online at <https://mpra.ub.uni-muenchen.de/2256/>

MPRA Paper No. 2256, posted 15 Mar 2007 UTC

Martingales, Detrending Data, and the Efficient Market Hypothesis

Joseph L. McCauley⁺, Kevin E. Bassler⁺⁺, and Gemunu H. Gunaratne⁺⁺⁺

Physics Department
University of Houston
Houston, Tx. 77204-5005
jmccauley@uh.edu

⁺Senior Fellow
COBERA
Department of Economics
J.E.Cairnes Graduate School of Business and Public Policy
NUI Galway, Ireland

⁺⁺Texas Center for Superconductivity
University of Houston
Houston, Texas 77204-5005

⁺⁺⁺Institute of Fundamental Studies
Kandy, Sri Lanka

Key Words: Martingales, Markov processes, detrending, memory, stationary and nonstationary increments, correlations, efficient market hypothesis.

Abstract

We discuss martingales, detrending data, and the efficient market hypothesis for stochastic processes $x(t)$ with arbitrary diffusion coefficients $D(x,t)$. Beginning with x -independent drift coefficients $R(t)$ we show that Martingale stochastic

processes generate uncorrelated, generally *nonstationary* increments. Generally, a test for a martingale is therefore a test for uncorrelated increments. A detrended process with an x -dependent drift coefficient is generally not a martingale, and so we extend our analysis to include the class of (x,t) -dependent drift coefficients of interest in finance. We explain why martingales look Markovian at the level of both simple averages *and* 2-point correlations. And while a Markovian market has no memory to exploit and presumably cannot be beaten systematically, it has never been shown that martingale memory cannot be exploited in 3-point or higher correlations to beat the market. We generalize our Markov scaling solutions presented earlier, and also generalize the martingale formulation of the efficient market hypothesis (EMH) to include (x,t) -dependent drift in log returns. We also use the analysis of this paper to correct a misstatement of the 'fair game' condition in terms of serial correlations in Fama's paper on the EMH. We end with a discussion of Levy's characterization of Brownian motion and prove that an arbitrary martingale is topologically inequivalent to a Wiener process.

1. Introduction

Recently [1] we focused on the condition for long time correlations like and including fractional Brownian motion (fBm), which is stationarity of the increments in a stochastic process $x(t)$ with variance nonlinear in the time. There, we derived the 2-point and 1-point densities including the transition density for fBm. We will point out below that there are nonMarkov systems where the pair correlations cannot be distinguished from those of a Markov process, but time series with stationary increments (like fBm) exhibit long time memory that can be seen at the level of pair

correlations: fBm cannot be mistaken for a Markov process at the 2-point level. We correspondingly emphasized that neither 1-point averages nor Hurst exponents can be used to identify the presence or absence of history-dependence in a time series, or to identify the underlying stochastic process (see [2] for the conclusion that an equation of motion for a 1-point density cannot be used to decide if a process is Markovian or not). In the same paper, we pointed out that the opposite class, systems with no memory at all (Markov processes) and with x -independent drift coefficients generate uncorrelated, typically nonstationary increments. The conclusions in [1] about Markov processes are more general than we realized at the time. Here, we generalize that work by focusing on martingales.

In applications to finance, by “ x ” we always mean $x(t)=\ln(p(t)/p_c)$ where $p(t)$ is a price at time t and p_c is a reference price, the consensus price or ‘value’ [3]. The consensus price p_c is simply the price that determines the peak of the 1-point returns density $f_1(x,t)$. The reason why *log increments* $x(t;T)=\ln p(t+T)/p(t)$ and price differences $\Delta p=p(t+T)-p(t)$ generally *cannot* be taken as ‘good’ variables describing a stochastic process (neither theoretically nor in data analysis) is explained below in part 4. It is impossible for a martingale, excepting the special case of a variance linear in the time t , to develop either stochastic dynamics or probability theory based on *increments* $x(t;T)$ or Δp , because if the increments are nonstationary, as they generally are, then the starting time t matters and consequently histograms derived empirically from time series assuming that the starting time doesn’t matter exhibit ‘significant artifacts’ like fat tails and spurious Hurst exponents [3,4]. In contrast, in a system with long time autocorrelations (like fBm), the stationary increment $x(t;T)=x(t+T)-x(t)=x(T)$, ‘in distribution’, is a perfectly good variable. But real markets [4], rule out such long time autocorrelations. Stated briefly, if

increments are nonstationary then the 1-point density that describes the increments is not independent of t (see part 5 below).

Next, we define the required underlying ideas.

2. Conditional expectations with memory

Imagine a collection of time series generated by an unknown stochastic process that we would like to discover via data analysis. Simple averages require only a 1-point density $f_1(x,t)$, e.g., $\langle x^n(t) \rangle = \int x^n f_1(x,t) dx$. No dynamical process can be identified by specifying merely either the 1-point density or a scaling exponent [1]. Both conditioned and unconditioned two-point correlations, e.g., $\langle x(t)x(t+T) \rangle = \int dy dx y x f_2(y,t+T;x,t)$, require a two point density $f_2(y,t+T;x,t)$ for their description and provide us with limited information about the class of dynamics under consideration.

Consider a collection of time series representing repeated runs of a single stochastic process $x(t)$. Empirically, we can only strobe the system finitely many times, so measurements of $x(t)$ take the form of $\{x(t_k)\}$, $k=1,\dots,n$ where n is the number of measurements/observations made in one run. Many repeated runs are required in order to get histograms reflecting the statistics of the process. If we can extract good enough histograms from the collection of time series (if there are many runs, and if each run contains enough points), then we can then try to extract the hierarchy of probability densities $f_1(x,t)$, $f_2(x_2,t_2;x_1,t_1)$, ..., $f_k(x_k,t_k;\dots,x_1,t_1)$ where $k \ll n$ (where f_1 implicitly reflects a specific choice of initial condition in data analysis). To get adequate histograms for f_n one would then need a much longer time series. If the memory in the process is discrete in size, then the minimum n number of densities that one needs in the hierarchy

depends on the length N of the memory sequence in the system (for a Markov process, $N=2$). In what follows $f_n(x_n, t_n; \dots; x_1, t_1)$ denotes the probability density for the sequence (x_n, \dots, x_1) at observation times (t_n, \dots, t_1) , where we generally take $t_1 < \dots < t_n$.

Conditional probability densities p_k or transition probability densities, can then be defined as [5,6]:

$$f_2(x_2, t_2; x_1, t_1) = p_2(x_2, t_2 | x_1, t_1) f_1(x_1, t_1), \quad (1)$$

$$f_3(x_3, t_3; x_2, t_2; x_1, t_1) = p_3(x_3, t_3 | x_2, t_2, x_1, t_1) p_2(x_2, t_2 | x_1, t_1) f_1(x_1, t_1), \quad (2)$$

and more generally as

$$f_n(x_n, t_n; \dots; x_1, t_1) = p_n(x_n, t_n | x_{n-1}, t_{n-1}; \dots; x_1, t_1) \dots p_2(x_2, t_2 | x_1, t_1) f_1(x_1, t_1), \quad (3)$$

where p_n is the 2-point conditional probability density to find x_n at time t_n , given the last observed point (x_{n-1}, t_{n-1}) and the previous history $(x_{n-2}, t_{n-2}; \dots; x_1, t_1)$. When memory is present in the system then one cannot use the simplest 2-point transition density p_2 to describe the complete time evolution of the dynamical system that generates $x(t)$.

In a Markov process the picture is much simpler. A Markov process [5,6] has no memory aside from the last observed point in the time series. In this case one loosely says that the system has no memory. There, we have

$$f_n(x_n, t_n; \dots; x_1, t_1) = p_2(x_n, t_n | x_{n-1}, t_{n-1}) \dots p_2(x_2, t_2 | x_1, t_1) f_1(x_1, t_1), \quad (4)$$

because all transition rates p_n , $n>2$, are built up as products of p_2 ,

$$p_k(x_k, t_k | x_{k-1}, t_{k-1}; \dots; x_1, t_1) = p_2(x_k, t_k | x_{k-1}, t_{k-1}), \quad (5)$$

for $k=3,4, \dots$, and so p_2 cannot depend on an initial state (x_1, t_1) or on any previous state other than the last observed point (x_{k-1}, t_{k-1}) . Only in the absence of memory does the 2-point density p_2 describe the complete time evolution of the dynamical system. E.g., we can prove that for an arbitrary process with or without memory

$$p_{k-1}(x_k, t_k | x_{k-2}, t_{k-2}; \dots; x_1, t_1) = \int dx_{k-1} p_k(x_k, t_k | x_{k-1}, t_{k-1}; \dots; x_1, t_1) p_{k-1}(x_{k-1}, t_{k-1} | x_{k-2}, t_{k-2}; \dots; x_1, t_1) \quad (6)$$

and therefore that

$$p_2(x_3, t_3 | x_1, t_1) = \int dx_2 p_3(x_3, t_3 | x_2, t_2; x_1, t_1) p_2(x_2, t_2 | x_1, t_1), \quad (7)$$

whereas the Chapman-Kolmogorov (CK) equation for a Markov process follows with $p_n=p_2$ for $n=2,3,\dots$, from (6) so that

$$p_2(x_3, t_3 | x_1, t_1) = \int dx_2 p_2(x_3, t_3 | x_2, t_2) p_2(x_2, t_2 | x_1, t_1). \quad (8)$$

The Markov property is expressed by $p_n=p_2$ for all $n \geq 3$, the complete lack of memory excepting the last observed point. *The CK Equation (8) is a necessary but not sufficient condition for a Markov process [7,8].*

A Markov process apparently defines a 1-parameter semi-group $U(t_2, t_1)$ of transformations [10], the semi-group

property is easy to prove for time translationally invariant systems (meaning that $p_2(x_n, t_n; x_{n-1}, t_{n-1}) = p_2(x_n, t_n - t_{n-1}; x_{n-1}, 0)$), but time translational invariance is not a property of FX data [4] and will not be assumed here. In any case, the group combination law is given by the CK eqn. (8), which easily can be used to prove associativity. Associativity expresses path independence of any sequence transformations. The identity element is defined by the equal times transition density

$$p_2(y, t | x, t) = \delta(y - x). \quad (9)$$

Processes with memory generally do obey the CK eqn. Instead, the class of path-dependent time evolutions is defined by the entire hierarchy eqns. (3,6), for $n=2,3,4,\dots$. That the transition density for fBm, e.g., obeys no CK eqn. is shown implicitly in Appendix B of [11], where the authors show that for general Gaussian processes one obtains the semi-group property iff. the Gaussian describes a Markov process. Without the CK eqn. one cannot derive a Fokker-Planck pde for the transition density from a Kramers-Moyal expansion [5].

Memory-dependent processes in statistical physics have been discussed by Hänggi and Thomas [11]. They point out that

$$p_2(x_3, t_3 | x_2, t_2) = \frac{\int dx_1 p_3(x_3, t_3 | x_2, t_2; x_1, t_1) p_2(x_2, t_2 | x_1, t_1) f_1(x_1, t_1) dx_1}{\int p_2(x_2, t_2 | x_1, t_1) f_1(x_1, t_1) dx_1} \quad (10)$$

is a functional of the initial state $f_1(x_1, t_1)$ in which the system was prepared at the initial time t_1 , unless the process is Markovian. In a nonMarkov system one can superficially hide this dependence on state preparation by choosing the

initial condition to be $f_1(x_1, t_1) = \delta(x_1)$ (that initial condition is inherent in the standard definition of fBm with initial time $t_1 = -\infty$ [1]). If, instead, we would or could choose $f_1(x_1, t_1) = \delta(x_1 - x'_0)$ at $t_1 = 0$, e.g., then we obtain $p_2(x_3, t_3; x_2, t_2) = p_3(x_3, t_3; x_2, t_2, x'_0)$, introducing a dependence on x'_0 in both the drift and diffusion coefficients. So in this case, what appears superficially as p_2 is really a special case of p_3 . The authors of [11] point out that the origin of memory in statistical physics is often a consequence of averaging over other, slowly changing, variables. We will return to this point in the section below on the efficient market hypothesis.

Systems with memory may lack translational invariance in x and/or time t , but there are drift-free Markov systems that lack translational invariance in both x and t because of nonstationary increments arising from an (x, t) dependent diffusion coefficient [1]. Next, we exhibit a more general class of Markov systems that break both 'space and time' translational invariance than those with Hurst exponent scaling of the 1-point density $f_1(x, t)$ and the diffusion coefficient $D(x, t)$ discussed earlier in ref. [1,9]. In general, scaling of the 1-point density f_1 does not yield scaling of either f_n or p_n for $n \geq 2$ (see ref. [1,9] for examples, both nonMarkovian and Markovian).

A class of Markov scaling solutions with scaling more general than Hurst exponent scaling [1,3,9], is given as follows: let

$$f_1(x, t) = \sigma_1^{-1}(t) F(u) \quad (11)$$

with initial condition $f_1(x, 0) = \delta(x)$, where $u = x / \sigma_1(t)$, with variance

$$\sigma^2(t) = \langle x^2(t) \rangle = \sigma_1^2(t) \langle x^2(1) \rangle. \quad (12)$$

Then with the diffusion coefficient scaling as

$$D(x,t) = (d\sigma_1^2 / dt)\bar{D}(u) \quad (13)$$

where $d\sigma_1/dt > 0$ is required, $f_1(x,t)$ satisfies the Fokker-Planck pde

$$\frac{\partial f_1}{\partial t} = \frac{1}{2} \frac{\partial^2 (Df_1)}{\partial x^2} \quad (14)$$

and yields the scale invariant part of the solution

$$F(u) = \frac{C}{\bar{D}(u)} e^{-\int u du / \bar{D}(u)}. \quad (15)$$

An example is given by Hurst exponent scaling $\sigma_1(t) = t^H$, $0 < H < 1$. A piecewise constant drift $R(t)$ can be included in our result by replacing x by $x - \int R(s) ds$ in u [1,9].

The Green function $g(x,t;x_o,t_o)$ of (14) for an arbitrary initial condition $(x_o,t_o) \neq (0,0)$ does not scale [9], but then the 2-point transition density $p_2(x_2,t_2; x_1,t_1)$ for fBm does not scale either, reflecting as it does an arbitrary point in a time series $(x_1,t_1) \neq (x_o,t_o) = (0,-\infty)$. In all cases scaling, when it occurs, can only be seen in the special choice of conditional density $f_1(x,t) = p_2(x,t;0,t_o)$ with $t_o = 0$ for a Markov process, and $t_o = -\infty$ for fBm.

The same 1-point density $f_1(x,t)$ may describe nonMarkovian processes because a 1-point density taken alone, without the information provided by the transition densities, defines no specific stochastic process and may be generated by many different completely unrelated processes, including systems

with long time increment autocorrelations like fBm [1]. We will show below that a 2-point density delineates fBM from a martingale, but that pair correlations cannot be used to distinguish an arbitrary martingale from a drift-free Markov process.

Finally, note also that

$$\begin{aligned} f_{n-1}(x_{n-1}, t_{n-1}; \dots; x_1, t_1) &= \int dx_n f_n(x_n, t_n; \dots; x_1, t_1) \\ &= \int dx_n p_n(x_n, t_n | x_{n-1}, t_{n-1}; \dots; x_1, t_1) f_{n-1}(x_{n-1}, t_{n-1}; \dots; x_1, t_1) \end{aligned} \quad (16)$$

so that

$$\int dx_n p_n(x_n, t_n | x_{n-1}, t_{n-1}; \dots; x_1, t_1) = 1. \quad (17)$$

3. Absence of trend and martingales

By a trend, we mean that $d\langle x(t) \rangle / dt \neq 0$, conversely, by lack of trend we mean that $d\langle x(t) \rangle / dt = 0$. If a stochastic process can be detrended, then $d\langle x \rangle / dt = 0$ is possible via a transformation of variables but one must generally specify which average is used to define $\langle x \rangle$. If the drift coefficient $R(x, t)$ depends on x , then detrending with respect to a specific average generally will not produce a detrended series if a different average is then used (e.g., one can choose different conditional averages, or an absolute average). To push this problem under the rug until the last section of the paper, we restrict in what follows to processes that can be detrended once and for all by a simple subtraction. I.e., we assume for the time being a trivial drift coefficient but allow for nontrivial diffusion coefficients. This case is of interest both theoretically and for FX data analysis.

A trivial drift coefficient $R(t)$ is a function of time alone. A nontrivial drift coefficient $R(x,t)$ depends on x , on (x,t) , or on (x,t) plus memory $\{x\}$, and is defined for Ito processes by

$$R(x,t,\{x\}) \approx \frac{1}{T} \int_{-\infty}^{\infty} dy (y-x) p_n(y, t+T; x, t, \{x\}) \quad (18)$$

as T vanishes, where $\{x\}$ denotes the history dependence in p_n , e.g. with $y=x_n$ and $x=x_{n-1}$ $(y, t+T, x, t; \{x\})$ denotes $(x_n, t_n; x_{n-1}, t_{n-1}, x_{n-2}, t_{n-2}, \dots, x_1, t_1)$ with $y=x_n$ and $x=x_{n-1}$. If $R(x,t)=0$ then

$$\int_{-\infty}^{\infty} dy y p_n(y, t+T; x, t, \{x\}) = x, \quad (19)$$

so that the conditional average over x at a later time is given by the last observed point in the time series, $\langle x(t+T) \rangle_{\text{cond}} = x(t)$. This is the notion of a fair game: there is no systematic change in x on the average as t increases, $d\langle x(t) \rangle_{\text{cond}} / dt = 0$. The process $x(t)$ is generally nonstationary, and the condition (19) is called a *local martingale* [12]. *The possibility of vanishing trend, $d\langle x \rangle / dt = 0$, implies a local martingale $x(t)$, and vice-versa.*

This is essentially the content of the Martingale Representation Theorem [13], which states that an arbitrary martingale can be built up from a Wiener process $B(t)$, the most fundamental martingale, via stochastic integration ala Ito,

$$x(t) = \int b(x(s), s; \{x\}) dB(s). \quad (20)$$

There is no drift term in (20), in the stochastic differential equation (sde)

$$dx(t) = b(x(t), t; \{x\})dB(t) \quad (21)$$

the diffusion coefficient,

$$D(x, t, \{x\}) \approx \frac{1}{T} \int dy (y - x)^2 p_n(y, t | x, t - T, \{x\}) \quad (22)$$

as T vanishes, is given by $D=b^2$. In a Markov system the drift and diffusion coefficients depend on (x,t) alone, have no history dependence. Ito calculus based on martingales has been developed systematically by Durrett, including the derivation of Girsanov's Theorem for arbitrary diffusion coefficients $D(x,t)$ [12]. Many discussions of Girsanov's Theorem [13,14] implicitly rule out the general case (19) where $D(x,t)$ may depend on x as well as t . In this paper we do not appeal to Girsanov's theorem because the emphasis is on application to data analysis, to detecting martingales in empirical data. A new and simplified proof of Girsanov's theorem for variable diffusion coefficients will be presented elsewhere [15].

It's quite easy to write down a diffusion coefficient with memory of the initial state. For a Markov process the scaling processes follow from assuming that

$$D(x, t) = |t|^{2H-1} \bar{D}(u), u = |x|/|t|^H \quad (22b)$$

and exhibit no history. If, however, we should instead write

$$D(x - x_0, t - t_0) = |t - t_0|^{2H-1} \bar{D}(u), u = |x - x_0|/|t - t_0|^H \quad (22c)$$

then by (10) above this follows from (22) with $n=3$, where (x_0, t_0) is the initial state in which the system was prepared. As we pointed out above, variable diffusion systems are not translation invariant, and neither is FX data [4]. To

emphasize the point, (22c) generates a nonMarkovian martingale.

There are stochastic processes that are inherently biased, and fBm provides an example. There, although the absolute average vanishes $\langle x(t) \rangle = \int \langle x \rangle_c f_1(x,t) dx = 0$, the conditional average yields $d\langle x \rangle_c / dt \neq 0$ where the time dependence arises from long time correlations rather from a drift term: in fBm one obtains [1]

$$\langle x(t) \rangle_{\text{cond}} = \int dy p_2(x, s | x, t) = C(t, s)x \quad (23)$$

instead of the martingale condition (19). Here, $d\langle x \rangle_c / dt = x dC / dt \neq 0$ because the factor $C(t, s) \neq 1$ is proportional to the autocorrelation function $\langle x(s)x(t) \rangle$ where the stationarity of increments guaranteeing long time memory was built in [1]. Such processes cannot be 'detrended' ($R(x, t) = 0$ by construction in fBm [1,16]) because what appears locally to be a trend in a conditional average is simply the strongly correlated behavior of the entire time series.

Note next that subtracting an average drift $\int \langle R \rangle dt$ from a process $x(t)$ defined by x -dependent drift term plus a Martingale,

$$x(t) = x(t-T) + \int_{t-T}^t R(x(s), s; \{x\}) ds + \int b(x(s), s; \{x\}) dB(s), \quad (24)$$

does not produce a martingale. Here, if we replace $x(t)$ by $x(t) - \int \langle R \rangle dt$ where the average drift term defined conditionally from some initial condition (x_1, t_1) ,

$$\langle R \rangle = \int dx R(x, t) p_2(x, t | x_1, t_1) \quad (25)$$

depends on t alone we do not get drift free motion, and choosing absolute or other averages of R will not change this. In financial analysis, e.g., $\langle R \rangle$ may represent an average from the opening return x_1 at opening time t_1 up to some arbitrary intraday return x at time t . The subtraction yields

$$x(t) = x(t-T) + \int_{t-T}^t R(x(s), s; \{x\}) ds - \int_{t-T}^t \langle R \rangle ds + \int b(x(s), s; \{x\}) dB(s) \quad (26)$$

and is not a martingale unless R is independent of x : we obtain $\langle x(t) \rangle_c = x$ iff. $x = x_0$. The general problem of an (x, t) dependent drift $R(x, t)$ in financial applications will be discussed in the last section of this paper.

Again, in what follows we assume a trivial drift $R(t)$ that has been subtracted, so that by $x(t)$ we really mean $x(t) - \int R(t) dt$.

So we can, for our present purposes, divide stochastic processes into those that satisfy the martingale condition

$$\langle x(t) \rangle_{\text{cond}} = x(t_0), \quad (27)$$

where $\langle \dots \rangle_{\text{cond}}$ denotes the conditional average (19), and those that do not. Those that do not satisfy (19) can be classified further into processes that consist of a nontrivial (i.e., (x, t) -dependent) drift plus a martingale (20), and those (including fBm) that are not defined by an underlying martingale.

Summarizing the idea of a martingale, given any set of n points in a time series, $\{x(t_k)\}$, $k=1, \dots, n$, where $t_n > t_{n-1} > \dots > t_2 > t_1$ and the hierarchy of transition densities p_n , the idea of a Martingale is that the best systematic forecast of the future [17] is the conditional average $\langle x(t_k) \rangle_{\text{cond}} = x(t_{k-1})$. I.e., our

expectation of the future is determined by the last observed point in the time series,

$$\int x_n p_n(x_n, t_n | x_{n-1}, t_{n-1}, \dots, x_1, t_1) dx_n = x_{n-1}, \quad (28)$$

all previous observations (x_{n-1}, \dots, x_1) don't contribute. This feature makes a martingale as near as possible to a drift-free Markov process without eliminating the possibility of memory. The conditions that must be satisfied in order that a martingale follows are derived in the next section. The point here is that at the level of simple averages the history dependence cannot be detected. We will show in the next section that the history dependence also cannot appear in pair correlations, making any history in a martingale hard to detect empirically. We understand the condition for a local martingale (19) as the condition that bias-free motion occurs.

The simplest, best known example of a martingale is a drift-free Markov process, where there is no memory at all, i.e., where D depends on (x, t) *alone* completely independent of any and all history simply because (see eqn. (22) above) the transition density $p_2(y, s; x, t)$ depends on the one, single past state (x, t) alone, and on no other earlier states.

Finally, the Ito sde (22) with or without drift included can be used to derive a Fokker-Planck pde for a stochastic process with memory. The Fokker-Planck pde is usually derived from the CK eqn. for a Markov process as an approximation, but this is not necessary. The derivation of the Fokker-Planck pde from the Ito sde, without assuming a C-K eqn. a priori, is provided in [18,19] goes through even if the drift and diffusion coefficients R and D are memory dependent. In that case one has a pde for a 2-point conditional probability p_n depending on a history of $n-2$ earlier states. The

derivation is given in the Appendix for the benefit of the reader.

4. Stationary vs. nonstationary increments

Let us preface this section with a comment: in contrast with what is assumed in the econophysics and finance literature, we know of only two stochastic processes with both finite variance and stationary increments: the Wiener process and fractional Brownian motion. Furthermore, we know of no finance data with stationary increments. Furthermore, stationary increments is a very different condition than either time or space translational invariance. Nonstationary increments are ubiquitous in both theory and data analysis.

In this section we generalize an argument in [1] that assumed Markov processes with trivially removable drift $R(t)$. In fact, that argument was based on no specifically Markovian assumption and applies quite generally to nonMarkovian martingales. In the analysis that follows, we assume a drift-free nonstationary process $x(t)$ with the initial condition $x(t_0)=0$, so that the variance is given by $\sigma^2=\langle x^2(t)\rangle=\int x^2 f_1(x,t)$. By the increments of the process we mean $x(t;T) = x(t+T)-x(t)$ and $x(t;-T)=x(t)-x(t-T)$.

We state in advance that we assume that $[-\infty < x < \infty]$, that there are no boundary conditions that would lead to statistical equilibrium. All processes considered are nonstationary ones.

Stationary increments are defined by

$$x(t + T) - x(t) = x(T), \quad (29)$$

‘in distribution’, and by nonstationary increments [1,3,4,5] we mean that

$$x(t + T) - x(t) \neq x(T). \quad (30)$$

in distribution. When (29) holds, then given the density of ‘positions’ $f_1(x,t)$, we also know the density $f_1(x(T),T)=f_1(x(t+T)-x(t),T)$ of increments independently of the starting time t . Whenever the increments are nonstationary then any analysis of the increments inherently requires the two-point density, $f_2(x(t+T),t+T;x(t),t)$. From the standpoint of theory there exists no 1-point density of increments $f(x;T),T$ depending on T alone, independent of t , and spurious 1-point histograms of increments are typically constructed empirically by assuming that the converse is possible [4]. Next, we place an important restriction on the class of stochastic processes under consideration.

According to Mandelbrot, so-called ‘efficient market’ has no memory that can be *easily* exploited in trading [17]. Beginning with that idea we can assert the necessary but not sufficient condition, the absence of increment autocorrelations,

$$\langle (x(t_1) - x(t_1 - T_1))(x(t_2 + T_2) - x(t_2)) \rangle = 0, \quad (31)$$

when there is no time interval overlap, $t_1 < t_2$ and $T_1, T_2 > 0$. This is a much weaker condition and far more interesting than asserting that the increments are statistically independent. We will see that this condition leaves the question of the dynamics of $x(t)$ open, except to rule out processes with increment autocorrelations, specifically stationary increment processes like fBm [1,20], but also processes with correlated nonstationary increments like the

time translationally invariant Gaussian transition densities described in [2].

Consider a stochastic process $x(t)$ where the increments (31) are uncorrelated. From this condition we easily obtain the autocorrelation function for positions (returns), sometimes called 'serial autocorrelations'. If $t > s$ then

$$\langle x(t)x(s) \rangle = \langle (x(t) - x(s))x(s) \rangle + \langle x^2(s) \rangle = \langle x^2(s) \rangle > 0, \quad (32)$$

since with $x(t_0)=0$ $x(s)-x(t_0)=x(s)$, so that $\langle x(s)x(t) \rangle = \langle x^2(s) \rangle$ is simply the variance in x . Given a history $(x(t), \dots, x(s), \dots, x(0))$, or $(x(t_n), \dots, x(t_k), \dots, x(t_1))$, (32) reflects a martingale property:

$$\begin{aligned} \langle x(t_n)x(t_k) \rangle &= \int dx_n \dots dx_1 x_n x_k p_n(x_n, t_n | x_n, t_n, \dots, x_n, t_n, \dots) p_{n-1}(\dots) \dots p_{k+1}(\dots) f_k(\dots) \\ &= \int x_k^2 f_k(x_k, t_k; \dots; x_1, t_1) dx_k \dots dx_1 = \int x^2 f_1(x, t) dx = \langle x_k^2(t_k) \rangle \end{aligned} \quad (33)$$

where

$$\int x_m dx_m p_m(x_m, t_m | x_{m-1}, t_{m-1}; \dots; x_1, t_1) = x_{m-1}. \quad (34)$$

*Every martingale generates uncorrelated increments and conversely, and so for a Martingale $\langle x(t)x(s) \rangle = \langle x^2(s) \rangle$ if $s < t$.*¹

In a martingale process, the history dependence cannot be detected at the level of 2-point correlations, memory effects can at best first appear at the level 3-point correlations

¹ Note that (32,33) hold for time translationally invariant martingales, where $p_2(x, t; y, s) = p_2(x, t-s; y, 0)$. One can easily check this for a drift-free Gaussian Markov process. I.e., time translational invariance does not imply that $\langle x(t)x(s) \rangle$ is a function of $t-s$ alone. Time translational invariance of p_n , $n \geq 2$, does not imply that a statistical equilibrium density $f_1(x)$ exists and is approached asymptotically by $f_1(x, t)$ [21]. I.e., a time translationally invariant martingale on $[-\infty, \infty]$ cannot yield a stationary process, cannot lead to statistical equilibrium.

requiring the study of a transition density p_3 . Here, we have not postulated a martingale, instead we've deduced that property from the lack of pair wise increment correlations. But this is only part of the story. What follows next is crucial for avoiding mistakes in data analysis [4].

Combining

$$\langle (x(t+T) - x(t))^2 \rangle = \langle x^2(t+T) \rangle + \langle x^2(t) \rangle - 2\langle x(t+T)x(t) \rangle \quad (35)$$

with (34), we get

$$\langle (x(t+T) - x(t))^2 \rangle = \langle x^2(t+T) \rangle - \langle x^2(t) \rangle \quad (36)$$

which depends on *both* t and T , excepting the case where $\langle x^2(t) \rangle$ is linear in t . Uncorrelated increments are generally nonstationary. ***Therefore, martingales generate uncorrelated, typically nonstationary increments.*** So, at the level of pair correlations a martingale with memory cannot be distinguished empirically from a drift-free Markov process. *To see the memory in a martingale one must study at the very least the 3-point correlations.* The increments of a martingale may be stationary iff. the variance is linear in t (we restrict ourselves to the consideration of processes with finite variance). For $H=1/2$ the Ito integral eqn. for a scaling martingale x yields (see eqn. (26') in [9])

$$x(t+T) - x(t) = \int_0^T \sqrt{D(x(s+t)/|s+t|^{1/2})} dB(s)$$

where stationarity of increments of $B(t)$ was used. So although the t -dependence disappears from the average (36) when $H=1/2$, we cannot prove that it disappears for all moments $\langle (x(t+T)-x(t))^n \rangle$. I.e., we cannot prove from Ito

calculus alone that scaling martingales with $H=1/2$ have stationary increments, although simulations for the exponential process indicate stationary increments [4].

We've emphasized earlier [1] that stationary increments $x(t,T)=x(t+T)-x(t)=x(T)$ with finite variance $\langle x^2(t) \rangle < \infty$ generate the long time increment autocorrelations characteristic of fBm [1,16,20], whereas stationary uncorrelated increments with infinite variance occur in Levy processes [22,23]. Stationary Gaussian processes with correlated nonstationary increments are constructed in [2].

A martingale $x(t)$ has no drift, and conditioned on the return $x(t_0)$ yields $\langle x(t) \rangle_{\text{cond}} = x(t_0)$. That is, $x(t)$ not only has no trend but the conditional average is in addition 'stuck' at the last observed point in a time series,

$$\int x_n p_n(x_n, t_n | x_{n-1}, t_{n-1}; \dots; x_1, t_1) dx_n = x_{n-1}. \quad (37)$$

Since $x(t)$ represents the return or 'gain', one further toss of the coin produces no expected gain.

Summarizing, we've shown explicitly that fBm is not a martingale [1], while every Markov process with trivial drift $R(t)$ can be transformed into a (local) Martingale via the substitution of $x(t) - \int R dt$ for $x(t)$: Ito sdes with vanishing drift describe local martingales [12]. A martingale may have memory, and we've provided a model diffusion coefficient to illustrate the appearance of memory (any drift or diffusion coefficient depending on a state (x', t') other than the present state (x, t) exhibits memory, so diffusive models with memory are quite easy to construct). We've shown that uncorrelated increments are nonstationary unless the variance is linear in t . This means that looking for memory in two point correlations is useless: at that level of description a martingale with memory will look Markovian. To find the

memory in a martingale one must study the transition densities p_n and correlations for $n \geq 3$. This has not been discussed in the literature, so far as we know.

As a preliminary step to discussing the EMH, consider a Martingale process $x(t)$. The best forecast of any later return is the expected return

$$\int x_k p_2(x_k, t_k | x_{k-1}, t_{k-1}; \dots; x_1, t_1) dx_k = x_{k-1}, \quad (38)$$

so that no gain is expected in sequential time intervals, no matter how much you know about the past. I.e., if the same sequence (x_{n-1}, \dots, x_1) was observed at some other time in the past and a return $x_n \gg x_{n-1}$ had then occurred, we have no reason to expect that accident/fluctuation to be repeated. The best forecast of x_n is still $\langle x_n \rangle_{\text{cond}} = x_{n-1}$. Since we can average over x_{k-1}, \dots, x_1 , we can also predict/forecast that

$$\langle \langle x_k \rangle \rangle = \int x_k p_k(x_k, t_k | x_{k-1}, t_{k-1}; \dots; x_1, t_1) p_{k-1}(x_{k-1}, t_{k-1} | x_{k-2}, t_{k-2}; \dots; x_1, t_1) dx_k dx_{k-1} = \langle x_{k-1} \rangle = x_{k-2} \quad (39)$$

etc., and finally

$$\langle \langle \dots \langle x_k \rangle \dots \rangle \rangle = \int x_2 p_2(x_2, t_2 | x_1, t_1) dx_2 = x_1. \quad (40)$$

Summarizing, the progression from statistical independence to Markov processes to Martingales can be understood as a systematic reduction in restrictions. For statistical independence, the n -point density factors, $f_n(x_n, \dots, x_1) = f_n(x_n) \dots f_1(x_1)$. A Markov process generalizes this by allowing f_n to be determined by p_2 and f_1 alone, $f_n(x_n, \dots, x_1) = p_2(x_n; x_{n-1}) \dots p_2(x_2; x_1) f_1(x_1)$. Every drift-free Markov process is a martingale, $\langle x(t_n) \rangle_c = x_{n-1}$. The most general martingale keeps only the last condition and permits

memory, $p_n \neq p_2$ for $n \geq 3$. In this way we have a successive progression of complication in processes. All three classes of processes have in common that the increment autocorrelations vanish. But for statistical independence $\langle x(s)x(t) \rangle = 0$, whereas for martingales $\langle x(s)x(t) \rangle = \langle x^2(s) \rangle$ if $s < t$. Fractional Brownian motion and other systems with long time increment autocorrelations fall completely outside this hierarchy.

Here's a different summary. Wiener processes have statistically independent, stationary increments with variance linear in time. Consequently, a Wiener process has a time translationally invariant transition density. One can generalize the Wiener process in at least three different directions. Time translationally invariant processes include Mori-Zwanzig processes and the approach to statistical equilibrium, but these are not Ito processes (the noise is correlated). In Zwanzig-Mori processes, the increments are generally correlated and nonstationary. Processes with stationary correlated increments and variance nonlinear in the time yield the long time autocorrelations characteristic of fBm. These processes are neither of the Ito nor Zwanzig-Mori type. A Wiener process is a martingale. One can generalize to non-Wiener martingales (drift-free Ito processes), and from there to general Ito processes with drift.

5. The Efficient Market Hypothesis

We begin by summarizing our viewpoint for the reader. Real finance markets are hard to beat, arbitrage possibilities are hard to find and, once found, tend to disappear fast. In our opinion the EMH is simply an attempt to mathematize the idea that the market is very hard to beat. If there is no useful information in market prices, then those prices can be counted as noise, the product of 'noise trading'. A

martingale formulation of the EMH embodies the idea that the market is hard to beat, is overwhelmingly noise, but leaves open the question of hard to find correlations that might be exploited for exceptional profit.

A strict interpretation of the EMH is that there are no correlations, no patterns of any kind, that can be employed *systematically* to beat the average return $\langle R \rangle$ reflecting the market itself: if one wants a higher return, then one must take on more risk. A Markov market is unbeatable, it has no systematically repeated patterns, no memory to exploit. We will argue below that the stipulation should be added that in discussing the EMH we should consider only normal, liquid markets, meaning very liquid markets with small enough transactions that approximately reversible trading is possible on a time scale of seconds [3]. Otherwise, 'Brownian' market models do not apply. Liquidity, 'the money bath' created by the noise traders whose behavior is reflected in the diffusion coefficient [3], is somewhat qualitatively analogous to the idea of the heat bath in thermodynamics [24]: the second by second fluctuations in $x(t)$ are created by the continual noise trading.

Mandelbrot [17] proposed a less strict and very attractive definition of the EMH, one that directly reflects the fact that financial markets are hard to beat but leaves open the question whether the market can be beaten in principle at some high level of insight. He suggested that a martingale condition on returns realistically reflects the notion of the EMH. A martingale may contain memory, but that memory can't be easily exploited to beat the market precisely because the expectation of a martingale process $x(t)$ at any later time is simply the last observed return. In addition, as we've shown above, pair correlations in increments cannot be exploited to beat the market either. The idea that memory may arise (in commodities, e.g.) from other variables (like

the weather) [17] corresponds in statistical physics [11] to the appearance of memory as a consequence of averaging over other, more slowly changing, variables in the larger dynamical system.

The martingale (as opposed to Markov) version of the EMH is also interesting because technical traders assume that certain price sequences give signals either to sell or buy. In principle, that is permitted in a martingale. A particular price sequence $(p(t_n), \dots, p(t_1))$, were it quasi-systematically to repeat, can be encoded as returns (x_n, \dots, x_1) so that a conditional probability density $p_n(x_n; x_{n-1}, \dots, x_1)$ could be interpreted as providing a risk measure to buy or sell. By 'quasi-repetition' of the sequence we mean that $p_n(x_n; x_{n-1}, \dots, x_1)$ is significantly greater than a Markovian prediction. Typically, technical traders make the mistake of trying to interpret random price sequences quasi-deterministically, which differs from our interpretation of 'technical trading' based on conditional probabilities (see Lo et al [25] for a discussion of technical trading claims, but based on a non-martingale, non-empirically based model of prices). With only a conditional probability for 'signaling' a specific price sequence, an agent with a large debt to equity ratio can easily suffer the Gamblers' Ruin. In any case, we can offer no advice about technical trading, because the existence of market memory has not been firmly established (the question is left open by the analysis of ref. [25]), liquid finance markets look pretty Markovian so far as we've been able to understand the data [4], but one can go systematically beyond the level of pair correlations to try to find memory. Apparently, this remains to be done, or at least to be published.

If we return to the hypothetical models (22b,c), the case where $x_0 \neq 0$ implies a finite value $p_c \neq 0$, so that (22c) describes a hypothetical market where there is persistence of memory

of some previous valuation. Such memory could reflect heavy trading around a particular price and can, of course, be lost in the course of time. The writer remembers well the period of a few months ca. 1999 when CPQ sold for around \$22, and was traded often in the range \$18-\$25 before crashing further. Whether that provides an example is purely speculation at this point.

Fama [26] took Mandelbrot's proposal seriously and tried to test finance data at the simplest level for a fair game condition. We continue our discussion by first correcting a mathematical mistake made by Fama (see the first two of three unnumbered equations at the bottom of pg. 391 in [26]), who wrongly concluded in his discussion of martingales as a fair game condition that $\langle x(t+T)x(t) \rangle = 0$. Here's his argument, rewritten partly in our notation. Let $x(t)$ denote a 'fair game'. With the initial condition chosen as $x(t_0) = 0$, then we have the unconditioned expectation $\langle x(t) \rangle = \int x dx f_1(x, t) = 0$ (there is no drift). Then the so-called 'serial covariance' is given by

$$\langle x(t+T)x(t) \rangle = \int x dx \langle x(t+T) \rangle_{\text{cond}(x)} f_1(x, t). \quad (41)$$

Fama states that this vanishes because $\langle x(t+T) \rangle_{\text{cond}} = 0$. This is impossible: by a fair game we mean a Martingale, the conditional expectation is $\langle x(t+T) \rangle_{\text{cond}} = \int y dy p_2(y, t+T; x, t) = x = x(t) \neq 0$, and so Fama should have concluded instead that $\langle x(t+T)x(t) \rangle = \langle x^2(t) \rangle$ as we showed in the last section. Vanishing of (41) would be true of statistically independent variables but is violated by a 'fair game'. Can Fama's argument be salvaged? Suppose that instead of $x(t)$ we would try to use the *increment* $x(t, T) = x(t+T) - x(t)$ as variable. Then $\langle x(t, T)x(t) \rangle = 0$ for a Martingale, as we showed in part 4. However, Fama's argument still would not be generally correct because $x(t, T)$ cannot be taken as a 'fair game' variable unless the variance

is linear in t , and in financial markets the variance is not linear in t [3,4]. Fama's mislabeling of time dependent averages (typical in economics and finance literature) as 'market equilibrium' has been corrected elsewhere [24].

In our discussion of the EMH we shall not follow the economists' tradition and discuss three separate forms (weak, semi-strong, and strong [27]) of the EMH, where a hard to test or effectively nonfalsifiable distinction is made between three separate classes of traders. We specifically consider only normal liquid markets with trading times at multiples of 10 min. intervals so that a Martingale condition holds [4]. Normal market statistics overwhelmingly (with high probability, if not 'with measure one') reflect the noise traders [3], so we consider *only* normal liquid markets and ask whether noise traders produce signals that one might be able to trade on systematically. The question whether insiders, or exceptional traders like Buffett and Soros, can beat the market probably cannot be tested scientifically: even if we had statistics on such exceptional traders, those statistics would likely be too sparse to draw a firm conclusion (see [3,4] for a discussion of the difficulty of getting good enough statistics on the noise traders, who dominate a normal market). Furthermore, it is not clear that they beat liquid markets, some degree of illiquidity seems to play a significant role there. Effectively, or with high probability, there is only one type trader under consideration here, the noise trader. Noise traders provide the liquidity [28], their trading determines the form of the diffusion coefficient $D(x,t;\{x\})$ [3], where $\{x\}$ reflects any memory present. The question that we emphasize is whether, given a Martingale created by the noise traders, a normal liquid market can still be beaten systematically.

One can test for martingales and for violations of the EMH at increasing levels of correlation. At the level $n=1$, the level of

simple averages, the ability to detrend data implies a Martingale. At the level $n=2$, vanishing increment autocorrelations [4] implies a martingale. Both conditions are consistent with Markov processes and with the EMH. A positive test for a martingale *with memory* at the level $n \geq 3$ would eliminate Markov processes, and perhaps would violate the EMH as well. So far as we're aware, this case has not yet been proposed or discussed in the literature. If such correlations exist and would be traded on, then a finance theorist would argue that they would be arbitrage=ged away, changing the market statistics in the process. If true, then this would make the market even more effectively Markovian.

A Markov market cannot be systematically beaten, it has no memory of any kind to exploit. Volatility clustering [17] and so-called 'long term dependence' [29] appear in Markov models [30], are therefore not necessarily memory effects. In the folklore of finance it's believed that some traders are able to make money from volatility clustering, which is a Markovian effect with a nontrivial variable diffusion coefficient $D(x,t)$, e.g. $D(x,t)=t^{2H-1}(1+abs(x)/t^H)$ [30], so one would like to see the formulation of a trading strategy based on volatility clustering to check the basis for that claim.

Testing the market for a nonMarkovian martingale is nontrivial and apparently has not been done: tests at the level of pair correlations leave open the question of higher order correlations that may be exploited in trading. Whether the hypothesis of a martingale as EMH will stand the test of higher orders correlations exhibiting memory remains to be seen. In the long run, one may be required to identify a very liquid 'efficient market' as Markovian.

Finally, martingales typically generate nonstationary increments. This means that it is generally impossible to use

the increment $x(t,T)$ (or the price difference $p(t+T)-p(t)$) as a variable in the description of the underlying dynamics. The use of a returns or price increment as variable in data analysis generates spurious Hurst exponents [4,31] and spurious fat tails whenever the time series have nonstationary increments [3,4]. The reason that an increment cannot serve as a 'good' coordinate is that it depends on the starting time t : let $z=x(t;T)$. Then

$$f(z,t,t+T) = \int f_2(y,t+T;x,t)\delta(z-y+x)dx dy \quad (42)$$

is not independent of t , although attempts to construct this quantity as histograms in data analysis via 'sliding windows' implicitly presume t -independence [4,31]. If the increments are stationary then $z=y-x=x(T)$ and we obtain a well defined density $f(z,T)$. When the increments are nonstationary then f depends on t and (42) can be seen a failed attempt to coarsegrain. Correspondingly, there exists no Langevin eqn. for nonstationary increments. A Langevin eqn. for the increments can be obtained when the variance is linear in t , so that the increments are both stationary and uncorrelated; the increment is then independent of t and serves as a 'good' coordinate. But in the general case of stationary increments with finite variance, unless the variance is linear in t there are long time correlations that destroy the fair game/martingale property. Nearly all existing data analyses are based on a method of building histograms called 'sliding windows' [4]. Sliding a window from one value of t to another to read off $x(T)$ from $x(t,T)=x(t+T)-x(t)=x(T)$ inherently assumes that the increments $x(t,T)$ are stationary (see [31] for the original discussion of the importance of nonstationary increments in FX data analysis).

6. Martingales as EMH for nontrivial drift coefficients

In our analysis [4] of Euro-Dollar 1999-2004 FX data, the average drift is a small constant that can be ignored. We can't rule out that that result may be era dependent. What would happen if an x -dependent drift were important? E.g., in martingale option pricing an x -dependent drift $R(x,t)=r-D(x,t)/2$ is theoretically necessary [24,32], where r is the risk free interest rate (or more generally the cost of carry [23]). In reality option pricing via the exponential distribution has been successful with the neglect of that term [24,32].

However, consider a market like the U.S. stock markets from 1994-2000, where the average drift $\langle R \rangle$ should describe the bubble. If an x -dependent drift is a necessary consideration, then the condition for a Martingale as the EMH must be slightly modified.

With an x -dependent drift $R(x,t)$ the stochastic integral equation for the market consists of a drift term plus a martingale,

$$x(t+T) = x(t) + \int_t^{t+T} R(x(s),s)ds + \int_t^{t+T} \sqrt{D(x(s),s)}dB(s). \quad (43)$$

Whether or not R and/or D contain memory is at this stage unimportant. We can define an average drift

$$\langle R \rangle = \int dx R(x,t) p_2(x,t|x_0,t_0), \quad (44)$$

reflecting e.g. an intraday average [4] conditioned on the daily initial conditions. If we can subtract the drift from x then the resulting process is not a martingale. The best we can obtain in this case is the restricted condition

$$\int y dy p_2(y, t | x_0, t_0) = x_0 \quad (45)$$

where we can, e.g., take as initial condition the initial return at opening time t_0 each day. However, the nice condition of uncorrelated increments is lost,

$$\langle x(t, -T)x(t, T) \rangle = \left\langle \int_{t-T}^t ds (R(x(s), s) - \langle R \rangle) \int_t^{t+T} dw (R(x(w), w) - \langle R \rangle) \right\rangle \neq 0 \quad (46)$$

so we no longer have a clear and easy test on empirical returns data to rule out long time correlations.

To remedy this state of affairs, we're forced to use price as variable. Assume that $R(x, t) = \mu - D(x, t)/2$, which reflects the assumption that the basic market equation of motion is

$$dp = \mu p dt + \sqrt{p^2 d(p, t)} dB(t) \quad (47)$$

with $d(p, t) = D(x, t)$ determined empirically, where μ is the expected 'interest rate' on the financial instrument under consideration. In ref. [25] a nonmartingale Bachelier-type model was assumed, $p^2 d(p, t) = \text{constant}$, and 'patterns' were assumed without proof to be encoded in a nonlinear drift coefficient. Next, using as returns variable $y = x - \mu t$, with $S = pe^{-\mu t}$, we get a price martingale

$$dS = \sqrt{S^2 e(S, t)} dB(t) \quad (48)$$

where (by Ito calculus) $e(S, t) = d(p, t) = D(x, t)$. The condition to be tested empirically to establish this model is therefore $\langle S(t, t-T)S(t, T) \rangle = 0$, where the increments $S(t, T) = S(t+T) - S(t)$ will generally be nonstationary with $\langle S(t+T)S(t) \rangle = \langle S^2(t) \rangle$ if $T > 0$. If there is a drift coefficient with memory, then this model cannot be established. In the case of (47) the memory

must be reflected in the diffusion coefficient. This possibility has not been studied in the finance literature. Of course, to set the idea to work one must first get an accurate estimate for μ , a nontrivial task, empirically seen.

7. Levy's definition of Brownian motion, a cautionary note

Levy's characterization of "Brownian motion" (meaning the Wiener process) is stated in various equivalent ways in the literature (pg. 46 in Friedman [10], pg. 75 Durrett [12], pg. 204 in Steele [14], and pg. 111 in Durrett [34]) We can identify the careless reading of that theorem as the source of the false expectation expressed in much of the finance literature that an arbitrary martingale is equivalent by a change of time variable to a Wiener process (see pg. 204-5 in Steele [14] for that mistake, but see also pg. 75 in [12] for the same claim). Levy's definition can be stated as follows [10]: with the assumptions that $Y(t)$ and $Y^2(t)-t$ are both martingales, then $Y(t)$ is a Wiener process within a change of time variable. Here's the most general construction of a martingale from Ito calculus: let $x(t)$ be any Ito process $dx=R(x,t)dt+\sqrt{D(x,t)}dB(t)$. A local martingale $Y(t)=G(x,t)$ can be constructed by setting the drift term equal to zero in Ito's lemma (requiring that $G(X,t)$ satisfies Kolmogorov's backward time pde subject to initial and boundary conditions) and is generated by the sde

$$dY = \frac{\partial G}{\partial x} \sqrt{D(x,t)} dB. \quad (48)$$

Durrett [12] shows how to construct and do Ito calculus with martingales, a generalization of the standard case where Ito differentials and stochastic integration are developed for Wiener processes [10,12,13,14]. For a martingale Y , the easy to prove integration by parts formula becomes [12]

$$Y(t) - Y(t_0) = \int (dY)^2 + 2 \int Y dY, \quad (49)$$

where $(dY)^2 = E(x,t)dt$ with $E(x,t) = G'^2(x,t)D(x,t)$, showing that $Y^2(t) - \int (dY)^2$ is a martingale. This reduces to the Wiener martingale $Y^2(t) - t$ iff. $G'(x,t)\sqrt{D(x,t)} = 1$. E.g., for the drift-free exponential process [9] with $H=1/2$ and $x(0)=0$, $\langle x^2(t) \rangle = 2t$, showing that $\langle x^2(t) \rangle - 2t$ is a martingale, and therefore $\langle x^2(t) \rangle - t$ is not.

Durrett [12] emphasizes continuity of paths in his discussion of Levy's theorem. Scaling Markov processes [9] processes are generated by a drift-free sde with by $D(x,t) = |t|^{2H-1}D(u)$ where $u = |x|/|t|^H$, and satisfy the required conditions [21] for uniqueness and continuity of paths $x(t)$ if the diffusion is not stronger than quadratic, $D(u) = 1+u^n$ with $n \leq 2$, and if $t > 0$. The restriction to $t > 0$ would seem problematic, but we've shown by direct construction [9] that Green functions $g(x,t;y,s)$, $g(x,t;y,t) = \delta(x-y)$, exist for those processes at least for $y=0$, $s=0$, independent of n , so long as the indefinite integral $\int du/D(u)$ is finite. The exponential process is generated by $D(u) = 1+u$.

To complete the proof, we can show that the integrability requirements for the transformation of an arbitrary martingale $X(t)$, $dX = \sqrt{D(X,t)}dB(t)$, to a Wiener process $B(t)$ are not satisfied. Assume a transformation $Y(t) = G(X,t)$ such that $dY = \mu(t)dt + \sigma(t)dB$, i.e., Y is to be a time change on a Wiener process with drift, where $\sigma(t) \neq 1$ defines a time change on the Wiener process. From Ito's lemma we obtain

$$\frac{\partial G}{\partial X} \sqrt{D(X,t)} = \sigma(t) \quad (50)$$

$$\frac{\partial G}{\partial t} + \frac{D(X,t)}{2} \frac{\partial^2 G}{\partial X^2} = \mu(t)$$

and therefore

$$\frac{\partial G}{\partial t} = \mu(t) + \frac{\sigma(t)}{8} D^{-1/2} \frac{\partial D}{\partial X}. \quad (51)$$

The integrability condition

$$\frac{\partial^2 G}{\partial t \partial X} = \frac{\partial^2 G}{\partial X \partial t} \quad (52)$$

then yields

$$\frac{d\sigma}{dt} = \sigma \left[-\frac{\partial D / \partial t}{2D} - \frac{1}{4} \frac{\partial^2 D}{\partial X^2} + \frac{(\partial D / \partial X)^2}{8D} \right] = 0. \quad (53)$$

With $D(X,t)$ specified in advance, this equation produces a factor $\sigma(t)$ independent of X iff. $D(X,t)$ is independent of X . In that case

$$\sigma(t) = C \sqrt{D(t)} \quad (54)$$

yields merely a time change on standard Brownian motion $B(t)$ (meaning the Wiener process). Steele (pg. 205 in [14]) explicitly and apparently unknowingly restricts himself to this case, and the discussion of Girsanov's theorem in references [13,14] is also restricted to this case by virtue of the assumption that adding a drift term R to a Wiener

process yields another Wiener process (that is possible iff. the drift coefficient R is independent of x !). As Durrett [12] shows while using notation that is misleading for a physicist (" $\langle X \rangle$ " is not an average of X but rather means $\int (dX)^2 = \int E(X,t) dt$, e.g.), the correct statement of the Girsanov theorem is that removing an arbitrary drift term A via the Cameron-Martin-Girsanov transformation from a martingale $X(t)$ plus the drift A , $X(t)+A$, yields another martingale $M(t)$, and we see clearly that, in general, is neither of these martingales a Wiener process. "Intrinsic time" of the sort assumed by Durrett and Steele is discussed *explicitly* for the case where the diffusion coefficient $D(t)$ depends on t alone by McKean (pg. 29 in [34]). The idea of 'intrinsic time', a special time variable where $H=1/2$ so that increments are stationary, is constructed locally by Gallucio et al in an empirical analysis [31], where we know that the diffusion coefficient depends on both x and t [4].

If we ask which time translationally invariant diffusions, $dX = \sqrt{D(X)} dB$, map to a Wiener process, then (53) yields

$$-D \frac{\partial^2 D}{\partial X^2} + \frac{(\partial D / \partial X)^2}{2} = cD \quad (55)$$

with c a constant. This pde has at least one solution, $D(X) = aX^2$ with $a > 0$ a constant and $c = 0$. We obtain the transformation $Y = \ln X$ mapping the lognormal process $X(t)$ to the Wiener process $Y(t) = -(a/2)t + \sqrt{a}B(t)$. So Wiener processes with different time scales map to wiener processes, and the lognormal process maps to a wiener process. Aside from those special cases, the pot is empty.

Sumarizing, we've shown that *arbitrary martingales are topologically inequivalent to Wiener processes*: there is no global transformation $Y = G(X,t)$ of an arbitrary martingale X to a Wiener process. This is analogous to nonintegrability in

deterministic nonlinear dynamics, where chaotic and complex motions are topologically inequivalent to globally integrable ones. Locally, every Ito process reduces to a Wiener process with drift, and this is analogous to local integrability in dynamical systems theory where all deterministic motions satisfying a Lifshitz condition can be mapped locally to translations at constant speed on a lower dimensional manifold [34,35]. Assuming in the literature that arbitrary martingales are equivalent to Wiener processes trivializes martingales, and also leads to mistakes in calculations of first passage times, or 'hitting times': having provided us with the correct general formalism for stochastic calculus based on martingales, Durrett (eqn. (1.5) on pg. 212 of [36]) wrongly assumes with no explanation that Levy's theorem guarantees that an arbitrary martingale is merely a time transformation on a Wiener process.

Acknowledgement

Kevin E. Bassler is supported by the NSF through grants #DMR-0406323 and #DMR-0427938. Gemunu H. Gunaratne is supported by the NSF through grant #DMS-0607345 and by TLCC. Joseph L. McCauley is grateful to a referee of [1] for encouraging us to extend our analysis to include the EMH, to Harry Thomas for sending us four of his papers on the time evolution of nonMarkovian systems and to extremely helpful criticism, comments, and suggested corrections via email, and to Enrico Scalas for pointing us to Doob's assertion that a Chapman-Kolmogorov Equation is not a sufficient condition to make a process Markovian [32], which led us to ref. [7,8]. JMC is also especially grateful to Giulio Bottazzi, who in an email discussion [37] suggested integrability conditions similar to those of part 7 by attempting to construct a transformation from an arbitrary

Ito process with drift $R(x,t)$ to a Wiener process with $\mu(t)=0$ and $\sigma(t)=1$.

Appendix

Beginning with the sde (21) but with drift included,

$$dx = R(x, t; \{x, t\})dt + \sqrt{D(x, t; \{x, t\})}dB, \quad (A1)$$

where $\{x, t\}$ denotes a history of a finite nr. k of states $(x_k, t_k; \dots, x_1, t_1)$, consider the time evolution of any dynamical variable $A(x)$ that does not depend explicitly on t (e.g., $A(x)=x^2$). The sde for A is given by Ito's lemma [21,24]

$$dA = \left(R \frac{\partial A}{\partial x} + \frac{D}{2} \frac{\partial^2 A}{\partial x^2} \right) dt + \frac{\partial A}{\partial x} \sqrt{D(x, t)} dB \quad (A2)$$

With

$$x(t) = x(s) + \int_s^t R(x(q), q; \{x\})dq + \int_s^t \sqrt{D(x(q), q; \{x\})}dB(q) \quad (A3)$$

we form the conditional average

$$\langle A \rangle_t = \int p_n(x, t; y, s; x_k, t_k; \dots; x_1, t_1) dy \quad (A4)$$

where $n=k+2$. Then

$$\langle dA \rangle = \left(\left\langle R \frac{\partial A}{\partial x} \right\rangle + \left\langle \frac{D}{2} \frac{\partial^2 A}{\partial x^2} \right\rangle \right) dt \quad (A5)$$

Using $\langle dA \rangle / dt = d\langle A \rangle_t / dt$ and integrating by parts while ignoring the boundary terms¹, we obtain

$$\int dx A(x) \left[\frac{\partial p_n}{\partial t} + \frac{\partial(Rp_n)}{\partial x} - \frac{1}{2} \frac{\partial^2(Dp_n)}{\partial x^2} \right] = 0, \quad (\text{A6})$$

so that for an arbitrary dynamical variable $A(x)$ we get the Fokker-Planck pde (Kolmogorov's second eqn.)

$$\frac{\partial p_n}{\partial t} = - \frac{\partial(Rp_n)}{\partial x} + \frac{1}{2} \frac{\partial^2(Dp_n)}{\partial x^2}. \quad (\text{A7})$$

for the 2-point transition density depending on a finite history. See Friedman [10] for a rigorous derivation of Kolmogorov's first and second equations and the Chapman-Kolmogorov eqn. from Ito's lemma, and see also McCauley [19] for the explanation why finite memory is not excluded in that derivation.

References

1. J. L. McCauley , G.H. Gunaratne, & K.E. Bassler, *Hurst Exponents, Markov Processes, and Fractional Brownian Motion*, *Physica A* (2007), in press.
2. P. Hänggi, H. Thomas, H. Grabert, and P. Talkner, *J. Stat. Phys.* **18**, 155, 1978.

¹ If the transition density has fat tails, then higher moments will diverge. There, one must be more careful with the boundary terms.

3. J. L. McCauley, K.E. Bassler, & G.H. Gunaratne, *On the Analysis of Time Series with Nonstationary Increments* in *Handbook of Complexity Research*, ed. B. Rosser, 2007.
4. K.E. Bassler, J. L. McCauley, & G.H. Gunaratne, *Nonstationary Increments, Scaling Distributions, and Variable Diffusion Processes in Financial Markets*, 2006.
5. R.L. Stratonovich. *Topics in the Theory of Random Noise*, Gordon & Breach: N.Y., tr. R. A. Silverman, 1963.
6. M.C. Wang & G.E. Uhlenbeck in *Selected Papers on Noise and Stochastic Processes*, ed. N. Wax, Dover: N.Y., 1954.
7. W. Feller, *The Annals of Math. Statistics* **30**, No. 4, 1252, 1959.
8. M. Courbage & D. Hamdan, *The Annals of Probability* **22**, No. 3, 1662, 1994.
9. K.E. Bassler, G.H. Gunaratne, & J. L. McCauley , *Physica A* **369**, 343 (2006).
10. A. Friedman, *Stochastic Differential Equations and Applications*, pg. 23, Academic, N.Y., 1975.
11. P. Hänggi and H. Thomas, *Zeitschr. Für Physik* **B26**, 85, 1977.
12. R. Durrett, *Brownian Motion and Martingales in Analysis*, Wadsworth, Belmont, 1984.
13. M. Baxter and A. Rennie,. *Financial Calculus*, Cambridge, Cambridge, 1995.

14. J.M. Steele, *Stochastic Calculus and Financial Applications*. Springer-Verlag, N.Y., 2000.
15. Joseph L. McCauley, *Stochastic processes for physics and finance*, book manuscript to be submitted (2007).
16. B. Mandelbrot & J. W. van Ness, *SIAM Rev.* **10**, 2, 422, 1968.
17. B. Mandelbrot, *J. Business* **39**, 242, 1966.
18. J. L. McCauley, *Markov vs. nonMarkovian processes: A comment on the paper 'Stochastic feedback, nonlinear families of Markov processes, and nonlinear Fokker-Planck equations'* by T.D. Frank, submitted (2007).
19. J. L. McCauley, *Fokker-Planck and Chapman-Kolmogorov equations for Ito processes with finite memory*, submitted (2007).
20. P. Embrechts and M. Maejima, *Self-similar Processes*, Princeton University Press, Princeton, 2002.
21. L. Arnold, *Stochastic Differential Equations*, Krieger, Malabar, 1992.
22. E. Scalas, R. Gorenflo and F. Mainardi, *Physica A* **284**, 376, 2000.
23. F. Mainardi, M. Raberto, R. Gorenflo and E. Scalas *Physica A* **287**, 468, 2000.
24. J.L. McCauley, *Dynamics of Markets: Econophysics and Finance*, Cambridge, Cambridge, 2004.
25. A. W. Lo, H. Mamaysky, and J. Wang, *J. Finance* **LV**, Nr. 4, 1705, 2000.

26. E. Fama, *J. Finance* **25**, 383-417, 1970.
27. Johannes A. Skjeltorp, *Scaling in the Norwegian stock market*, *Physica A* **283**, 486-528, 2000.
28. F. Black, *J. of Finance* **3**, 529, 1986.
29. C.C. Heyde & N.N. Leonenko, *Adv. Appl. Prob.* **37**, 342, 2005.
30. K.E. Bassler, G.H. Gunaratne, & J. L. McCauley, work in progress (2007).
31. S. Galluccio, G. Caldarelli, M. Marsili, and Y.-C. Zhang, *Physica A* **245**, 423, 1997.
32. J. L. McCauley, G.H. Gunaratne, & K.E. Bassler, *Martingale option pricing*, submitted, 2006.
33. J. L. Snell, *A Conversation with Joe Doob*, <http://www.dartmouth.edu/~chance/Doob/conversation.html>; *Statistical Science* **12**, No. 4, 301, 1997.
34. H. P. McKean, *Stochastic Integrals*, AMS Chelsea, Providence, 2000.
35. V.I. Arnol'd, *Ordinary Differential Equations*, transl. by R. Cooke, Springer, N.Y., 1992.
36. J.L. McCauley, *Classical Mechanics: flows, transformations, integrability and chaos*, Cambridge Univ. Pr.: Cambridge, 1997.
37. R. Durrett, *Stochastic Calculus*, CRC, Boca Raton, 1996.

38. G. Bottazzi, private communication, March, 2006