



Munich Personal RePEc Archive

Trends in preschool lead exposure, mental retardation, and scholastic achievement: association or causation?

Rick Nevin

11. November 2008

Online at <https://mpra.ub.uni-muenchen.de/35339/>

MPRA Paper No. 35339, posted 11. December 2011 02:17 UTC

Trends in Preschool Lead Exposure, Mental Retardation, and Scholastic Achievement: Association or Causation?

Rick Nevin

November 11, 2008

Notice: This is an author's version of a work that was accepted for publication by *Environmental Research*. Changes that resulted from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in [*Environmental Research*, Volume 109, Issue 3, April 2009, Pages 301-310.](#)

*Trends in Preschool Lead Exposure, Mental Retardation, and Scholastic Achievement:
Association or Causation?*

Abstract

This study shows that 1936-1990 preschool blood lead trends explain 65% of the 1948-2001 variation in USA mental retardation (MR) prevalence, 45% of the 1953-2003 variation in the average Scholastic Achievement Test (SAT) verbal score, and 65% of the 1953-2003 variation in the average SAT math score. These temporal relationships are characterized by best-fit time lags (highest R^2 and t-value for blood lead) consistent with lead-induced cognitive damage in the first year of life: A 12-year lag for school-age MR, and a 17-year lag for SAT scores. Recent shifts in age-specific MR prevalence are consistent with recent trends in preschool blood lead. SAT and MR trends by race are consistent with racial differences in how 1960s slum clearance affected childhood exposure to severe lead paint hazards. SAT trends by Hispanic origin are consistent with an especially sharp fall in preschool blood lead in New York City since 1970.

Key Words: Lead Poisoning, IQ, Education

1. Introduction

This study examines whether trends in preschool lead exposure can explain trends in school-age mental retardation (MR) prevalence and Scholastic Achievement Test (SAT) scores. Racial, Hispanic origin, and birth cohort differences in lead exposure are compared with subsequent racial, Hispanic origin, and age-specific shifts in MR prevalence and SAT scores.

The link between MR and lead poisoning was first reported in the 1940s (Byers & Lord, 1943). In the 1960s, lead poisoning was associated only with childhood blood lead above 60 ug/dL (micrograms of lead per deciliter of blood) but subsequent research links lower blood lead to elevated MR risk (Marlowe, 1995; David et al., 1976; 1982). Many studies also show an inverse relationship between blood lead and IQ and academic achievement (U.S. Centers for Disease Control and Prevention, 1991; Lanphear et al. 2005; Miranda et. al, 2007). Higher blood lead is associated with larger absolute losses in IQ, but the impact of each ug/dL increase is greater at lower levels. Preschool blood lead of 10 ug/dl is associated with a loss of 7.4 IQ points relative to children with blood lead of 1 ug/dl (Canfield et al, 2003). Each ug/dl increase from 10-15 ug/dl lowers IQ by 0.323 points and each ug/dl above 15 ug/dl lowers IQ by 0.232 points (Schwartz, 1994). Therefore, blood lead of 40 ug/dL lowers IQ by almost 15 points versus blood lead of 1 ug/dL, and blood lead of 60-80 ug/dL reduces IQ by about 20 to 25 points.

MR is characterized by significant limitations in intellectual functioning and adaptive behavior. The specific IQ associated with “limitations in intellectual functioning” has varied by state, but special education for MR generally requires IQ below 70-75 (National Research Council, 1982). By definition, average IQ is 100, and 25% of the population has IQ below 90, so a loss of 15 to 25

IQ points due to blood lead of 40-80 ug/dL suggests lead exposure could substantially affect the percent of children satisfying the MR condition of significant intellectual limitations. IQ is a good predictor of academic achievement (Neisser et. al, 1996), so the loss of 7.4 IQ points due to blood lead of 10 versus 1 ug/dl also impacts achievement.

Elevated blood lead can be due to lead paint chip ingestion, inhaled air lead, and other pathways, but lead in paint and gasoline had especially pervasive effects due to contaminated dust ingested via hand-to-mouth activity as children crawl (U.S. Environmental Protection Agency, 1986). The lead share of USA pigments fell from near 100% in 1900 to 35% by the mid-1930s (Meyer & Mitchell, 1943), and the USA banned lead paint after 1977, but 80% of pre-1940 and 46% of 1940-59 homes still had some interior lead paint in 1999. (Jacobs et al. 2002) Average preschool blood lead tracked trends in leaded gas use from the 1930s through the 1980s as air lead fallout contaminated dust while lead paint exposure changed slowly with slow changes in the housing stock (Thomas et. al. 1999; Nevin, 2007). Average USA preschool blood lead was 15 ug/dL in the late-1970s, but fell sharply from 1975-87 due to the leaded gas phase-out (Pirkle et al., 1994). Since 1990, preschool blood lead prevalence over 10 ug/dL has tracked trends in lead dust hazards from lead paint (Jacobs & Nevin, 2006; Nevin & Jacobs, 2006; Nevin et al., 2007).

Many city children suffered severe lead poisoning from 1950-70 due to additive exposure to air lead and deteriorated lead paint in city slums. Atmospheric emissions from gas lead affected blood lead even in rural areas, but traffic caused severe city exposure because 55% of emissions settled within 20 km of the roadway (Organization for Economic Cooperation and Development, 1993). Most 1950-70 slum housing was built around 1900 when the use of heavily leaded interior

paint was common. Lead dust hazards are still present in 61% of all homes with deteriorated interior lead paint (Jacobs et al. 2002), and lead paint chip ingestion is often a factor in severe lead poisoning: In 1989-90, after the leaded gas phase-out, children with x-ray evidence of paint chip ingestion had average blood lead of 63 ug/dL. (McElvaine et al. 1992) Slum demolition reduced severe lead paint hazards in the 1960s, but 25% of city children tested still had blood lead over 40 ug/dL when gas lead use peaked around 1970. Gilsinn (1972) found that substandard housing prevalence explained 95% of local variation in children over 40 ug/dL in 1970, but the percent of children over 40 ug/dL relative to substandard housing prevalence was higher in New York City due to higher air lead in larger cities.

Blacks accounted for 15% of central city households in 1960, but occupied 56% of substandard city housing, and the percent of all central city blacks in substandard housing was 25% in 1960 and 16% in 1966 (Kristof, 1968; Koebel, 1996). Per capita gas lead use fell from 1956-1962 but hit new highs from 1966-1974, when 62% of blacks under six-years-old lived in central cities, versus 24% of whites under age six. (U.S. Census, 1960-90) Average blood lead for black two-year-olds in Chicago and New York City fell about 30% from 1970-78, but the late-1970s USA average for black children ages 6-36 months was still 50% above the white average (Agency for Toxic Substances and Disease Registry, 1988).

2. Data and Methods

Early 20th Century trends in MR prevalence are compared with prior trends in paint lead use. Regressions compare post-1935 preschool blood lead trends (Nevin, 2007) with subsequent MR and SAT score trends. Post-1987 trends in preschool blood lead prevalence over 10 ug/dL are

compared with 1991-2006 trends in 6-11-year-old and 12-17-year-old MR prevalence. MR and SAT trends by race (National Research Council, 2002; U.S. Department of Education, 1993) are compared to 1950-1970 urban renewal and gas lead trends (Nevin, 2000; U.S. Department of Housing and Urban Development, 1971). Hispanic SAT trends (U.S. Department of Education, 1997-2007) are compared with post-1970 blood lead trends by Hispanic subpopulation location.


The 1936-99 preschool blood lead estimates in this analysis are anchored by National Health and Nutrition Examination Survey (NHANES) data for 1976-80, 1988-91, 1992-94, and 1998-2000 (Thomas et al, 1999; Pirkle et al, 1994; U.S. Centers for Disease Control and Prevention, 1997 & 2000). Average preschool blood lead before and between survey years were extrapolated from NHANES data and year-to-year percent changes in 1980-88 air lead, 1946-76 refinery lead use, and 1936-46 (leaded) gas use. (U.S. Environmental Protection Agency, 2001; U.S. Census, 1975; Nevin, 2000 & 2007) Although children exposed to lead paint and leaded gas had greater risk of elevated blood lead (because lead ingestion is additive), national trends in *average* blood lead closely tracked air lead and leaded gas use trends (Thomas et al, 1999) due to slow changes in lead paint exposure after the 1930s. Average blood lead estimates used in this analysis are also consistent with available 1956-76 data on national, large urban area, small city, and rural blood lead levels (Nevin, 2007; Organization for Economic Cooperation and Development, 1993).

The calculation of “public school” MR reflects children in special education for MR at the start of each academic year as a percent of total enrollment. Annual data are available for 1976-2005 and straight-line interpolation was used to derive pre-1976 trends from 3-6 year interval data (U.S. Department of Education, 1993 & 1997-2007).

Age-specific MR prevalence reflects children in an age bracket in special education for MR as a percent of total population in that age bracket (U.S. Department of Education, 1995-2007). Age-specific MR trends from 1991-2006 are compared to earlier NHANES data on children ages one to five years, and 1997-2004 surveillance data on children under seven years, with blood lead over 10 ug/dL (US Centers for Disease Control and Prevention, 1997; 2000; 2007a; 2007b).

This analysis reports SAT data by calendar years at the start of academic years (most students take the SAT in the first half of their senior year). SAT data are available for 1966-2004 based on the same SAT norm, and comparable 1951-1965 estimates were derived from data based on an older norm (College Board, 1987-2005; 2002). Figure 1 shows trend estimates for the percent of test-takers who speak a foreign language at home and the percent who took SAT prep courses outside of school. Students who speak a foreign language at home have lower scores (especially verbal scores) and prep courses are expected to increase scores. Prep course trends reflect survey data (Powers, 1998), straight-line interpolation between survey years, and assume no pre-1977 or post-1995 change. Trends for foreign language reflect reported 1986-2004 data and an estimated 1970-86 rise and no pre-1970 change, based on trends in Hispanic enrollment and language spoken at home. A foreign language was spoken at home by 8% of 18-24-year-olds in 1979 and 14% in 1989; Spanish was spoken by 60% of these youths in 1979; and the Hispanic share of 16-24-year-old college enrollment was 3.1% in 1972 (the earliest available data) and 4.5% in 1979 (U.S. Department of Education, 2004; 1997-2007).

Fig. 1: Trend Data and Estimates for SAT Prep and Students Speaking Foreign Language at Home




Legend: An increase in the percent of students who speak a foreign language at home should lower SAT scores (especially verbal scores) and an increase in the percent taking prep courses outside of school should increase scores. Prep course trends reflect survey data, straight-line interpolation between five survey years, and assume no change before 1977 or after 1995. Trends for foreign language spoken at home reflect reported 1986-2004 data, and pre-1986 estimates assume a steady 1970-1986 rise and no pre-1970 change, based language spoken at home and Hispanic enrollment trends. (Powers, 1998; College Board, 1987-2005; U.S. Department of Education, 2004; 1997-2007)

Regressions were run with preschool blood lead lags of 1 to 45 years to identify best-fit lags for MR prevalence and SAT scores, where the “best-fit” lag has the highest significance (t-value) for blood lead and percent of variation explained (R^2). Regression results are reported for best-fit lags based on MR data through 2001 and SAT data through 2003. Graphs show the best-fit lag relationship for blood lead versus MR through 2006 and SAT trends through 2004. To illustrate best-fit lags, regression R^2 is also graphed across time lags for MR prevalence and SAT scores.

3. Results


Figure 2 shows public school MR increased from 0.06% in 1914 to 0.38% in 1935, but was little changed from 1935-39, and fell from 1939-47. Figure 3 shows MR then rose from 0.37% in 1948 to a peak of 2.16% in 1976 and fell to 1.09% in 2006, tracking blood lead with a 12-year lag.

Fig. 2: 1914-1952 Public School MR Prevalence



Legend: Public school MR prevalence increased from 0.06% in 1914 to 0.38% in 1935, as IQ testing increased, but MR prevalence was little changed from 1935-1952, and actually fell from 1939-1947, after a 38% decline in per capita use of lead in paint from 1920-1930, before the rise in gas lead use. (U.S. Department of Education, 1993)


Fig. 3: Preschool Blood Lead vs. Public School MR Prevalence with 12 year Lag



Legend: Public school MR prevalence was 0.37% in 1938, peaked at 2.16% in 1976, and fell to a 42-year low of 1.09% in 2006, tracking preschool blood lead trends with a 12-year lag. (Nevin, 2007; U.S. Department of Education, 1993 & 1997-2007)

Figure 4 graphs R^2 across time lags for which 1936-99 blood lead is statistically significant, showing the 12-year MR time lag is the “best-fit” (highest R^2). Table 1 shows regression results for the best-fit 12-year lag for blood lead versus 1948-2001 MR. (SAT results in Figure 4 and Table 1 are discussed below)

Fig. 4: R² Values across Time Lags for 1953-2003 SAT and 1948-2001 MR Regressions



Legend: R² changes in a striking pattern across regressions comparing 1948-2001 MR prevalence and 1953-2003 SAT scores with different time lags for 1936-1990 preschool blood lead. MR shows no correlation with blood lead with lags of less than 2 or more than 21 years, and R² peaks at a 12-year lag. SAT math R² peaks at an 18-year lag in a simple regression. When the prep and foreign language variables are added to the model, R² peaks at a 17-year lag for math and verbal scores, for test-takers around age 17. These best-fit lags are all consistent with lead-induced cognitive damage in the first year of life, when children ingest lead contaminated dust as they crawl.


Table 1: Regressions for Preschool Blood Lead vs. Mental Retardation Prevalence and Average SAT Math and Verbal Scores

Dependent Variable	Independent Variable	Lag (Years)	Coefficient	Standard Error	t-Value	p-Value	R ²	N
Mental Retardation	Intercept		0.00498	0.00097	5.13	<.0001	0.646	54
	Preschool Blood Lead	12	0.00066	0.00007	9.74	<.0001		
SAT Verbal	Intercept		584.8	3.78	154.61	<.0001	0.900	51
	Preschool Blood Lead	17	-2.1	0.18	-11.81	<.0001		
	SAT Prep		485.4	114.35	4.25	0.0001		
	Foreign Language		-615.0	84.00	-7.32	<.0001		
SAT Verbal	Intercept		574.0	3.27	175.63	<.0001	0.861	51
	Preschool Blood Lead	17	-1.7	0.18	-9.54	<.0001		
	Foreign Language		-268.1	22.59	-11.87	<.0001		
SAT Verbal	Intercept		550.7	5.12	107.49	<.0001	0.454	51
	Preschool Blood Lead	17	-2.2	0.35	-6.38	<.0001		
SAT Math	Intercept		527.9	3.00	175.70	<.0001	0.677	51
	Preschool Blood Lead	17	-1.3	0.14	-9.37	<.0001		
	SAT Prep		189.6	90.84	2.09	0.0423		
	Foreign Language		-135.1	66.73	-2.03	0.0485		
SAT Math	Intercept		523.7	1.82	287.24	<.0001	0.647	51
	Preschool Blood Lead	17	-1.2	0.12	-9.48	<.0001		

Legend: Regressions were run for preschool blood lead versus mental retardation prevalence and average SAT math and verbal scores with time lags of 1 to 45 years, and the best-fit lag (highest R² & t-value for blood lead) was 12 years for mental retardation prevalence and 17 years for average SAT math and verbal scores. The percent of students taking SAT prep courses outside of school and the percent who speak a foreign language at home were also significant in the SAT regressions.


Overall public school MR was relatively stable at 1.2%-1.3% from 1991-2004, but Figure 5 shows age 6-11-year-old MR fell from 1% in 1996 to 0.7% in 2006, and age 12-17-year-old MR prevalence fell from 1.27% in 2001 to 1.13% in 2006. The percent of 18-21-year-olds in special education for MR was 0.4% in 1996 and 2002-2006. Figure 6 shows 6-8-year-old MR declined about 35% from 1993-2006, as 17-year-old MR rose 7%. Age 6-21-year-old black student MR fell from 2.11% in 1998 to 1.69% in 2006, as age 6-21 white MR fell from 0.78% to 0.63%.

Fig. 5: 1991-2006 Age-Specific MR Prevalence Trends



Legend: Age 6-11-year-old MR prevalence fell from 1% in 1996 to 0.7% in 2006, and 12-17-year-old prevalence fell from 1.27% in 2001 to 1.13% in 2006, as the percent of the age 18-21 population enrolled in special education for MR remained at 0.4%. (U.S. Department of Education, 1995-2007) The 6-11-year-old and 12-17-year-old trends are both consistent with a national decline in preschool blood lead prevalence over 10 ug/dL since the late-1980s.

Fig. 6: 1993-2006 Percent Change in MR Prevalence by Student Age




Legend: Ages 6, 7, and 8 recorded 1993-2005 MR prevalence declines of about 35%, as prevalence declined 28% at age 9, 22%-23% at ages 10 and 11, but declined only 4% at ages 16 and 18 and rose 7% at age 17 as more MR students now stay in school until they earn a high school diploma or certificate. (U.S. Department of Education, 1995-2007)

NHANES data show the percent of preschool children with blood lead over 10 ug/dL fell from 88% in 1976-80, to 8.6% in 1988-91, 4.4% in 1991-94, and 1.6% in 1999-2002. Surveillance tests of children with blood lead of 10-15 ug/dL also fell over 40% from 1997-2002, as children with tests of 15-20 ug/dL fell over 50%, and children over 20 ug/dL fell over 55%. Surveillance data show an especially large decline since 1997 in black children with blood lead over 10 ug/dL and over 20 ug/dL (Meyer et al., 2003).

Figure 4 graphs SAT regression R^2 across all time lags for which the independent variables are statistically significant and have the expected sign. R^2 peaks at an 18 year lag across SAT math simple regressions, but the prep course and language variables are significant only with a 16 or 17 year blood lead lag, and multiple regression R^2 peaks at a 17 year lag. R^2 peaks at 23 years for SAT verbal simple regressions, but the prep and language variables are significant only with blood lead lags of 12-19 years, and multiple regression R^2 peaks at a 17 year lag. Figure 7 shows

the fall and rise of SAT math and verbal scores have tracked the rise and fall of blood lead with a 17-year lag. Table 1 shows blood lead with a 17-year lag is highly significant in regressions with and without SAT prep and foreign language variables. R^2 is 64.7% in a simple regression for SAT math versus blood lead with a 17 year lag, and adding the prep and language variables only increases R^2 to 67.7%. R^2 is 45.4% in a simple regression for SAT verbal versus blood lead with a 17 year lag, and adding the prep and language variables increases R^2 to 90%. R^2 is 86.1% in a verbal regression with just the language variable and a 17-year lag for blood lead.


Fig. 7: Preschool Blood Lead vs. SAT Math and Verbal Scores with 17 year Lag



Legend: The fall and rise of average SAT math and verbal scores have tracked the rise and fall of blood lead with a 17-year lag. The decline in scores was blunted by a surge from 1977-1986 in the percent of students taking prep courses outside of school, and the later rise has been slowed by a surge in the percent of students who speak a foreign language at home (especially affecting verbal scores), but the effect of preschool blood lead is still clearly evident. (Nevin, 2007; College Board, 1987-2005; 2002)

Figure 8 shows per capita gas lead use was little changed as urban renewal projects in execution increased 100-fold from the early-1950s through the 1960s. A single urban renewal project often involved demolition of large numbers of dilapidated slum units. Overall public school MR fell from 2.2% in 1976 to 1.2% in 1992, as black student MR fell from 4.1% to 2.5%. White MR fell from 1.3% in 1976 to 1.07% in 1986 then rose to 1.15% in 1992. Average black SAT scores increased substantially from 1975-87, as white SAT scores fell slightly.

Fig. 8: 1950-1970 Trends in Urban Renewal Projects and Per Capita Gas Lead



Legend: Per capita gas lead use was little changed as urban renewal projects increased 100-fold from the early-1950s through the 1960s, where a single project often involved demolition of a large number of dilapidated slum units with severe lead paint hazards. Slum demolition was especially extensive in black neighborhoods, reducing the racial difference in average preschool lead exposure. Subsequent racial convergences in MR prevalence, SAT and National Assessment of Educational Progress scores, juvenile burglary arrest rates, and adult incarceration rates can be traced to this 1960s racial convergence in preschool lead exposure. (Nevin, 2000; U.S. Department of Housing and Urban Development, 1971)

USA preschool blood lead fell after the mid-1970s, but Mexico did not reduce gas lead use until the late-1980s, affecting foreign-born Mexican-Americans. Native-born Mexican-Americans are also disproportionately located in border areas that had cross-border gas lead exposure, whereas the Puerto Rican population is disproportionately located in New York City (U.S Census, 2001). New York City had extremely high air lead in the 1960s, followed by an especially sharp drop in blood lead after 1970. In 1986, average SAT math and verbal scores for college-bound Puerto Ricans were 23 and 21 points lower, respectively, than average scores for Mexican Americans. In 2002, the average math score for Puerto Ricans was 4 points below the Mexican American average, and their average verbal score was 8 points higher than the Mexican American average.

4. Discussion

The increase in MR from 1914-1935 is consistent with expanding use of IQ tests in schools (Plucker, 2003), but MR prevalence was little changed from 1935-39 and declined from 1939-48. That 1940s MR decline occurred among students born after per capita use of lead in USA paint fell 58% from 1914-30 (Nevin, 2000), and before any substantial rise in gas lead exposure.

Best fit lags for MR and SAT trends versus 1936-90 blood lead are consistent with lead-induced cognitive damage in the first year of life. Blood lead coefficients are highly significant at best-fit lags, with the expected positive sign for MR and negative signs for SAT scores. SAT prep and foreign language are also statistically significant with the expected signs, but their inclusion has little impact on the value or significance of the blood lead coefficient. The explanatory power of language and prep is small in the math regression, raising R^2 from 64.7% in a simple regression to 67.7% in a multiple regression. Adding SAT prep and foreign language variables does raise verbal regression R^2 from 45% to 90%, mostly due to the language variable. (SAT scores fell in 2005, but this appears to be related to a new SAT test that year, as the same academic year saw the rival ACT record its biggest increase in average scores in 20 years.)

Racial trends in MR and SAT scores are consistent with birth cohort slum clearance trends that disproportionately affected black children. Average preschool blood lead peaked around 1970, but severe lead poisoning prevalence must have peaked before slum demolition reduced severe lead paint hazards in dilapidated housing. The 1976 MR peak is consistent with a 12-year lag after a peak in severe lead poisoning around 1964, and the fall in MR from 1976-92 was almost entirely due to a decline among black children. Hauser (1998) shows that the racial difference in

National Assessment of Educational Progress (NAEP) scores also narrowed at different times for ages 9, 13, and 17-years-old, but black gains at each age were traced to changes across 1962-73 birth years. The 17-year-old NAEP racial trends are similar to the 1975-87 racial convergence in SAT scores. These MR, SAT, and NAEP trends are all consistent with a 1960s convergence in the relative severity of black and white preschool lead exposure. In absolute terms, however, black children still have higher MR prevalence and lower average SAT and NAEP scores, consistent with black children continuing to have higher average blood lead.

Office of Civil Rights (OCR) data for 1978-79 and earlier MR research found that 75%-80% of all MR students were “Educable Mentally Retarded” (EMR), with adaptive limitations (largely based on teacher observations) and IQ above 55. (National Research Council, 1982) Most EMR cases were of unknown cause, and prevalence was high among low income children who lived in city slums. In 1970, 25% of city children tested had blood lead over 40 ug/dL, and substandard housing explained 95% of local variation in children over 40 ug/dL, so the percent over 40 ug/dL must have been much higher before 1960s slum clearance. Blood lead of 40 ug/dL is associated with a loss of 15 IQ points, so the large percentage of 1960s children over 40 ug/dL must have pushed many of those children below the IQ level associated with EMR. The 1978-79 OCR data and earlier research found higher EMR prevalence among 10-14-year-olds than 5-9-year-olds, suggesting EMR diagnosis was directly related to academic demands of school. Age 15-19-year-old prevalence was half the 10-14-year-old prevalence, as EMR students were more likely to drop out (and lose the MR label once they left school). Recent MR data show a dramatic shift in age-specific MR risk, with peak prevalence at 14 years in 1993 and 16 years in 2005, consistent with younger students showing earlier effects of the ongoing decline in preschool blood lead.

There are several limitations to the data used in this analysis. Public school MR prevalence is somewhat affected by temporal changes in the percent of the school-age population enrolled in public schools, but enrollment trends do not appear to explain MR trends: Grade 9-12 enrollment was 70% of the 14- to 17-year old population in 1952, 80% to 85% from 1963-2001, and 87% in 2005. (U.S. Department of Education, 1993 & 1997-2007) The calculation of age-specific MR also understates total population prevalence for older ages because the numerator excludes MR individuals not enrolled in public school special education, but this mainly affects 18-21-year-old prevalence. The blood lead trends used in this analysis are subject to blood lead survey random error and some estimation error, but potential error is small relative to the large temporal changes in preschool blood lead. This analysis cannot control for temporal changes in public school MR related to changes in MR diagnosis, or education attainment trends that could affect the average parental IQ for students taking the SAT, but trends in education attainment and special education diagnoses (discussed below) do not appear to explain the findings reported here.

Education Attainment Trends versus SAT, MR, and other Special Education Trends

The 1970s SAT decline was initially thought to reflect a change in SAT test-taker population, as increased access to college reduced the average ability of students taking the SAT. Long-term trends now show the opposite relationship between achievement, as measured by the SAT, and educational attainment. SAT scores were relatively stable from 1960-68, as the 16-24-year-old status dropout rate (percent not in school and without a diploma or GED) fell from 27% to 16%, and the 16-24-year-old college enrollment rate of high school completers (enrolled within 12 months of diploma or GED) rose from 45% to 55%. SAT scores fell as the status dropout rate

declined to 14% in 1973 but stayed at 13.9% or higher until 1983, and as the college enrollment rate stayed below 51% from 1972-80 and did exceed the 1968 rate until 1985. The subsequent rise in SAT scores occurred as the dropout rate fell from 13.7% in 1983 to 9.4% in 2005 and the college enrollment rate rose from 55% in 1984 to 68.6% in 2005.

MR students who left school as dropouts fell 23% from 1993-2005 as those graduating with a certificate or diploma rose 97% and 32%, respectively. Some research has suggested falling MR and rising “specific learning disability” prevalence from 1976-90 reflects more restrictive MR criteria shifting mild MR pupils to the specific learning category (U.S. Department of Education, 1991), but this hypothesis implies an increase in average MR severity, which is inconsistent with subsequent gains in MR student attainment.

Shattuck (2006) argues diagnostic substitution could explain the 1994-2003 fall in 6-11-year-old MR and concurrent rise in 6-11-year-old autism, but this hypothesis cannot explain why MR fell from 1976-92. Moreover, diagnostic substitution is not consistent with high autism prevalence among white and upper-income pupils (U.S. Centers for Disease Control and Prevention, 2006) because MR prevalence has always been higher for black and low-income pupils. Also, 17-year-old autism and MR both rose from 1993-2005, as more MR students remained in school.

A multi-site national study has shown that borderline and mild MR students also recorded IQ averaging 5.6 points lower on the 1991 WISC-III versus their earlier scores on the 1974 WISC-R (Kanaya et. al., 2003). This could explain why both 6-11-year-old and 12-17-year-old MR increased in the early-1990s: As more schools adopted the WISC-III in MR assessments, more

students recorded IQ scores associated with MR, when many of those students would have had WISC-R IQ above the level associated with MR. MR student performance on the 1991 WISC-III versus the 1974 WISC-R suggest the 1993-2005 gains in MR student attainment are the result of real gains in cognitive ability in the population classified as MR.

Correlation and Causation Conclusions

This analysis poses a specific variation of the question addressed by Sir Austin Bradford Hill (1965) in "The Environment and Disease: Association or Causation?"

Our observations reveal an association between two variables, perfectly clear-cut and beyond what we would care to attribute to the play of chance. What aspects of that association should we especially consider before deciding that the most likely interpretation of it is causation?

Preschool lead exposure, MR, and SAT trends are reviewed below in the context of the nine “viewpoints” Bradford Hill recommended in assessing evidence of causation.

(1) Strength: Bradford Hill illustrates this viewpoint with evidence that cigarette smokers have a lung cancer death rate nine times that of nonsmokers, and a coronary thrombosis death rate “no more than twice” that of nonsmokers. He notes there is evidence of causation in both cases, but the strength of the lung cancer relationship provides stronger evidence. The association between blood lead and MR trends falls between these two benchmarks: MR risk for children born near the early-1960s peak exposure to air lead and lead paint hazards in slums was five to six times the risk for children born in the late-1930s. The strength of the blood lead, MR, and SAT trend association is also reflected in high regression R^2 and the significance of blood lead coefficients.

(2) Consistency: Has the association “been repeatedly observed by different persons, in different places, circumstances and times?” (Hill, 1965) MR and SAT trends show a strong association with 1936-90 USA blood lead trends, encompassing changes in a large, diverse, national population across four time zones and over five decades. Declining MR from 1939-47, tracking earlier declines in lead paint, shows the same association back to 1920s birth cohorts. These trends are consistent with controlled studies showing a causal association between lead poisoning and MR in individual children. (Byers & Lord, 1943; Marlowe, 1995; David et al., 1976; 1982) The negative correlation between blood lead and SAT trends is consistent with research demonstrating an inverse relationship between preschool blood lead and IQ, and research showing IQ is an excellent predictor of academic achievement. (Schwartz, 1994; U.S. Centers for Disease Control and Prevention, 1991; Lanphear et al. 2005; Miranda et. al, 2007; Canfield et al, 2003; Neisser et. al, 1996) The long-term nationwide association between blood lead, MR, and SAT trends is consistent with slum clearance trends and subsequent racial trends in MR and in SAT and NAEP scores, and with large Puerto Rican SAT gains after a large fall in New York City preschool blood lead.

(3) Specificity. MR and SAT trends show a relationship between preschool blood lead and two manifestations of the specific outcome of impaired cognitive ability, reflecting cognitive impacts across the preschool blood lead distribution. The 1936-90 average blood lead trend reflects a temporal shift in the entire preschool blood lead distribution, including a rise and fall in severe lead poisoning prevalence especially associated with MR, and in marginally elevated blood lead prevalence associated with marginally lower IQ and academic achievement.

(4) Temporality. At a minimum, temporality asks if the suspected cause preceded the effect, but best-fit time lags for MR and SAT scores present an especially compelling case of temporality. While time series comparisons can result in coincidental correlations, R^2 changes in a striking pattern across regressions comparing MR and SAT scores with different time lags for 1936-90 blood lead trends. MR shows no correlation with blood lead with lags of less than 2 or over 21 years – the blood lead coefficient in such regressions is insignificant - and R^2 peaks at a 12-year lag, for students around 12 years old. SAT math scores show no significant relationship with blood lead with lags of less than 7 or over 26 years, and simple regression R^2 peaks at an 18-year lag. With prep and foreign language variables added to the model, R^2 peaks at a 17-year lag for both math and verbal scores, for test-takers around 17 years old. These MR and SAT lags are consistent with lead-induced cognitive damage in the first year of life. Age-specific MR trends reinforce the temporality evidence of causation. Research in the 1980s found MR prevalence peaked in elementary and junior high school, but prevalence peaked at age 14 in 1993 and age 16 in 2006. Declines in 6-11-year-old MR from 1996-2006 and age 12-17 MR from 2002-2006 track the same documented birth cohort decline in preschool blood lead after the late-1980s.

(5) Biological gradient (dose-response relationship): Controlled studies show a dose-response relationship between preschool blood lead and IQ later in life: A loss of 7.4 IQ points due to blood lead of 10 versus 1 ug/dl, and 15 to 25 IQ points associated with blood lead of 40 to 80 ug/dL. (Schwartz, 1994; Canfield et al, 2003) Preschool blood lead trends show a corresponding population-dose-response relationship with population MR and SAT scores. Blood lead rose over four-fold from the 1930s through the 1960s and MR prevalence increased by a similar order of magnitude with a 12-year lag. The fall in MR from the 1970s to 1992 was also roughly

proportionate to the earlier decline in preschool blood lead, before the introduction of a new IQ test used in school evaluations in the early-1990s.

(6) Biological Plausibility. Documented neurochemical, subcellular, and cellular effects of preschool blood lead establish a biological basis for impairments in IQ, learning, and behavior.

These effects, which are not entirely independent, due to the integrated nature of the nervous system, include: (1) inhibition of mitochondria, disrupting energy metabolism; (2) alteration of calcium mediated neurotransmitter release of postsynaptic receptors; (3) reduction in myelin formation; (4) effects on synaptogenesis, presynaptic terminals and postsynaptic receptors; (5) changes in neurotransmitter release, including but not limited to dopamine; (6) abnormalities in electrophysiological activity for both cortex and brain stem including increased latencies and decreased amplitudes of evoked potentials; and (7) effects on brain development related to immaturity of the blood-brain barrier during gestation. (Banks et al., 1997)

(7) Coherence. A causal interpretation “should not seriously conflict with the generally known facts of the natural history and biology of the disease” (Hill, 1965). The association between blood lead, MR, and SAT trends reveals an ecological coherence with controlled studies linking individual lead exposure to cognitive damage. The racial convergence in MR prevalence and SAT and NAEP scores across slum clearance birth years is also consistent with the known history of heavily leaded paint use in older housing, and lead dust and paint chip ingestion risks in severely deteriorated city slums. Lead exposure trends also reveal a coherent relationship with international trends in IQ and social behavior that is correlated with IQ (Nevin 2000; 2007).

The cognitive and behavioral effects of preschool lead exposure can also reconcile the seeming incoherence of controversial research on IQ correlates and recent behavior trends. An analysis of 1979-90 National Longitudinal Survey of Youth (NLSY) data found that dropping out of high school and incarceration were both strongly associated with low IQ, independent of race or socioeconomic status (Herrnstein & Murray, 1994). NLSY participants, ages 14-22 years in 1979, constituted a large representative sample of Americans born in the late-1950s and early-1960s. The high school dropout rate (percent who failed to get a diploma or GED by ages 25-33 years) was 55% for white NLSY participants with IQ below 75, 35% for those with IQ of 75-90, 6% for IQ of 90-110, and less than 1% for IQ over 110. The fall in dropout rates since 1990 means these IQ-related dropout risks cannot be true today. The low NLSY dropout rate for those with IQ above 90 also means the fall in dropout rates must reflect an especially sharp decline for those with IQ below 90. This conclusion is also consistent with the data showing MR student dropouts fell 23% from 1993-2005, as MR students graduating with a diploma or certificate increased.


Less than 1% of white NLSY males with IQ above 110 were incarcerated prior to 1980, versus 3% of those with IQ of 90-110, and 7% of those with IQ below 90. The fall in USA crime since 1990 means these incarceration risks cannot be true today. The low incarceration rate for NLSY males with IQ above 90 also means the crime decline must reflect an especially sharp decline for those with IQ below 90. The 1990s crime decline was led by a sharp fall in juvenile offending, consistent with the decline in MR led by younger students. The 1960s rise in crime was also led by surging juvenile offending, tracking the earlier rise of preschool blood lead (Nevin, 2007), consistent with the 1948-76 rise in MR. The 1960s birth years linked to a racial convergence in

MR prevalence and SAT and NAEP scores are also linked to a racial convergence in juvenile burglary arrest rates. Controlled studies also link preschool lead exposure to delinquent behavior and criminal violence (Denno, 1990; Dietrich et al., 2001 Needleman et al., 1996; 2003). Blood lead trends explain most of the substantial variation in crime rates in the USA, Canada, Britain, Australia, New Zealand, West Germany, France, Italy, and Finland across several decades (Nevin, 2007). Crime rates track blood lead with similar time lags within each nation: A 23-year lag for violent crime, consistent with the typical age of violent offenders, and an 18-year best-fit lag for burglary, consistent with the typical age of property crime offenders.

The NLSY IQ-incarceration correlation has been interpreted as evidence that inherited low IQ is a cause of criminal offending (Gottfredson, 1998). The coherence viewpoint casts doubt on this causal interpretation, because inherited IQ cannot explain crime trends, and IQ is largely stable after childhood whereas criminal offending peaks around ages 15-25 years. Adult white matter growth has been linked to a reduced risk of impulsive behavior as people age (Bartzokis et al., 2001) and juvenile delinquency linked to the impact of preschool lead exposure on impulsivity (Needleman et al., 2003). Preschool lead exposure causes gray matter damage linked to IQ loss, and white matter damage related to behavior, suggesting the NLSY IQ-incarceration correlation could reflect separate lead effects. IQ loss may be permanent, but white matter growth to age 50 could ameliorate lead-induced damage, which could explain lower offending rates for older adults, and the impact of preschool blood lead on national crime rates with best-fit time lags that reflect the typical age of offenders (Nevin, 2007).

Coherent trends in blood lead, education, and crime are now evident in 2000-2006 incarceration rates by age and race (Sabol et al. 2007; Beck & Harrison., 2001). Figure 9 shows incarceration rates fell from 2000-06 for all men under age 30, reflecting the decline in preschool blood lead since the mid-1970s, but incarceration rates are rising for men over 40, born when lead poisoning was epidemic. The overall 30-39-year-old male incarceration rate rose slightly from 2000-06, but the 30-39-year-old rate for black males fell 12%, reflecting the same slum clearance birth years associated with a racial convergence in MR, SAT and NAEP scores, and juvenile burglary arrest rates.

Fig. 9: 2000-2006 Percent Change in Age-Specific Male Incarceration Rates



Legend: Declining incarceration rates from 2000-2006 for males under 30 reflect declining preschool blood lead since the mid-1970s. Incarceration rates are still rising for those over age 40, born when childhood lead poisoning was epidemic. The overall age 30-39 male incarceration rate rose slightly from 2000-2006, but the age 30-39 rate for black males fell 12%, reflecting slum clearance birth years also associated with a racial convergence in MR prevalence and juvenile burglary arrest rates. (Sabol et al. 2007; Beck & Harrison., 2001)

(8) Experimental evidence: If, “because of an observed association some preventive action is taken ... [and it does] in fact prevent ... Here the strongest support for the causation hypothesis may be revealed.” (Hill, 1965) Regulations to eliminate the use of lead in gasoline and to reduce lead paint hazards in older housing anticipated societal benefits in excess of regulatory costs because of observed associations between preschool lead poisoning and MR, between even small

elevations in blood lead and IQ, and between IQ and educational attainment and lifetime earnings. Ecological trends examined here suggest the magnitude of preschool lead exposure's impact on MR and education achievement and attainment trends may have been underestimated in the past, but the direction of that impact was recognized as the basis for deliberate regulatory interventions that have achieved predicted results.

(9) Analogy. “With the effects of thalidomide and rubella before us we would surely be ready to accept slighter but similar evidence with another drug or another viral disease in pregnancy.” (Hill, 1965) Similarly, the effects of blood lead trends before us should lend urgency to eliminating the remaining risk of preschool lead exposure and blood lead levels once considered safe. Some nations still use leaded gas, and children and pregnant women in many nations are still exposed to industrial lead emissions, occupational lead hazards, lead-glazed ceramics, and lead-contaminated home remedies and cosmetics. Lead paint hazards are the greatest remaining USA risk, and a simple window replacement strategy would yield lead hazard reduction benefits plus energy savings from efficient windows that far exceed window replacement costs. (Jacobs & Nevin, 2006; Nevin & Jacobs, 2006; Nevin et al, 2007)

Bradford Hill concluded by addressing *The Case for Action*, with words that clearly apply to the case for moving aggressively to eliminate all preschool lead exposure:

All scientific work is incomplete – whether it be observational or experimental. All scientific work is liable to be upset or modified by advancing knowledge. That does not confer upon us a freedom to ignore the knowledge we already have, or to postpone the action that it appears to demand at a given time. (Bradford Hill, 1965)

References

- Agency for Toxic Substances and Disease Registry (1988) *The Nature and Extent of Lead Poisoning in Children in the United States*, U.S. Department of Health and Human Services
- Banks, E., Ferretti, L., & Shucar, D. (1997). Effects of low level lead exposure on cognitive function in children: A review of behavioral, neuropsychological and biological evidence, *Neurotoxicology*, 18, 237-81
- Bartzokis G, Beckson M, Lu PH, Nuechterlein KH, Edwards N, Mintz J. (2001) "Age-related changes in frontal and temporal lobe volumes in men: a magnetic resonance imaging study", *Archives of General Psychiatry*, 58, 461-465.
- Beck, A. & Harrison, P. (2001) *Prisoners in 2000*, Bureau of Justice Statistics, NCJ 188207, Retrieved March 9, 2006, from: <http://www.ojp.usdoj.gov/bjs/pub/pdf/p00.pdf>
- Byers R. & Lord E. (1943) Late effects of lead poisoning on mental development. *Am J Dis Child* 66:471–494.
- Canfield, R., Henderson, C., Cory-Slechta, D., Cox, C., Jusko, T., & Lanphear, B. (2003). Intellectual impairment in children with blood lead levels below 10 µg/dL. *The New England Journal of Medicine*, 348, 1517-1522.
- College Board (1987-2005) *College-Bound Seniors*
- College Board (2002) Summary Data for Students Administered the SAT Test 1951-52 through 1970-71, Fax from Ellen Sawtell, Data Coordinator and Analyst.
- David, O., Hoffman S., McGann, B., Sverd, J., & Clark, J. (1976). Low lead levels and mental retardation, *Lancet* 2:1376-9
- David, O., Grad, G., McGann, B., & Koltun, A. (1982). Mental retardation and "nontoxic" lead levels, *Am J Psychiatry* 139:806-809

- Denno, D. W. (1990). *Biology and Violence*, Cambridge University Press, NY, NY
- Dietrich, K., Ris, M., Succop, P., Berger, O., & Bornschein R.. (2001). Early Exposure to Lead and Juvenile Delinquency, *Neurotoxicology and Teratology*, 23, 511-518
- Gilsinn, J., (1972) *Estimates of the Nature and Extent of Lead Paint Poisoning in the United States*, U.S. Department of Commerce, National Bureau of Standards.
- Gottfredson L. (1998) The general intelligence factor, *Scientific American*, 9, 24-29
- Hauser, R. (1998) Trends in Black-White Test Score Differentials: I. Uses and Misuses of NAEP/SAT Data, In Neisser, U., *The rising curve: Long-term gains in IQ and related measures*. Washington, DC: American Psychological Association
- Herrnstein, R., & Murray, C. (1994) *The Bell Curve: Intelligence and Class Structure in American Life*. The Free Press, New York.
- Hill, AB. (1965) The environment and disease: association or causation? *Proc R Soc Med*. 58:295–300.
- Jacobs, D., Clickner, R., Zhou, J., Viet, S., Marker, D., Rogers, J., Zeldin, D., Broene, P., & Friedman, W. (2002). The prevalence of lead-based paint hazards in U.S. housing, *Environmental Health Perspectives*, 110, 599-606.
- Jacobs, D. & Nevin, R., (2006). “Validation of a 20-year forecast of US childhood lead poisoning: Updated prospects for 2010”, *Environmental Research* 103: 352-364.
- Kanaya T., Scullin, M., & Ceci, S. (2003) The Flynn Effect and U.S. Policies: The Impact of Rising IQ Scores on American Society Via Mental Retardation Diagnoses, *American Psychologist*, 58: 778-90.
- Koebel, C., (1996). *Urban Redevelopment, Displacement and the Future of the American City*. Community Affairs Office, Federal Reserve Bank of Richmond.

- Kristof, F.S., (1968). *Urban Housing Needs through the 1980's: an Analysis and Projection*. The National Commission on Urban Problems, Research Report No. 10.
- Lanphear B.P., Hornung, R., Khoury, J., Yolton, K., Baghurst, P., Bellinger, D., Canfield, R.L., Dietrich, K.N., Bornschein, R., Greene, T., Rothenberg, S.J., Needleman, H.L, Lourdes, S., Wasserman, G., Graziano, J., Roberts, R. (2005). Low-level environmental lead exposure and children's intellectual function: An international pooled analysis. *Environ Health Perspect* 113:894-899
- Marlowe, M. (1995) *The Violation of Childhood: Toxic Metals and Developmental Disabilities, The Journal of Orthomolecular Medicine*, 10, 79-86.
- McElvaine, M. DeUngria, E., Matte, T., Copley, C., and Binder, S. (1992) "Prevalence of radiographic evidence of paint chip ingestion among children with moderate to severe lead poisoning, St. Louis, Missouri, 1989 through 1990," *Pediatrics*, 89, 740-742.
- Meyer, H. & Mitchell, A. (1943). Lead and zinc pigments and zinc salts, pp. 165-178, In *Minerals yearbook 1941*, US Geological Survey, Washington.
- Meyer, P., Pivetz, T., Dignam, T., Homa, D., Schoonover, J., & Brody, D. (2003) Surveillance for elevated blood lead levels among children 1997-2001, *Morbidity and Mortality Weekly Report* 52 (SS-10) 1-22.
- Miranda, M., Kim, D., Galeano, A., Paul, C., Hull, A., & Morgan, S. (2007) The Relationship between Early Childhood Blood Lead Levels and Performance on End-of-Grade Tests, *Environmental Health Perspectives*, 115, 1242-1247.
- National Research Council (1982). *Placing Children in Special Education: A Strategy for Equity*, National Academy Press, Washington DC.

- National Research Council (2002). *Minority Students in Special and Gifted Education*, National Academy Press, Washington DC.
- Needleman, H., Riess, J., Tobin, M., Biesecker GE, Greenhouse JB. I. (1996). Bone lead levels and delinquent behavior, *Journal of the American Medical Association*, 257, 363-369.
- Needleman, H., McFarland, C., Ness, R., Fienberg, S., & Tobin, M. (2003). Bone lead levels in adjudicated delinquents. A case control study. *Neurotoxicology and Teratology*, 24, 711-717.
- Neisser U., Boodoo, G., Bouchard, T., Boykin, A., Brody, N., Ceci, S., Halpern, D., Loehlin, J Perloff, R., Sternberg, J., Urbina, S. (1996) Intelligence: knowns and unknowns, *American Psychologist*, 51, 77-101.
- Nevin, R. (2000). How lead exposure relates to temporal changes in IQ, violent crime, and unwed pregnancy. *Environmental Research*, 83, 1-22.
- Nevin, R., 2007. Understanding international crime trends: the legacy of preschool lead exposure. *Environmental Research*, 104, 315-336.
- Nevin, R. & Jacobs, D. (2006) Windows of opportunity: lead poisoning prevention, housing affordability, and energy conservation, *Housing Policy Debate*, 17: 185-207.
- Nevin, R., Jacobs, D., Berg, M., & Cohen, J. (2007) Monetary Benefits of Preventing Childhood Lead Poisoning with Lead-Safe Window Replacement, *Environmental Research*, 106: 410-419.
- Organization for Economic Co-Operation and Development (1993) *Risk Reduction Monograph No. 1: Lead*.
- Pirkle, J., Brody, D., Gunter, E., Kramer RA, Paschal DC, Flegal KM, Matte TD.. (1994). The decline in blood lead levels in the United States. *Journal of the American Medical Association*, 272, 284-91

- Plucker, J. A. (Ed.). (2003). Human intelligence: Historical influences, current controversies, teaching resources. Retrieved January 6, 2008, from <http://www.indiana.edu/~intell>
- Powers, D. (1998) Preparing for the SAT 1: Reasoning Test – An Update, College Board.
- Sabol, W., Couture, H., Harrison, P., (2007) *Prisoners in 2006*, Bureau of Justice Statistics, NCJ 219416, Retrieved February 9, 2008, from: <http://www.ojp.usdoj.gov/bjs/pub/pdf/p06.pdf>
- Schwartz, J. (1994). Low-level lead exposure and children's IQ: A meta-analysis and search for a threshold, *Environmental Research*, 65, 42-55
- Shattuck P. (2006) The Contribution of Diagnostic Substitution to the Growing Administrative Prevalence of Autism in US Special Education, *Pediatrics* 117, 1028-1037, April 2006
- Thomas, V., Socolow, R., Fanelli, J., & Spiro, T. (1999) "Effects of reducing lead in gasoline: An analysis of the international experience," *Environmental Science and Technology*, 33, 3942-3947.
- U.S. Census. (1960-1990) *Statistical Abstract of the United States*.
- U.S. Census (2001). The Hispanic Population 2000.
- U.S. Census. (1975). Historical Statistics of the United States: Colonial Times to 1970. U.S. Government Printing Office.
- U.S. Centers for Disease Control and Prevention. (1991). *Preventing Lead Poisoning in Young Children*, Report No. 99-2230, Atlanta GA, U.S. Department of Health and Human Services.
- U.S. Centers for Disease Control and Prevention, (1997) Update: blood lead levels – United States, 1991-1994, *Morbidity and Mortality Weekly Report*, 46, 141-6.
- U.S. Centers for Disease Control and Prevention, (2000) Blood lead levels in young children – United States and selected states, 1996–1999, *Morbidity and Mortality Weekly Report*, 49, 1133-7.

U.S. Centers for Disease Control and Prevention (2006). Mental Health in the United States: Parental Report of Diagnosed Autism in Children Aged 4-17 Years - United States, 2003-2004, *Morbidity and Mortality Weekly Report*, 55, 481-486.

U.S. Centers for Disease Control and Prevention, (2007a) Interpreting and Managing Blood Lead Levels <10 µg/dL in Children and Reducing Childhood Exposures to Lead, *Morbidity and Mortality Weekly Report*, 56, 1-14.

<http://www.cdc.gov/MMWR/preview/mmwrhtml/rr5608a1.htm#tab1>

U.S. Centers for Disease Control and Prevention, (2007b) *CDC Surveillance Data, 1997-2006*. Retrieved December 21, 2007, from <http://www.cdc.gov/nceh/lead/surv/stats.htm>

U.S. Department of Education (1991) Delivering Special Education: Statistics and Trends, Revised ERIC Digest #E463

U.S. Department of Education (1993) *120 Years of American Education: A Statistical Portrait*.

U.S. Department of Education (2004) *Language Minorities and Their Educational and Labor Market Indicators—Recent Trends*

U.S. Department of Education (1997-2007) *Digest of Education Statistics: 1996- 2006*.

U.S. Department of Education (1995-2007) Office of Special Education Programs Annual Reports to Congress.

U.S. Department of Housing and Urban Development. (1971). *1970 HUD Statistical Yearbook*., U.S. Government Printing Office

U.S. Environmental Protection Agency (1986). *Air Quality Criteria for Lead: Volume I of IV*, Environmental Criteria and Assessment Office, EPA 600/8-83-028 a-d.

U.S. Environmental Protection Agency (2001). National Air Quality and Emissions Trends Report, 1999.