



Munich Personal RePEc Archive

## **Asymmetric Cournot duopoly: game complete analysis**

Carfi, David and Perrone, Emanuele

2012

Online at <https://mpra.ub.uni-muenchen.de/37093/>  
MPRA Paper No. 37093, posted 05 Mar 2012 03:33 UTC

# Asymmetric Cournot Duopoly: Game Complete Analysis

**David Carfi**

Department of Mathematics  
University of California at Riverside  
900 Big Springs Road, Surge 231 Riverside  
CA 92521-0135, USA.  
davidcarfi71@yahoo.it

**Emanuele Perrone**

University of Messina  
Via dei Verdi 75, 98122, Messina, Italy.  
e\_perrone@alice.it

**Abstract.** In this paper we apply the *Complete Analysis of Differentiable Games* (introduced by D. Carfi in [3], [6], [8], [9]; already employed by himself and others in [4], [5], [7]) and some new algorithms using the software *wxMaxima 11.04.0*, in order to reach a **total scenario knowledge (that is the total knowledge of the payoff space of the interaction)** of the classic Cournot Duopoly (1838), viewed as a complex interaction between two competitive subjects, in a particularly interesting **asymmetric case**. The software *wxMaxima* is an interface for the computer algebra system *Maxima*. *Maxima* is a system for the manipulation of symbolic and numerical expressions, including differentiation, sets, vectors and matrices.

**Keywords:** Asymmetric Cournot Duopoly, Normal-form Games, Software algorithms in Microeconomic Policy, Complete Analysis of a normal-form game, Pareto optima, valuation of Nash equilibriums, Bargaining solutions.

**JEL Classification:** D7, C71, C72, C78

## 1. Introduction

The Cournot Duopoly is a classic oligopolistic market in which there are two enterprises producing the same commodity and selling it in the same market. In this classic model, in a competitive background, the two enterprises employ as possible strategies the quantities of the commodity produced. The main solutions proposed in literature for this kind of duopoly are the *Nash equilibrium* and the *Collusive Optimum*, without any subsequent critical exam about these two kinds of solutions. The absence of any critical quantitative analysis is due to the relevant lack of knowledge regarding the set of all possible outcomes of this strategic interaction. On the contrary, by considering the Cournot Duopoly as a *differentiable game* (normal form games with differentiable payoff functions) and studying it by the topological methodologies introduced in Game Theory by D. Carfi, we obtain an exhaustive and complete vision of the entire payoff space of the Cournot game (this also in asymmetric cases with the help of *wxMaxima*) and this total view allows us to analyze critically the classic solutions and to find other ways of action to select Pareto strategies, in the **asymmetric cases too**. In order to illustrate the applications of this topological methodologies to the considered infinite game, several compromise decisions are considered, and we show how the complete study gives a real extremely extended comprehension of the classic model.

## 2. Formal description of a generalized Cournot normal form game

Our model of Cournot game is a non-linear two-players loss game  $G = (f, >)$  (see also [6], [8] and [9]). The two players/enterprises are called *Emil* and *Frances* (following J.P. Aubin's books [1] and [2]).

**Assumption 1 (strategy sets).** The two players produce and offer the same commodity in the quantities  $x \in \mathbb{R}_{\geq}$  for Emil and  $y \in \mathbb{R}_{\geq}$  for Frances. In more precise terms: the payoff function  $f$  of the game  $G$  is defined on a subset of the positive cone of the Cartesian plane  $\mathbb{R}^2$ , interpreted as a space of bi-quantities. We assume (by simplicity) that the set of all strategies, of each player, is the interval  $E = [0, +\infty[$ .

**Assumption 2 (asymmetry of the game).** The game  $G$  is not assumed necessarily symmetric with respect to the players. In other terms, the payoff pair  $f(x, y)$  is not assumed to be the symmetric of the pair  $f(y, x)$ .

**Assumption 3 (form of price function).** We suppose the price function,  $p$  from  $\mathbb{R}^2$  into  $\mathbb{R}$ , linear and defined by

$$p(x, y) = a - b_1x - b_2y \quad (2.1)$$

for every productive bi-strategy  $(x, y)$ , where  $a \geq 0$  is a fixed price and  $b_i > 0$  ( $i = 1, 2$ ) is the marginal coefficient corresponding to the production of the player  $i$ .

**Assumption 4 (form of cost functions).** Let the cost function  $C_1$  (defined on  $E^2$ ) of the first player be given by

$$C_1(x, y) = c_1x + e_1xy + d, \quad (2.2)$$

for every positive price pair  $(x, y)$  and let, analogously, the demand function of the second enterprise be given by

$$C_2(x, y) = c_2y + e_2xy + d, \quad (2.3)$$

for every positive bi-strategy  $(x, y)$ , with  $c > 0$  the marginal cost and  $d \geq 0$  the fixed cost. So we are considering a **Cobb-Douglas perturbation** of the classic linear costs.

**Assumption 5 (payoff functions).** First player's *net cost function* is defined, classically, by the revenue

$$\begin{aligned} f_1(x, y) &= C_1(x, y) - p(x, y)x = \\ &= c_1x + e_1xy + d - (ax - b_1xx - b_2xy) = \\ &= x(b_1x + (b_2 + e_1)y - (a - c_1)) + d = \\ &= x(b_1x + (b_2 + e_1)y - m_1) + d = \\ &= m_1x((b_1/m_1)x + ((b_2 + e_1)/m_1)y - 1) + d, \end{aligned} \quad (2.4)$$

for every positive bi-strategy  $(x, y)$ . Symmetrically, for Frances, the profit function is defined by

$$\begin{aligned} f_2(x, y) &= C_2(x, y) - p(x, y)y = \\ &= y((b_1 + e_2)x + b_2y - (a - c_2)) + d = \\ &= y((b_1 + e_2)x + b_2y - m_2) + d = \\ &= m_2y((b_1 + e_2)/m_2)x + (b_2/m_2)y - 1) + d, \end{aligned} \quad (2.5)$$

for every positive bistrategy  $(x, y)$ .

### 3. Study of the Cournot's normal form game

In the following we shall study the following particular case. We shall put:

$$m_2 = m_1 = 1; \quad b_1 = 2; \quad e_2 = -1; \quad b_2 = 1; \quad e_1 = 0,$$

so that, the bi-loss function is defined by

$$f(x, y) = (x(2x + y - 1), y(x + y - 1)) + (d, d),$$

for every bistrategy  $(x, y)$  of the game in the unbounded square  $E^2$ .

**Payoff function to examine.** When the fixed cost  $d$  is zero (this assumption determines only a “reversible” translation of the loss space), the bi-loss function  $f$  from the compact square  $[0, 1]^2$  into the bi-loss plane  $\mathbb{R}^2$  is defined by

$$f(x, y) = (x(2x + y - 1), y(x + y - 1)), \quad (3.1)$$

for every bistrategy  $(x, y)$  in the square  $S = [0, 1]^2$  which is the convex envelope of its vertices, denoted by A, B, C, D starting from the origin and going anticlockwise.

Now, we must find *the critical space of the game* and its image by the function  $f$ , before representing  $f(S)$ .

For, we determine (as explained in [3], [6], [8] and [9]) firstly the *Jacobian matrix* of the function  $f$  at any point  $(x, y)$  - denoted by  $J_f(x, y)$ . We will have, in vector form, the pair of gradients

$$J_f(x, y) = ((y + 4x - 1, x), (y, 2y + x - 1)), \quad (3.2)$$

and concerning the determinant of the above pair of vectors

$$\begin{aligned} \det J_f(x, y) &= (y + 4x - 1)(2y + x - 1) - xy = \\ &= 2y^2 + 8xy - 3y + 4x^2 - 5x + 1. \end{aligned} \quad (3.3)$$

The *Jacobian determinant* is zero at those points  $(x_1, y_1)$  and  $(x_2, y_2)$  of the strategy square such that

$$y_1 = -\sqrt{32x^2 - 8x + 1}/4 - 2x + 3/4, \quad (3.4)$$

and

$$y_2 = \sqrt{32x^2 - 8x + 1}/4 - 2x + 3/4. \quad (3.5)$$

We obtain two curves (Figure 3.1) whose union is *the critical zone of Cournot Game*.




Figure 3. 1. Critical zone of Cournot game

#### 4. Transformation of the strategy space

It is readily seen that the intersection points of the green curve with the boundary of the strategic square are the point  $K = (2/8, 0)$ .

**Remark.** The point  $K$  is the intersection point of the **green curve** with the **segment [A, B]**, its abscissa  $\mu$  verifies the non-negative condition and the following equation

$$\sqrt{32\mu_1^2 - 8\mu_1 + 2} = 3 - 8\mu_1, \quad (4.1)$$

this abscissa is so  $\mu_1 = 2/8$  (equal to 0,25).

We start from Figure 3.1.

The transformations of the bi-strategy square vertices and of the points  $H, K$  are the following:

- $A' = f(A) = f(0, 0) = (0, 0)$ ;
- $B' = f(B) = f(1, 0) = (1, 0)$ ;
- $C' = f(C) = f(1, 1) = (2, 1)$ ;
- $D' = f(D) = f(0, 1) = (0, 0)$ ;
- $H' = f(H) = f(0, 1/2) = (0, -1/4)$ ;
- $K' = f(K) = f(2/8, 0) = (-1/8, 0)$ .

Starting from Figure 3.1, with  $S = [0, 1]^2$ , we can do the transformation of its sides.

**Side [A, B].** Its parameterization is the function sending any point  $x \in [0, 1]$  into the point  $(x, 0)$ ; the transformation of this side can be obtained by transformation of its generic point  $(x, 0)$ , we have

$$f(x, 0) = (2x^2 - x, 0). \quad (4.2)$$

We obtain the segment with end points  $K'$  and  $B'$ , with parametric equations

$$X = 2x^2 - x \text{ and } Y = 0, \quad (4.3)$$

with  $x$  in the unit interval.

**Side [B, C].** Its parameterization is

$$(x = 1, y \in [0, 1]);$$

the figure of the generic point is

$$f(1, y) = (y + 1, y^2). \quad (4.4)$$

We can obtain the parabola passing through the points  $B', C'$  with parametric equations

$$X = y + 1 \text{ and } Y = y^2. \quad (4.5)$$

**Side [C, D].** Its parameterization is

$$(x \in [0, 1], y = 1);$$

the transformation of its generic point is

$$f(x, 1) = (2x^2, x). \quad (4.6)$$

We obtain the parabola passing through the points  $D', C'$  with parametric equations

$$X = 2x^2 \text{ and } Y = x, \quad (4.7)$$

with  $x$  in the unit interval.

**Side [A, D].** Its parameterization is

$$(x = 0, y \in [0, 1]);$$

the transformation of its generic point is

$$f(0, y) = (0, y^2 - y). \quad (4.8)$$

We obtain the segment with end points A' and H', with parametric equations

$$X = 0 \text{ and } Y = y^2 - y. \quad (4.9)$$

Now, we find *the transformation of the critical zone*. The parameterization of the critical zone is defined by the equations

$$y_1 = -\sqrt{(32x^2 - 8x + 1)/4 - 2x} + 3/4, \quad (\text{see (3.5)})$$

and

$$y_2 = \sqrt{(32x^2 - 8x + 1)/4 - 2x} + 3/4. \quad (\text{see (3.6)})$$

The parametrization of the **GREEN ZONE** is

$$(x \in [0, 1], y = y_1);$$

the transformation of its generic point is

$$f(x, y_1) = (x(2x + y_1 - 1), y_1(x + y_1 - 1)), \quad (4.10)$$

a parametrization of the **YELLOW ZONE** is

$$(x \in [0, 1], y = y_2);$$

the transformation of its generic point is

$$f(x, y_2) = (x(2x + y_2 - 1), y_2(x + y_2 - 1)). \quad (4.11)$$

The transformation of the Green Zone has parametric equations

$$X = x(-\sqrt{(32x^2 - 8x + 1)/4 - 1/4}) \quad (4.12)$$

and

$$Y = (-\sqrt{(32x^2 - 8x + 1)/4 - 2x} + 3/4) * (-\sqrt{(32x^2 - 8x + 1)/4 - x - 1/4}), \quad (4.13)$$

and the transformation of the Yellow Zone has parametric equations

$$X = x(\sqrt{(32x^2 - 8x + 1)/4 - 1/4}) \quad (4.14)$$

and

$$Y = (\sqrt{(32x^2 - 8x + 1)/4 - 2x} + 3/4) * (\sqrt{(32x^2 - 8x + 1)/4 - x - 1/4}). \quad (4.15)$$

We have two colored curves in *green* and *black* (Figure 4.1). The *black* curve is break by a point of discontinuity  $T$  obtained by resolving the following equation

$$x(\sqrt{32x^2 - 8x + 1})/4 - 1/4 = 0; \tag{4.16}$$

the solutions of the above equation are

$$x_1 = 1/4, x_2 = 0, \tag{4.17}$$

and then, replacing them in the parametrical equations of the critical zone numbered as 4.15 we obtain

$$T_1 = 0$$

and

$$T_2 = -1/8$$




Figure 4.1. Payoff space of Cournot game

## 5. Non-cooperative friendly phase

Examining the Figure 4.1 we will note that the game has two extremes, that is the *shadow minimum*  $\alpha = (-1/8, -1/4)$  and the *maximum*  $\beta = C' = (2, 1)$ .

The *Pareto minimal boundary of the payoff space*  $f(S)$  is the curve passing through the points  $K'$  and  $H'$  colored in *green* showed in the Figure 5.1.

The *Pareto maximal boundary of the payoff space*  $f(S)$  coincides with the *maximum*  $\beta = C' = (2, 1)$ .

Both Emil and Frances do not control the Pareto minimal boundary; they could reach the point  $K'$  and  $H'$  of the boundary, but the solution is not many satisfactory for them. In fact, a player will suffer the maximum loss.




Figure 5. 1. Extrema of the Asymmetric Cournot game




Figure 5. 2. Extrema of the Symmetric Cournot game

**Remark.** Comparing the figure 5.1 with the figure 5.2 we may observe that the benefit to the community in case of asymmetry decreases; in fact the area contained in the first quadrant is greater than the symmetric case and the area contained in the third quadrant is smaller than the symmetric one. In other terms, when the Cournot duopoly becomes an asymmetric games is easier to have a loss.



## 6. Properly non-cooperative (egoistic) phase

Now, we will consider *the best reply correspondences* between the two players Emil and Frances.

If Frances produces the quantity  $y$  of the commodity, Emil, in order to reply rationally, should minimize his *partial cost function*

$$f_1(\cdot, y) : x \mapsto x(2x + y - 1), \quad (6.1)$$

on the compact interval  $[0, 1]$ .

According to the Weierstrass theorem, there is at least one Emil's strategy minimizing that partial cost function and, by Fermàt theorem, the *Emil's best reply strategy to Frances' strategy*  $y$  is the only quantity

$$B_1(y) = x^* := 1/4(1 - y). \quad (6.2)$$

Indeed, the partial derivative

$$f_1(\cdot, y)'(x) = 4x + y - 1, \quad (6.3)$$

is negative for  $x < x^*$  and positive for  $x > x^*$ .

So, the Emil's best reply correspondence is the function  $B_1$  from the interval  $[0, 1]$  into the interval  $[0, 1]$ , defined by

$$y \mapsto 1/4(1 - y). \quad (6.4)$$

If Emil produces the quantity  $x$  of the commodity, Frances, in order to reply rationally, should minimize his *partial cost function*

$$f_2(x, \cdot) : y \mapsto y(x + y - 1), \quad (6.5)$$

on the compact interval  $[0, 1]$ .

According to the Weierstrass theorem, there is at least one Frances' strategy minimizing that partial cost function and, by Fermàt theorem, the *Frances' best reply strategy to Emil's strategy*  $x$  is the only quantity

$$B_2(x) = y^* := 1/2(1 - x). \quad (6.6)$$

Indeed, the partial derivative

$$f_2(x, \cdot)'(y) = 2y + x - 1, \quad (6.7)$$

is negative for  $x < x^*$  and positive for  $x > x^*$ .

So, the Frances' best reply correspondence is the function  $B_2$  from the interval  $[0, 1]$  into the interval  $[0, 1]$ , defined by  $x \mapsto 1/2(1 - x)$ .

The *Nash equilibrium* is the fixed point of the multifunction  $B$  - associated with the pair of two reaction functions  $(B_2, B_1)$  - defined from the Cartesian product of the domains into the Cartesian product of the codomains (in inverse order), by  $B : (x, y) \mapsto (B_1(y), B_2(x))$ , that is the only bi-strategy  $(x, y)$  such that

$$x = 1/4(1 - y), y = 1/2(1 - x), \quad (6.8)$$

that is the point  $N = (1/7, 3/7)$  - as we can see also from Fig. 6.1 - which gives a bi-loss  $N' = (-2/49, -9/49)$ , as Figures 6.2 will show. The Nash equilibrium is not completely satisfactory, because it is not a Pareto optimal bi-strategy, but it represents the only properly non-cooperative game solution.




Figure 6. 1. Nash Equilibrium of Cournot game




Figure 6. 2. Payoff at Nash equilibrium of Cournot game

## 7. Defensive and offensive phase

Players' conservative values are obtained through their *worst loss functions*.

**Worst loss functions.** On the square  $S = [0, 1]^2$ , *Emil's worst loss function* is defined by

$$f^{\#_1}(x) = \sup_{y \in F} x(2x + y - 1) = 2x^2, \quad (7.1)$$

its minimum will be

$$v^{\#_1} = \inf_{x \in E} (f^{\#_1}(x)) = \inf_{x \in E} (2x^2) = 0, \quad (7.2)$$

attained at the conservative strategy  $x^{\#} = 0$ .

*Frances' worst loss function* is defined by

$$f^{\#_2}(y) = \sup_{x \in E} y(x + y - 1) = y^2, \quad (7.3)$$

its minimum will be

$$v^{\#_2} = \inf_{y \in F} (f^{\#_2}(y)) = \inf_{y \in F} (y^2) = 0 \quad (7.4)$$

attained at the unique conservative strategy  $y^{\#} = 0$ .

**Conservative bivalue.** The *conservative bivalue* is

$$v^{\#} = (v^{\#_1}, v^{\#_2}) = (0, 0).$$

*The worst offensive multifunctions* are determined by the study of the *worst* loss functions.

The Frances' worst offensive reaction multifunction  $O_2$  is defined by  $O_2(x) = 1$ , for every Emil's strategy  $x$ ; indeed, fixed an Emil's strategy  $x$  the Frances' strategy 1 maximizes the partial cost function  $f_1(x, \cdot)$ . The Emil's worst offensive correspondence versus Frances is defined by  $O_1(y) = 1$ , for every Frances' strategy  $y$ .

*The dominant offensive strategy* is 1 for both players, indeed the offensive correspondences are constant.

*The core of the payoff space* (in the sense introduced by J.P. Aubin) is the Pareto minimal boundary, contained in the cone of lower bounds of the conservative bi-value  $v^{\#}$ ; the conservative bi-value don't give us a bound for the choice of cooperative bistrategies.

*The conservative knot* of the game is the point  $(0, 0)$ .

## 8. Cooperative phase

When there is an agreement between the two players, *the best compromise solution (in the sense introduced by J.P. Aubin)* showed graphically in the Figures 8.1.

Besides, the best compromise solution coincides with the *core best compromise*, with the *Nash bargaining solution*, with the *Kalai-Smorodinsky bargaining solution*. It coincides also with *the transferable utility solution* which is the unique Pareto strategy that minimized the aggregate utility function  $f_1 + f_2$ , this can be easily viewed by geometric evidences considering on the payoff universe the levels of that aggregate function, which are affine lines parallel to the vector  $(1, -1)$ .




Figure 8. 1. Conservative study. The core and the Kalai-Smorodinsky payoff of Cournot game

## References

1. Aubin Jean Pierre, *Mathematical Methods of Game and Economic Theory* (Revised Edition), (North-Holland).
2. Aubin Jean Pierre, *Optima and Equilibria*, (Springer Verlag, 1998).
3. Carfi David, *Topics in Game Theory*, (Edizioni Il gabbiano, 2010).
4. Carfi David, Schilirò Daniele, *Crisis in the Euro area: cooperative game solutions as new policy tools*, TPREF - Theoretical and Practical Research in Economic Fields, Summer issue 2011  
<http://mpra.ub.uni-muenchen.de/27138/>
5. Carfi David, Ricciardello Angela, *An algorithm for payoff space in  $C^1$  games*, AAPP | Physical, Mathematical, and Natural Sciences, Vol. LXXXVIII, issue 1, 2010  
<http://cab.unime.it/journals/index.php/AAPP/article/view/C1A1001003/>
6. Carfi David, *Differentiable game complete analysis for tourism firm decisions*, Proceedings of THE 2009 INTERNATIONAL CONFERENCE ON TOURISM and WORKSHOP on Sustainable tourism within High Risk areas of environmental crisis, Messina, April 22/25 (2009)  
<http://mpra.ub.uni-muenchen.de/29193/>
7. Carfi David, *Globalization and Differentiable General Sum Games*, Proceeding of 3rd International Symposium Globalization and convergence in economic thought, Faculty of Economics - Communication and Economic Doctrines Department - The Bucharest Academy of Economic Studies - Bucharest December 11/12 (2009)
8. Carfi David, *Payoff space and Pareto boundaries of  $C^1$  Games*, APPS vol. 2009  
<http://www.mathem.pub.ro/apps/v11/a11.htm>
9. Carfi David, *Complete study of linear infinite games*, Proceedings of the International Geometry Center - Proceeding of the International Conference "Geometry in Odessa 2009", Odessa, 25 - 30 May 2009 - vol. 2 n. 3 (2009)  
<http://proceedings.d-omega.org/>
10. Carfi David, *Optimal boundaries for decisions*, Atti dell'Accademia Peloritana dei Pericolanti - Classe di Scienze Fisiche, Matematiche e Naturali Vol. LXXXVI, (2008)  
<http://cab.unime.it/journals/index.php/AAPP/article/view/376>
11. Carfi David, Ricciardello Angela, *Non-reactive strategies in decision-form games*, AAPP | Physical, Mathematical, and Natural Sciences, Vol. LXXXVII, issue 1, pp. 1-18  
<http://cab.unime.it/journals/index.php/AAPP/article/view/C1A0902002/0>
12. Carfi David, *Decision-form games*, Proceedings of the IX SIMAI Congress, Rome, 22 - 26 September 2008, Communications to SIMAI congress, vol. 3, (2009) pp. 1-12  
<http://cab.unime.it/journals/index.php/congress/article/view/307>
13. Carfi David, *Reactivity in decision-form games*, MPRA - paper 29001, University Library of Munich, Germany  
<http://mpra.ub.uni-muenchen.de/29001/>
14. Carfi David, Ricciardello Angela, *Mixed Extensions of decision-form games*, MPRA - paper 28971, University Library of Munich, Germany  
<http://mpra.ub.uni-muenchen.de/28971/>