



Munich Personal RePEc Archive

Bayesian Averaging of Classical Estimates in Asymmetric Vector Autoregressive (AVAR) Models

Albis, Manuel Leonard F. and Mapa, Dennis S.

School of Statistics, University of the Philippines Diliman

2014

Online at <https://mpra.ub.uni-muenchen.de/55902/>
MPRA Paper No. 55902, posted 12 May 2014 07:03 UTC

Bayesian Averaging of Classical Estimates in Asymmetric Vector Autoregressive (AVAR) Models¹

Manuel Leonard F. Albis² and Dennis S. Mapa, PhD.³
School of Statistics, University of the Philippines Diliman, Quezon City

ABSTRACT

The estimated Vector AutoRegressive (VAR) model is sensitive to model misspecifications, such as omitted variables, incorrect lag-length, and excluded moving average terms, which results in biased and inconsistent parameter estimates. Furthermore, the symmetric VAR model is more likely misspecified due to the assumption that variables in the VAR have the same level of endogeneity. This paper extends the Bayesian Averaging of Classical Estimates, a robustness procedure in cross-section data, to a vector time-series that is estimated using a large number of Asymmetric VAR models, in order to achieve robust results. The combination of the two procedures is deemed to minimize the effects of misspecification errors by extracting and utilizing more information on the interaction of the variables, and cancelling out the effects of omitted variables and omitted MA terms through averaging. The proposed procedure is applied to simulated data from various forms of model misspecifications. The forecasting accuracy of the proposed procedure was compared to an automatically selected equal lag-length VAR. The results of the simulation suggest that, under misspecification problems, particularly if an important variable and MA terms are omitted, the proposed procedure is better in forecasting than the automatically selected equal lag-length VAR model.

Keywords: BACE, AVAR, Robustness Procedures

¹ The authors are grateful to the Statistical Research and Training Center (SRTC) for the thesis fellowship grant for the completion of this study, and to the participants of the Colloquium on the Statistical Sciences at the School of Statistics, University of the Philippines, for their valuable comments.

² Assistant Professor, School of Statistics, University of the Philippines Diliman.
Email Address: manuel_albis@hotmail.com

³ Associate Professor and Director for Research, School of Statistics and Affiliate Associate Professor of Economics, School of Economics, University of the Philippines Diliman

1. Introduction

The Vector Autoregressive (VAR) model by Sims (1980) became a popular tool for forecasting a group of interrelated economic variables because of its ease of use. However, Braun & Mittnik (1993) showed that the ordinary least squares (OLS) coefficients VAR estimates are sensitive to misspecification errors due to omitted variables, incorrect lag-length, and excluded moving average (MA) terms. This results in having biased and inconsistent estimators and creates problems in forecasting and the estimation of the impulse response function (IRF) and variance decompositions. If these problems are not considered in the modeling procedure, then the results of the VAR model may be misleading. A certain degree of caution must be emphasized for the purpose of policy and decision making under these circumstances.

The effects of excluded MA terms in the VAR model are alleviated by using a large number of lags. However, the problem of omitting an important variable in the VAR model is the hardest to solve. This problem is common in practice partly because of the true model is usually unknown. .

Furthermore, the VAR model itself is misspecified. It assumes the lags of all variables in the system are the same or symmetric. This is a problem in applied research since variables tend to have different degrees of endogeneity. Keating (1993, 1995 & 2000) addressed this problem by allowing unequal lag length or asymmetry in the VAR model (AVAR).

Another way of dealing with these problems is to use model averaging that is deemed to produce robust results under problems of model misspecifications. Strachan & van Dijk (2007) were the first to apply Bayesian Model Averaging (BMA) on VAR. The authors assumed prior distribution for each parameter in the model. In analyzing cross-section data, Sala-i-Martin (1997) proposed the Averaging of Classical Estimates (ACE) that uses the

likelihood function of the regression model as weights in averaging the OLS estimates, where the average is taken across all models generated in the context of the Extreme Bounds Analysis of Leamer (1983). Sala-i-Martin, Doppelhofer & Miller (2004) formulated the Bayesian Averaging of Classical Estimates (BACE) that uses the posterior model probability as weights for the OLS estimates that needs only one prior information – the number of variables in the true model.

The main objective of this paper is to develop a modeling procedure that will yield robust VAR forecasts by the use of BACE on the forecasts of AVAR models using less assumption, particularly on the parameter's prior distributions of popular Bayesian VAR methods. The combination of the two procedures, the BACE and the AVAR, is expected to minimize the effects of misspecification errors by extracting and utilizing more information on the interaction of the variables, and cancelling out the effects of omitted variables and omitted MA terms through averaging. The paper also aims to determine the forecasting performance of the BACE-AVAR method by applying it to stationary and deseasonalized vector of variables simulated from different data characteristics. The Modified Diebold-Mariano test and the relative MAPE will be used in the forecasting accuracy of the BACE-AVAR procedure with respect to a model with automatic selection procedure.

2. VAR, AVAR and BACE Procedures

This section will provide a background on VAR and AVAR models, how these models are specified and estimated, and how to measure their predictive accuracy. The BACE procedure in the context of cross-section data is also discussed.

2.1 Vector Autoregressive Moving Average (VARMA) Models

Following Lütkepohl (2004), consider the generalized form of the finite order VARMA(p, q) model that is given by:

$$\sum_{i=0}^p \mathbf{A}_i^* \mathbf{y}_{t-i} = \sum_{i=0}^q \mathbf{M}_i^* \mathbf{u}_{t-i} \quad (1)$$

where $\mathbf{y}_t = (y_{1t}, \dots, y_{Kt})'$, $t = 1, \dots, T$, is stationary, \mathbf{A}_i^* and \mathbf{M}_i^* are ($K \times K$) autoregressive and moving average coefficient matrices, respectively, and $\mathbf{u}_t = (u_{1t}, \dots, u_{Kt})'$ is a K -dimensional white noise process, that is $E(\mathbf{u}_t) = \mathbf{0}$ and

$$E(\mathbf{u}_t \mathbf{u}_h') = \begin{cases} \boldsymbol{\Sigma}_u, & \text{if } t = h \\ 0 & \text{otherwise,} \end{cases} \quad (2)$$

and $\boldsymbol{\Sigma}_u$ is positive definite.

Due to the difficulties in estimating a VARMA(p, q) model, it is a common practice among researchers to estimate VARMA($p, 0$) model, which is popularly known as the VAR(p) model. The VAR(p)⁴ model is commonly represented by:

$$\mathbf{y}_t = \sum_{i=1}^p \mathbf{A}_i^* \mathbf{y}_{t-i} + \mathbf{u}_t \quad (3)$$

where the terms are as defined in Equation (1). The parameters are usually estimated using ordinary least squares (OLS) for all equations in the system.

⁴ The VAR operator is stable and the process is stationary if $\det \mathbf{A}^*(z) \neq 0$, where $z \in \mathbb{C}$. If this is the case, then the VAR(p) model can also be expressed as $\mathbf{y}_t = \sum_{i=0}^{\infty} \boldsymbol{\Phi}_i \mathbf{u}_{t-i}$, where $\boldsymbol{\Phi}_0 = \mathbf{I}_K$, if $\mathbf{A}_0^* = \mathbf{M}_0^* = \mathbf{I}_K$, and $\boldsymbol{\Phi}_i = \sum_{j=1}^i \boldsymbol{\Phi}_{i-j} \mathbf{A}_j^*$, $i = 1, 2, \dots$, with $\mathbf{A}_j^* = 0$ for $j > p$. The $\boldsymbol{\Phi}_i$'s are popularly known as the impulse response function in the literature. In practice, researchers use the orthogonalized form of the IRF that can be expressed by $\boldsymbol{\Phi}_i^o = \boldsymbol{\Phi}_i \mathbf{L}$ where \mathbf{L} is a lower triangular matrix of the Cholesky decomposition of $\boldsymbol{\Sigma}$, that is $\boldsymbol{\Sigma} = \mathbf{L} \mathbf{L}'$. The interpretations of the VAR(p) model is coursed through the estimated IRF as it gives the reaction of the value of a variable when there is an abrupt change in the other variables.

Another approach in estimating the VAR model parameters is through the Bayesian VAR, which was introduced by Litterman (1980). This approach was extensively used in modeling and forecasting economic variables.⁵ The BVAR procedure involves setting the prior distributions of the parameters and running MCMC simulations. Sun & Ni (2003; 2004) indicated that the use of the non-informative Jeffrey’s prior in BVAR is likely to have over-estimated posterior mean variance. Their study also showed that the results of BVAR across different priors were different. This indicates that results of the BVAR are sensitive to the selected prior information.

2.2 Automatic Selection Procedure

In practice, the model builder usually starts with VAR(p^*) model where p^* is selected using an automatic selection procedure. This involves the estimation of all VAR(p) model for $p = 1, \dots, p^*$, and selecting the initial model that yields the “best” value of a pre-selected information criterion. The common information criteria are the Akaike Information Criterion (1973) that is given by:

$$AIC = \log|\hat{\Sigma}| + 2\frac{r}{T} \quad (4)$$

and the Bayesian Information Criterion by Schwarz (1978) that is of the form:

$$BIC = \log|\hat{\Sigma}| + \frac{r \log T}{T} \quad (5)$$

where r is the number of estimated parameters, K is the number of dependent variables in the vector, and T is the sample size. Hurvich & Tsai (1993) corrected the AIC for small samples and it has the form:

⁵ Some of the studies that used BVAR and variants of it are from Po, Chi, Shyu, & Hsiao (2002), Chen & Leung (2003), Ramos (2003), Carriero, Kapetanios, & Marcellino (2009).

$$AICc = \log|\hat{\Sigma}| + 2\frac{r}{T - r/K}. \quad (6)$$

Kadilar & Eldemir (2002) analyzed the performance of the popular information criteria by simulating VAR(1) and VAR(2) models, with and without seasonality. They showed that performance of the information criteria is better in VAR without seasonality than VAR with seasonality. They also noted the improvement in the performance of AIC as the number of variables in the VAR model, without seasonality, increases. However, the result for the AIC is reversed for VAR with seasonal data. Hence, the authors recommended not to use the AIC in the presence of seasonality in the VAR data. Overall, they ranked the performance of the information criteria from highest to lowest as: Schwarz (SIC), Hannan-Quinn (HQ), Akaike (AIC).

Waele & Broersen (2003) noted that the AIC is an unbiased estimate of the Kullback-Leibler discrepancy.⁶ However, for a finite sample size, it tends to over-fit the model by choosing a high number of lags, as discussed earlier. They also showed that the Kullback-Leibler discrepancy can be used as an information criterion which they call KIC. Seghouane (2006) proposed a refinement to the KIC, which the author called KICvc, where vc stands for vector correction. The KICvc performs better than the KIC in model specification for small sample sizes.

George, Sun & Ni (2004) developed a Bayesian stochastic search approach in determining the VAR model that can incorporate restrictions on the VAR coefficients and on the elements of the error covariance matrix. Korobilis (2010) developed an automatic variable selection procedure using the Gibbs sampler for linear and nonlinear VARs. Numerical simulations

⁶ For an in-depth discussion of the Kullback-Leibler discrepancy, see *“On Information and Sufficiency”* by Kullback & Leibler (1951) and *“Finite sample effects in vector autoregressive modeling”* by Waele & Broersen (2002).

indicated that both procedures select a satisfactory model with improved forecasting performance.

2.3 Asymmetric Vector Autoregressive (AVAR) Models

Hsiao (1981) was the first to suggest the estimation of VAR models with variables having unequal lag length. However, Keating (1993) argued that Hsiao's method depends on the inclusion sequence of the explanatory variables in the model and that Litterman's Bayesian (Litterman, 1980) approach gives biased parameter estimates, a minor issue in forecasting but a potential problem in determining macroeconomic structures. Keating introduced asymmetries in the lag lengths of the variables in the VAR system and named this as Asymmetric VAR (AVAR) model. The AVAR(p_1, \dots, p_k), can be written as

$$\mathbf{y}_t = \sum_{i=1}^{p^*} \mathbf{D}\mathbf{A}_i^* \mathbf{y}_{t-i} + \mathbf{u}_t \quad (7)$$

where $p^* = \max\{p_1, p_2, \dots, p_k\}$; $\mathbf{D} = \text{diag}\{1_{\{i \leq p_1\}}, 1_{\{i \leq p_2\}}, \dots, 1_{\{i \leq p_k\}}\}$, a diagonal matrix having indicator variables as elements such that $1_{\{i \leq p_j\}} = \{1 \text{ if } i \leq p_j; 0 \text{ otherwise}\}$; the rest are as previously described. The \mathbf{D} matrix restricts some of the parameters to zero and this matrix introduces the inequalities in the lag-length of the variables.

The AVAR model is a VAR model that permits unequal lag length for the variables in the equations. However, the lag specification should be the same across all equations in the system. Because of this, the AVAR gives a parsimonious model with a substantial reduction in the standard errors compared to the ordinary VAR. This translates to the clarity in the interpretations from the impulse response function and variance decompositions.

Keating also performed an automatic selection procedure over a set of AVAR models. For a vector of variables, the procedure estimates all possible AVAR models given a maximum

number of lags L , and for each estimated model, the selected information criterion is computed. The best AVAR model with the best information criterion is selected. There are L^k number of AVAR models needed to be estimated in the procedure. For convenience, the values of the AIC, SIC and HQ that are computed using Keating's procedure will be called KAIC, KSIC and KHQ respectively.⁷

Ozcicek & McMillin (1999) studied the performance of the popular information criteria, such as AIC, SIC, KAIC and KSIC, in determining the lag length of a VAR model. Using a variety of autoregressive data structure such as either short or long-lagged process, and symmetric and asymmetric lag lengths, the authors showed that AIC is best for symmetric data, since the other information criteria under-fits the model. The authors also showed that KAIC is the best criterion to use for asymmetric data and they proposed this criterion to be used in modeling since the lag length structure of the data is uncertain and most of the time asymmetric in theory.

2.4 Predictive Accuracy

Diebold and Mariano (1995) developed a test for predictive accuracy in forecasting that is not restricted to the quadratic loss function and can handle a wide variety of error characteristics. For the two forecasts $\{\hat{y}_{1t}\}_{t=1}^T$ and $\{\hat{y}_{2t}\}_{t=1}^T$ for the series $\{y_t\}_{t=1}^T$, let $\{e_{1t}\}_{t=1}^T$ and $\{e_{2t}\}_{t=1}^T$ be the associated forecast errors. The loss associated with a forecast at time t is given by the loss function $g(y_t, \hat{y}_{it})$ and the authors pointed out that the loss function is a direct function of the forecast errors, that is, $g(y_{it}, \hat{y}_{it}) = g(e_{it})$. The null hypothesis of the Diebold-Mariano test is $E(d_t) = 0$, where $d_t = g(e_{1t}) - g(e_{2t})$ is the loss differential. So

⁷ The names KAIC and KSIC are adapted from Ozcicek & McMillin (1999).

that if we have $\{d_t\}_{t=1}^T$, then under the assumption that the loss differential series is covariance stationary and short memory,

$$\sqrt{T}(\bar{d} - \mu) \xrightarrow{d} N(0, 2\pi f_d(0)), \quad (8)$$

where, $\bar{d} = \frac{1}{T} \sum_{i=1}^T d_t$ and $f_d(0) = \frac{1}{2\pi} \sum_{\tau=-\infty}^{\infty} \gamma_d(\tau)$, the spectral density of the loss differential at frequency zero, having $\gamma_d(\tau) = E\{(d_t - \mu)(d_{t-\tau} - \mu)\}$, the sample autocovariance of the loss differential to lag τ , and μ is the population mean of the loss differential.

Harvey, Leybourne and Newbold (HLN) (1997) corrected the Diebold-Mariano (DM) test for finite samples. The h -step ahead forecasts DM test statistic is given by:

$$DM = \bar{d}[\hat{V}(\bar{d})]^{-\frac{1}{2}} \quad (9)$$

where $\hat{V}(\bar{d})$ is the estimated variance of \bar{d} that is given by

$$\hat{V}(\bar{d}) \approx \frac{1}{T} \left[\hat{\gamma}_0 + 2 \sum_{k=1}^{h-1} \hat{\gamma}_k \right] \quad (10)$$

and, $\hat{\gamma}_k$ is the estimated autocovariance of \bar{d} that has the form:

$$\hat{\gamma}_k = \frac{1}{T} \sum_{t=k+1}^T (d_t - \bar{d})(d_{t-k} - \bar{d}) \quad (11)$$

The h -step ahead forecasts Modified DM test (MDM) is of the form:

$$MDM = \left[\frac{T + 1 - 2h + T^{-1}h(h-1)}{T} \right]^{\frac{1}{2}} DM \quad (12)$$

having a Student's t distribution with $T - 1$ degrees of freedom. The MDM test statistic was used in determining the forecasting performance of the BACE-AVAR procedure against the

forecasting performance of the VAR model that is selected automatically using an information criterion.

The Mean Absolute Percentage Error (MAPE) is a descriptive measure of predictive accuracy that is given by:

$$MAPE = \frac{1}{T} \sum_{t=1}^T \left| \frac{y_t - \hat{y}_t}{y_t} \right| \quad (13)$$

where $|\cdot|$ is the absolute value function. The MAPE was used in the study to measure the distance between the actual and predicted forecasts. The ratio of two MAPE is called the relative MAPE and is given by the form

$$RelMAPE_i = \frac{MAPE_i}{MAPE^*} \quad (14)$$

where $MAPE_i$ is from the i^{th} model and $MAPE^*$ is from the baseline model. Relative MAPE of less than 1 implies that the model that is being evaluated is better than the baseline model.

2.5 Bayesian Averaging of Classical Estimates

The BACE by Sala-i-Martin, et. al. (2004) computes the weighted average of the OLS coefficient estimates weighted by the probability that the model where it is estimated from is the true model. This approach also has an advantage over BMA, since it only needs the number of variables in the model as prior information under the assumption of equal prior inclusion probabilities for each variable, whereas the BMA must be given assumed prior distributions for all of the parameters.

The procedure involves estimating all regression models of the form

$$y = \beta_0 + \beta_{zj}z + \boldsymbol{\beta}'_{fj}\mathbf{f} + \boldsymbol{\beta}'_{xj}\mathbf{x}_j + \epsilon \quad (15)$$

where z is the variable of interest, \mathbf{f} is a vector of fixed variables that appear in all the regressions, and $\mathbf{x}_j \in \mathfrak{X}$ is a vector of variables taken from the \mathfrak{X} collection of all other variables under consideration.

If it is assumed that the prior inclusion probability of each variable in the model are equal, the prior probability of model j , denoted as $P(M_j)$, will be:

$$P(M_j) = \left(\frac{\bar{k}}{K}\right)^{k_j} \left(1 - \frac{\bar{k}}{K}\right)^{K-k_j} \quad (16)$$

where \bar{k} is the speculated number of variables in the true model, K is the total number of variables in the dataset, and k_j is the number of variables in the j^{th} model.

The weights that will be used in the averaging is the posterior probabilities of the M_j 's. The weight is a function of the prior probability and is given by:

$$P(M_j|y) = \frac{P(M_j)n^{-k_j/2} SSE_j^{-n/2}}{\sum_{i=1}^{2^k} P(M_i)n^{-k_i/2} SSE_i^{-n/2}} \quad (17)$$

where the SSE_j is the sum of squared errors in model j . Therefore, the posterior mean of $\boldsymbol{\beta}$ is given by:

$$E(\boldsymbol{\beta}|y) = \sum_{j=1}^{2^k} P(M_j|y)\hat{\boldsymbol{\beta}}_j \quad (18)$$

where $\hat{\boldsymbol{\beta}}_j$ is the estimated value of the vector of coefficients under OLS; and its corresponding posterior variance is of the form:

$$Var(\boldsymbol{\beta}|y) = \sum_{j=1}^{2^k} P(M_j|y)Var(\boldsymbol{\beta}|y, M_j) + \sum_{j=1}^{2^k} P(M_j|y)[\hat{\boldsymbol{\beta}}_j - E(\boldsymbol{\beta}|y)]^2 \quad (19)$$

3. The BACE-AVAR Procedure

In specifying the AVAR model, the procedure of Keating (1995) estimates L^K AVAR models given a maximum lag length of L that is set by the researcher. The best specification will be selected based on the model that gives the best value of an information criterion.

The prior probability $P(\mathbf{M}_j)$ will be assumed to be equal for all the equations in the j^{th} estimated VAR model. The formula for the prior probability for the AVAR model \mathbf{M}_j is given by:

$$P(\mathbf{M}_j) = \binom{L}{l_j} \left(\frac{\bar{l}}{L}\right)^{l_j} \left(1 - \frac{\bar{l}}{L}\right)^{L-l_j} \quad (20)$$

where, l_j is the total number of AR lag regressors for each of the equations in the j^{th} model, and \bar{l} is the assumed total number of lags of all the variables in the true model. To simplify the procedure, the value for \bar{l} can be given by running an automatic selection procedure over VAR models and setting \bar{l} based on the recommended number of lags. Alternatively, the researcher may run BACE-AVAR using a different \bar{l} .

The formula for the posterior probability will be:

$$P(M_{jq} | y) = \frac{P(\mathbf{M}_j) T^{-l_j/2} SSE_j^{-T/2}}{\sum_{i=1}^{L^K} P(\mathbf{M}_i) T^{-l_i/2} SSE_i^{-T/2}} \quad (21)$$

where SSE_j is the sum of squared errors of the AVAR model that is given by $SSE_j = \sum_{k=1}^K \sum_{i=1}^T (y_{ki} - \hat{y}_{ki})^2$.⁸ This will give a single weight for an estimated AVAR model depending on the ability of all its equations to fit their corresponding variables.

⁸ See Section 4.1 for the explanation in the power of SSE_j from $-T/2$ to $-0.1T/2$.

3.1 BACE-AVAR on Forecasted Values

The posterior probabilities will also be used in computing for the forecasts of the BACE-AVAR procedure. The forecasts will be the weighted mean of the forecast series produced by the AVAR models, with the posterior probabilities as its weights. The $(T + t)^{th}$ forecast for the vector of variables \mathbf{y}_{T+t} from the BACE-AVAR procedure is given by:

$$\hat{\mathbf{y}}_{T+t}^{BACE} = \sum_{j=1}^{L^K} \mathbf{P}_j \hat{\mathbf{y}}_{j,T+t} \quad (22)$$

where $\hat{\mathbf{y}}_{j,T+t}$ is the forecast at time $T + t$ of the j^{th} AVAR model, and \mathbf{P}_j is as discussed above. The corresponding standard error of $\hat{\mathbf{y}}_{T+t}^{BACE}$ is the average of the standard errors of the AVAR forecasts that is given by the formula

$$se(\hat{\mathbf{y}}_{T+t}^{BACE}) = \sum_{j=1}^{L^K} \mathbf{P}_j [se(\hat{\mathbf{y}}_{j,T+t})] \quad (23)$$

3.2 Summary of the Procedure

For a k -dimensional vector of stationary time-series variables y_i in \mathbf{y} , the procedure of the Bayesian Averaging of Classical Estimates in Asymmetric Vector Autoregressions (BACE-AVAR) is as follows:

1. Set L and \bar{l} , the maximum lag length for the AVAR models and the assumed symmetric lag length of the models. The \bar{l} can be set by using the automatic selection procedures for VAR models given a certain information criterion. L may be set as $L = \bar{l} + 3$, as adding a constant 3 to the lag length as specified by an automatic selection procedure is considered a rule of thumb among practitioners.

2. Estimate all possible AVAR models given L using OLS. The total number of models to be estimated is L^K .
3. For each of these models, compute the prior probability $P(\mathbf{M}_j)$ and the posterior probability $P(\mathbf{M}_j|y)$ that the j^{th} AVAR model has the correct specification.
4. For each model, obtain the forecasts along with their variances.
5. Compute the weighted forecasts and its corresponding variance at time t , \hat{y}_{T+t}^{BACE} and $se(\hat{y}_{T+t}^{BACE})$.

3.3 Performance of BACE-AVAR through Simulations

The performance of BACE-AVAR was assessed using simulations. A 3-dimensional vector time-series dataset⁹ was generated from a VAR or VARMA; for that particular dataset, all possible AVAR models were estimated given a maximum number of lags L , as well as their corresponding posterior probabilities of being the true model; the impulse responses and variable forecasts were weighted using these posterior model probabilities, and the results were compared with the true impulse response function and true forecasted values, respectively. The forecasting performance of the method was compared to the symmetric VAR that is selected by AIC for the cases with sample size 1000 and AICc for the other cases. The number of iterations for each case was set to 100.

⁹ A 4-dimensional vector time-series data will be generated in the cases where one important variable is omitted. See Section 3.3.1.4 for the discussion of Omitted Variable.

3.3.1 Specific Scenarios

The data were generated arbitrarily with the restriction that the time-series is stable and non-stationary. The BACE-AVAR was evaluated using cases summarized in Table 3.1. Overall, sixty scenarios were considered.

Table 3.1
Simulation Cases

Omitted Variable	MA Terms	u_t Covariance Matrix	Sample Size
1 With	1 With	1 Constant Variance 2	1 30 (1)
2 Without	2 Without	2 Constant Variance 3	2 50 (4)
		3 Constant Variance 5	3 100 (6)
			4 300 (7)
			5 1000 (8)

3.3.1.1 Sample Size

The range for the sample size is from 30 to 1000. The choice of range for the sample size is mainly due to the periodicity of the data that is being used in practice. A sample size of 30 can be viewed as an annual data; a sample size of 50, as quarterly data; a sample size of 100 as a monthly data; a sample size of 300 can be viewed as weekly data; and 1000 can be associated to daily data. Though these specifications were a generalization of common practice, the BACE-AVAR procedure does not limit the sample size with respect to the periodicity of the data. It is just more likely in practice that the sample size of the data is directly related to the period of the series, that is, higher sampling rate will have a larger sample size compared to lower sampling rates.

Furthermore, data collected on lower sampling rates have a higher amount of aggregated information than higher frequency samples. In mimicking this phenomenon, the AR part of the true model will have a lag order that is directly related to the sample size that will be obtained. The AR lag orders are enclosed in parenthesis beside the sample sizes in Table 3.1.

3.3.1.2 Covariance Matrix

Three cases will be set for the covariance matrix Σ of the innovation series $\mathbf{u}_t = (u_{xt}, u_{yt}, u_{zt})'$, or $\mathbf{u}_t = (u_{xt}, u_{yt}, u_{zt}, u_{ft})'$, in the case of an omitted variable. These will be limited to a diagonal matrix with equal elements, which were set to be two, three and five, that is, $\Sigma = i * I_K, i \in \{2, 3, 5\}$ and $K \in \{3, 4\}$. These values for the scalar covariance matrix were chosen to determine the performance of the BACE-AVAR procedure through different magnitudes of variances, relative to the performance of the automatically selected model. The starting values of the simulated data and the parameters for each replicate will be the same in order to have comparable results for the different specification of Σ .

3.3.1.3 Omitted Moving Average Terms

The case wherein there is an omitted MA term was also considered in the simulations. The MA term are restricted to lag order of one. The MA terms will be omitted in the modeling procedure.

3.3.1.4 Omitted Variable

It is likely in practice that some of the important variables in the system were not included in the modeling. It may be because that the variable is difficult to measure, the variable has not been measured, or the variable cannot be measured directly. In simulating this phenomenon, a 4-dimensional vector time-series will be generated, and only the three variables of interest

will be used in modeling. The AR and/or MA parameters of this omitted variable will be generated using the same procedure as the parameter generation of the variables of interest.

4. Results & Discussion

This chapter discusses the performance of the BACE-AVAR procedure in forecasting and in determining the interaction of the variables given some misspecification errors. The problems that were encountered in the simulation proper will be discussed first. The discussion of the results will then follow. In summary, the BACE-AVAR procedure has an advantage in forecasting over the automatically selected model under the problem of an omitted important variable and excluded MA term.

4.1 Preliminary Concerns on the Simulation

The formula for the posterior probability of the BACE procedure in cross-section data given Equation (17) involves the SSE being raised to the power $-n/2$, where n is the sample size of the cross-section data. In the BACE-AVAR procedure, the resulting posterior probability will be zero for a large sample size T due to the small size of the SSE . Therefore, the problem was counteracted by raising the SSE to $-(0.1T)/2$ on this part of the formula that is given in Equation (21) for the sample sizes $T = 300$ and $T = 1000$ since any power of the SSE that is less than $-100/2$ yields undefined posterior probabilities. However, this stands only as a temporary remedy to the problem. This issue posits that the order in which the SSE converges in the formula of the posterior probability may be different, in time-series data, from cross-section data, even if the model is estimated using OLS.

There were cases wherein the steps taken in order to have a simulated data that is stable and stationary do not work, since the generated parameters were selected at random. In order to guarantee a model that gives a stable data, a data burning of 10,000 time points were done.

Table 4.1 gives the lag length of the symmetric VAR models in the simulation. The number of AVAR models for models with lower lag length is small because $L = \bar{l} + 3$. This may affect the results of the averaging and may decrease the performance of the BACE-AVAR on smaller samples. Therefore, the performance for small samples of the BACE-AVAR procedure as stated in the results of the simulation may still be further improved.

Table 4.1
Average VAR Lag Lengths from Automatic Selection Procedure Using AICc

Sample Size	No Omitted Variable		With an Omitted Variable	
	No MA Term	With MA Term	No MA Term	With MA Term
30 (1)	1.03	1.36	1.07	1.33
50 (4)	3.02	3.46	2.44	2.86
100 (6)	5.66	6.27	4.85	5.4
300 (7)	7.04	8.93	7.27	8.64
1000* (8)	8.07	12.79	11.25	13.79

* AIC for Sample Size 1000

4.2 Forecasting Accuracy

This section will discuss the results of the forecasting accuracy of the BACE-AVAR procedure relative to the automatically selected model having the least value of AIC for large samples and least value of AICc for small samples, which will be hereafter called MINIC. In determining the forecasting accuracy of the procedures, the MDM test and the MAPE were used.

4.2.1 Modified Diebold-Mariano Test

The results for the forecasting accuracy are given in Table 4.2 and Table 4.3. The tables present the proportion of significant MDM test result at 10% level of significance that the BACE-AVAR procedure has a better measure of forecasting accuracy than the MINIC, and vice versa, for the different sample sizes, and by the loss functions that were used. The differences of the proportion of significant tests between the BACE-AVAR and the MINIC were also reported, as well as the average proportion of significant MDM tests for the variables of interest. The MDM test result for each of the variable of interest is given in Appendix A. Generally, the results of the MDM test indicate that the BACE-AVAR procedure is better than the MINIC in forecasting under the omitted variable problem. The BACE-AVAR procedure performs the same with respect to the MINIC across the different variance specification. This indicates that the BACE-AVAR procedure is not affected by the variance specification as specified in the simulations, relative to the automatically selected model.

In Table 4.2, for the case of no omitted variable and no omitted MA term, the BACE-AVAR is slightly better than the MINIC for small samples ($T = 30, 50$ and 100). However, its performance diminished for sample sizes where the posterior model probability was modified. For $T = 1000$, the forecasting accuracy of MINIC over the BACE-AVAR procedure is about 25%. For the case of no omitted variable but with an omitted MA term, the results indicate similar outcome as the previous case, but the improvement of the MINIC over the BACE-AVAR procedure now comes with smaller magnitude.

Table 4.2
MDM Test: Proportion of Significance for the Case of No Omitted Variable
 Alpha: 0.10; Number of Iterations: 100

Model	Sample Size	No MA Term		With MA Term	
		Absolute Loss	Squared Loss	Absolute Loss	Squared Loss

Significance of BACE over MINIC	30	14.3	15.7	14.0	18.7
	50	16.0	15.0	14.3	16.0
	100	10.7	11.3	12.7	15.0
	300	4.3	6.0	6.0	6.0
	1000	3.3	5.3	7.0	5.3
Significance of MINIC over BACE	30	10.0	13.3	13.3	13.0
	50	5.7	6.7	11.3	12.0
	100	10.3	12.3	6.3	8.0
	300	22.3	22.3	19.0	20.0
	1000	26.3	31.7	11.0	15.0
Difference (BACE- MINIC)	30	4.3	2.3	0.7	5.7
	50	10.3	8.3	3.0	4.0
	100	0.3	-1.0	6.3	7.0
	300	-18.0	-16.3	-13.0	-14.0
	1000	-23.0	-26.3	-4.0	-9.7

In Table 4.3, for the case of an omitted variable with no excluded MA term, the BACE-AVAR procedure has better forecasting accuracy than the MINIC except for $T = 30$. Furthermore, the magnitude of this improvement increases as the sample size increases. The improvement of the BACE-AVAR procedure drastically increases for sample sizes involving the modified posterior probability. For the case of omitting an important variable and MA term, the BACE-AVAR is better than the MINIC in terms of forecasting accuracy at around 15% to 20% depending on the sample size. The results emphasize that more information can still be extracted by the BACE-AVAR procedure for small samples by improving the weights or the posterior model probability.

Increasing the level of significance to 5% as stated in Tables A.2.1 to A.2.4 in Appendix A, yields the same interpretation that the MINIC is better than the BACE-AVAR procedure when there are no misspecification errors. In addition to this, the outputs also show that the result is reversed when there are misspecification errors. Increasing the level of significance to 1% as given in Tables A.3.1 to A.3.4 still has the same result, but the forecasting accuracy

of one procedure over the other may just be deemed as negligible due to the small values of the significant percentages. Full results are given in Appendix A.

Table 4.3
MDM Test: Proportion of Significance for the Case of an Omitted Variable
 Alpha: 0.10; Number of Iterations: 100

Model	Sample Size	No MA Term		With MA Term	
		Absolute Loss	Squared Loss	Absolute Loss	Squared Loss
Significance of BACE over MINIC	30	5.7	7.0	7.0	6.7
	50	10.7	10.3	11.0	11.7
	100	7.7	9.3	14.7	14.0
	300	19.3	17.7	20.7	23.0
	1000	16.3	20.7	15.0	18.0
Significance of MINIC over BACE	30	11.7	11.3	8.7	8.0
	50	5.0	6.0	5.0	8.7
	100	5.0	6.0	4.3	2.7
	300	2.3	2.0	2.3	3.0
	1000	2.0	2.0	2.7	1.3
Difference (BACE-MINIC)	30	-6.0	-4.3	-1.7	-1.3
	50	5.7	4.3	6.0	3.0
	100	2.7	3.3	10.3	11.3
	300	17.0	15.7	18.3	20.0
	1000	14.3	18.7	12.3	16.7

4.2.2 Mean Absolute Percentage Error

The forecasting accuracy was also measured descriptively by the relative MAPE. The relative MAPE forecasts of the BACE-AVAR procedure with respect to the MAPE forecasts of the MINIC are given in Table 4.4 and Appendix B. Relative MAPE values that are less than one imply that the BACE-AVAR forecasts are closer to the outsample data than the MINIC, whereas values greater than one indicate the opposite.

Table 4.4
Average Relative MAPE of Forecasts

Sample Size	No Omitted Variable		With Omitted Variable	
	No MA	With MA	No MA	With MA

30	0.6195	0.6390	1.0007	0.7803
50	0.7221	1.3715	0.9521	1.1204
100	1.1456	0.7054	1.2131	1.3044
300	0.5876	0.6308	0.4813	0.6042
1000	0.7547	1.5094	0.5429	0.7231

The result in Table 4.4 for the case of no omitted variable and no omitted MA term reveals that the forecasts of the BACE-AVAR procedure using the unmodified posterior probability are closer to the actual values compared to the forecasts of the MINIC by about 35% except for the sample size 50. For the case of $T = 300$ and 1000, the BACE-AVAR procedure also has the same performance over the MINIC. This indicates that even if the MDM test suggests that the MINIC is superior to the BACE-AVAR for cases of no misspecification, the forecasts of BACE-AVAR are closer to the actual values than the forecasts of the MINIC on the average.

For the case of no omitted variable but with an omitted MA term, it seems that the forecasts of the BACE-AVAR procedure is closer to the outsample data than that of the MINIC by about 35% except for $T = 50$. The result still holds for the cases of the modified posterior probability except for $T = 1000$. Thus, the BACE-AVAR procedure is, on the average, still at par or better in some cases than the MINIC based on the relative MAPE.

The BACE-AVAR procedure is better than or at par to the MINIC for the case wherein an important variable is omitted. The relative MAPE of BACE-AVAR to the MAPE of the MINIC reaches around 0.50 for the sample size of 1000 – a 50% improvement over the MINIC. All the relative MAPE of the sample sizes that were subjected to the modified posterior probability are less than one. This result is not contained only for the sample size of 300 and 1000 but is also evident for $T = 300$, except for $T = 100$, that exhibits a relative MAPE of 1.21. This may give an indication that the power of the SSE_j in Equation (21) may

not be a linear function of the sample size. Nevertheless, it is evident that the BACE-AVAR procedure can reach a 50% improvement over the MINIC in terms of MAPE forecasts.

For the case of an omitted variable and excluded MA term, the BACE-AVAR procedure improves over the MINIC for $T = 30, 300$ and 1000 . The relative MAPE for the sample size 50 and 100 is 1.12 and 1.30, respectively. Furthermore, the improvement of the MINIC over the BACE-AVAR procedure exhibits an upward trend from the sample size of 30 to 100, which are the sample sizes that use the original form of the posterior model probability. This may imply that correction on the posterior probability may also be applied to improve the overall performance of the BACE-AVAR procedure. Full results are given in Appendix B.

5. Conclusion

Misspecification problems in VAR modeling such as incorrect AR lag, excluded MA terms, and omitted relevant variables, are common in practice. The worst problem among those that were stated is omitting an important variable since it is immeasurable given the data on hand. Aside from that, it also gives biased and inconsistent estimates. The implication of this problem is crucial in policy evaluation since it will yield misleading forecasts and incorrect variable relationships.

This study presents the application of the BACE on AVAR models in forecasting in presence of misspecification errors. Simulations under different scenarios were done to evaluate the performance of the BACE-AVAR procedure over an automatically selected model using an information criterion.

The results suggest that the BACE-AVAR procedure produces more accurate forecasts than the automatically selected model using the corrected AIC when there is a problem of omitted

variables and omitted MA term. The forecasting accuracy of the BACE-AVAR procedure is better for large sample sizes. On the other hand, if there are no omitted variable and excluded MA term, the automatically selected model is better than the BACE-AVAR procedure. Given the results of the study, the BACE-AVAR procedure is recommended in forecasting.

It is recommended to simulate the performance of the BACE-AVAR procedure for processes generated from a structural VAR. It is also recommended to extend the BACE-AVAR procedure for cointegrated variables. But before working on these recommendations, it is suggested to improve the posterior model probability in order for the BACE-AVAR procedure to be at par with the automatically selected model when there is no misspecification error.

Bibliography

- Akaike, H. (1973). A New Look at the Statistical Model Identification. *IEEE Transactions on Automatic Control*, Vol. 19(No. 6), pp. 716-723.
- Akaike, H. (1974, Dec.). A new look at the statistical model identification. *Automatic Control, IEEE Transactions on*, Vol. 19(No. 6), pp. 716,723.
- Braun, P. A., & Mittnik, S. (1993, Oct.). Misspecifications in vector autoregressions and their effects on impulse responses and variance decompositions. *Journal of Econometrics*, Vol. 59(No. 3), pp. 319-241.
- Carriero, A., Kapetanios, G., & Marcellino, M. (2009). Forecasting Exchange Rates with a Large Bayesian VAR. *International Journal of Forecasting*, pp. 400-417.
- Chen, A.-S., & Leung, M. T. (2003). A Bayesian Vector Error Correction Model in Forecasting Exchange Rates. *Computers and Operations Research*, Vol. 30, pp. 887-900.
- Diebold, F. X., & Roberto, M. S. (1995, July). Comparing Predictive Accuracy. *Journal of Business & Economic Statistics*, Vol. 13(No. 3), pp. 253-263.
- Fackler, J. S., & Krieger, S. C. (1986, Jan). An Application of Vector Time Series Techniques to Macroeconomic Forecasting. *Journal of Business & Economic Statistics*, Vol. 4(No. 1), pp. 71-80.
- Hannan, E. J., & Quinn, B. G. (1979). The Determination of the Order of an Autoregression. *Journal of the Royal Statistical Society. Series B (Methodological)*, Vol. 41(No. 2), pp. 190-195.
- Harvey, D., Leybourne, S., & Newbold, P. (1997). Testing the Equality of Prediction Mean Squared Errors. *International Journal of Forecasting*, Vol. 13, pp. 281-291.
- Hsiao, C. (1981). Autoregressive Modelling and Money-Income Causality Detection. *Journal of Monetary Economics*, Vol. 7(No. 1), pp 85-106.
- Hurvich, C. M., & Tsai, C.-L. (1993). A Corrected Akaike Information Criterion for Vector Autoregressive Model Selection. *Journal of Time Series Analysis*, Vol. 14(No. 3), pp. 271-279.
- Jorda, O. (2005, March). Estimation and Inference of Impulse Responses by Local Projections. *The American Economic Review*, Vol. 95(No. 1), pp. 161-182.
- Kadilar, C., & Erdemir, C. (2002). Comparison of Performance Among Information Criteria in VAR and Seasonal VAR models. *Hacettepe Journal of Mathematics and Statistics*, Vol. 31, pp. 127-137.
- Keating, J. W. (1993). Asymmetric Vector Autoregression. *Proceedings of the Business and Economic Statistics Section* (pp. pp. 68-73). American Statistical Association.
- Keating, J. W. (1995). Vector Autoregressive Models with Asymmetric Lag Structure. *Working Paper*.

- Keating, J. W. (2000). Macroeconomic Modeling with Asymmetric Vector Autoregressions. *Journal of Macroeconomics*, Vol. 22(No. 1), pp. 1-28.
- Korobilis, D. (2010, March 4). *VAR forecasting using Bayesian variable selection*. Retrieved January 24, 2011, from Munich Personal RePec Archive: <http://mpra.ub.uni-muenchen.de/21124/>
- Kullback, S., & Leibler, R. A. (1951). On Information and Sufficiency. *Annals of Mathematical Statistics*, Vol. 22(No. 1), pp. 79-86.
- Leamer, E. E. (1983, March). Let's Take the Con Out of Econometrics. *The American Economic Review*, Vol. 73(No. 1), pp. 31-43.
- Litterman, R. B. (1980). A Bayesian Procedure for Forecasting With Vector Autoregressions. *Working Paper*.
- Lütkepohl, H. (1990, Feb.). Asymptotic Distributions of Impulse Response Functions and Forecast Error Variance Decompositions of Vector Autoregressive Models. *The Review of Economics and Statistics*, Vol. 72(No. 1), pp. 116-125.
- Lütkepohl, H. (2004). *Forecasting with VARMA Models*. Department of Economics, European University Institute.
- Ng, S., & Pierre, P. (2001, Nov.). Lag Length Selection and the Construction of Unit Root Tests with Good Size and Power. *Econometrica*, Vol. 69(No. 6), pp. 1519-1554.
- Ozcicek, O., & McMillin, W. D. (1999, April). Lag length selection in vector autoregressive models: symmetric and asymmetric lags. *Applied Economics*, Vol. 31(No. 4), pp. 517-524.
- Po, H. H., Chi, H. W., Shyu, J., & Hsiao, C. Y. (2002). A Litterman BVAR approach for production forecasting of technological industries. *Technological Forecasting and Social Change*, Vol. 70, pp. 67-82.
- Ramos, F. F. (2003). Forecasts of market shares from VAR and BVAR models: a comparison of their accuracy. *International Journal of Forecasting*, Vol. 19, pp. 95-100.
- Runkle, D. E. (1987, Oct.). Vector Autoregressions and Reality. *Journal of Business and Economic Statistics*, Vol. 5(No. 4), pp. 437-42.
- Sala-I-Martin, X. X. (1997, May). I Just Ran Two Million Regressions. *The American Economic Review*, Vol. 87(No. 2), pp. 178-183.
- Sala-I-Martin, X., Doppelhofer, G., & Miller, R. (2004, September). Determinants of Long-Term Growth: A Bayesian Averaging of Classical Estimates (BACE). *The American Economic Review*, Vol. 94(No. 4), pp. 813-835.
- Schwartz, G. (1978). Estimating the Dimension of a Model. *Annals of Statistics*, Vol. 6(No. 2), pp. 461-464.
- Schwarz, G. (1978). Estimating the Dimension of a Model. *Annals of Statistics*, Vol. 6(No. 2), pp. 461-464.

- Seghouane, A.-K. (2006, Oct.). Vector Autoregressive Model-Order Selection From Finite Samples Using Kullback's Symmetric Divergence. *Circuits and Systems I: Regular Papers, IEEE Transactions on*, Vol. 53(No. 10), pp. 2327-2335.
- Shibata, R. (1980). Asymptotically efficient selection of the order of the model for estimating parameters in a linear process. *Annals of Statistics*, Vol. 8(No. 1), pp. 147-146.
- Sims, C. A. (1980). Comparison of Interwar and Postwar Business Cycles: Monetarism Reconsidered. *The American Economic Review: Papers and Proceedings of the Ninety-Second Annual Meeting of the American Economic Association*. Vol. 70, No. 2, pp. 250-257. American Economic Association.
- Sims, C. A. (1980, Jan). Macroeconomics and Reality. *Econometrica*, Vol. 48(No. 1), pp. 1-48.
- Sims, C. A., & Zha, T. (1999, Sep.). Error Bands for Impulse Responses. *Econometrica*, Vol. 67(No. 5), pp. 1113-1155.
- Stock, J. H., & Watson, M. W. (1996, Jan.). Evidence on Structural Instability in Macroeconomic Time Series Relations. *Journal of Business & Economic Statistics*, Vol. 14(No. 1), pp. 11-30.
- Stock, J. H., & Watson, M. W. (2001). Vector Autoregressions. *The Journal of Economic Perspective*, Vol. 15(No. 4), pp. 105-115.
- Strachan, R., & van Dijk, H. K. (2007). Bayesian Model Averaging in Vector Autoregressive Processes with an Investigation of Stability of the US Great Ratios and Risk of a Liquidity Trap in the USA, UK and Japan. *MRG Discussion Paper Series 1407*.
- Sun, D., & Ni, S. (2004). Bayesian analysis of vector-autoregressive models with noninformative priors. *Journal of Statistical Planning and Inference*, Vol. 121, pp. 291-309.
- Tsay, R. S. (2005). *Analysis of Financial Time Series* (Second Edition ed.). Hoboken, New Jersey: John Wiley & Sons, Inc.
- Waele, S. d., & Broersen, P. M. (2002, Oct.). Finite sample effects in vector autoregressive modelling. *Instrumentation and Measurement, IEEE Transactions on*, Vol. 51(No. 5), pp. 917-992.
- Waele, S. d., & Broersen, P. M. (2003, Feb.). Order Selection for Vector Autoregressive Models. *Signal Processing, IEEE Transactions on*, Vol. 51(No. 2), pp. 427-433.

APPENDIX A

MDM Test: Proportion of Significance

Table A.1.1

Alpha: 0.10; Number of Iterations: 100
Case: No Omitted Variable; No MA Terms

Model	Sample Size	Absolute Loss Function				Squared Loss Function			
		Average (%)	V1	V2	V3	Average (%)	V1	V2	V3
Significance of BACE over MINIC	30	0.15	0.13	0.15	15.7	0.14	0.13	0.20	0.15
	50	0.18	0.15	0.15	15.0	0.16	0.15	0.14	0.18
	100	0.11	0.13	0.08	11.3	0.10	0.16	0.08	0.11
	300	0.06	0.05	0.02	6.0	0.08	0.07	0.03	0.06
	1000	0.01	0.03	0.06	5.3	0.05	0.05	0.06	0.01
Significance of MINIC over BACE	30	0.12	0.10	0.08	13.3	0.14	0.16	0.10	0.12
	50	0.08	0.04	0.05	6.7	0.10	0.06	0.04	0.08
	100	0.07	0.12	0.12	12.3	0.09	0.15	0.13	0.07
	300	0.23	0.22	0.22	22.3	0.21	0.27	0.19	0.23
	1000	0.28	0.26	0.25	31.7	0.33	0.32	0.30	0.28
Difference (BACE-MINIC)	30	0.03	0.03	0.07	2.3	0.00	-0.03	0.10	0.03
	50	0.10	0.11	0.10	8.3	0.06	0.09	0.10	0.10
	100	0.04	0.01	-0.04	-1.0	0.01	0.01	-0.05	0.04
	300	-0.17	-0.17	-0.20	-16.3	-0.13	-0.20	-0.16	-0.17
	1000	-0.27	-0.23	-0.19	-26.3	-0.28	-0.27	-0.24	-0.27

Table A.1.2

Alpha: 0.10; Number of Iterations: 100
Case: No Omitted Variable; With MA Terms

Model	Sample Size	Absolute Loss Function				Squared Loss Function			
		Average (%)	V1	V2	V3	Average (%)	V1	V2	V3
Significance of BACE over MINIC	30	14.0	0.10	0.14	0.18	18.7	0.12	0.24	0.20
	50	14.3	0.11	0.18	0.14	16.0	0.14	0.18	0.16
	100	12.7	0.12	0.16	0.10	15.0	0.15	0.19	0.11
	300	6.0	0.07	0.06	0.05	6.0	0.07	0.05	0.06
	1000	7.0	0.08	0.07	0.06	5.3	0.07	0.04	0.05
Significance of MINIC over BACE	30	13.3	0.18	0.12	0.10	13.0	0.19	0.10	0.10
	50	11.3	0.12	0.11	0.11	12.0	0.10	0.14	0.12
	100	6.3	0.08	0.05	0.06	8.0	0.11	0.07	0.06
	300	19.0	0.18	0.23	0.16	20.0	0.17	0.22	0.21
	1000	11.0	0.10	0.08	0.15	15.0	0.18	0.13	0.14
Difference (BACE-MINIC)	30	0.7	-0.08	0.02	0.08	5.7	-0.07	0.14	0.10
	50	3.0	-0.01	0.07	0.03	4.0	0.04	0.04	0.04
	100	6.3	0.04	0.11	0.04	7.0	0.04	0.12	0.05
	300	-13.0	-0.11	-0.17	-0.11	-14.0	-0.10	-0.17	-0.15
	1000	-4.0	-0.02	-0.01	-0.09	-9.7	-0.11	-0.09	-0.09

Table A.1.3

Alpha: 0.10; Number of Iterations: 100

Case: With Omitted Variable; No MA Terms

Model	Sample Size	Absolute Loss Function				Squared Loss Function			
		Average (%)	V1	V2	V3	Average (%)	V1	V2	V3
Significance of BACE over MINIC	30	5.7	0.09	0.03	0.05	7.0	0.11	0.06	0.04
	50	10.7	0.07	0.10	0.15	10.3	0.10	0.11	0.10
	100	7.7	0.08	0.09	0.06	9.3	0.08	0.11	0.09
	300	19.3	0.14	0.22	0.22	17.7	0.15	0.20	0.18
	1000	16.3	0.17	0.15	0.17	20.7	0.23	0.21	0.18
Significance of MINIC over BACE	30	11.7	0.11	0.09	0.15	11.3	0.09	0.10	0.15
	50	5.0	0.09	0.04	0.02	6.0	0.09	0.06	0.03
	100	5.0	0.07	0.04	0.04	6.0	0.10	0.05	0.03
	300	2.3	0.04	0.01	0.02	2.0	0.03	0.01	0.02
	1000	2.0	0.03	0.01	0.02	2.0	0.02	0.02	0.02
Difference (BACE-MINIC)	30	-6.0	-0.02	-0.06	-0.10	-4.3	0.02	-0.04	-0.11
	50	5.7	-0.02	0.06	0.13	4.3	0.01	0.05	0.07
	100	2.7	0.01	0.05	0.02	3.3	-0.02	0.06	0.06
	300	17.0	0.10	0.21	0.20	15.7	0.12	0.19	0.16
	1000	14.3	0.14	0.14	0.15	18.7	0.21	0.19	0.16

Table A.1.4

Alpha: 0.10; Number of Iterations: 100

Case: With Omitted Variable; With MA Terms

Model	Sample Size	Absolute Loss Function				Squared Loss Function			
		Average (%)	V1	V2	V3	Average (%)	V1	V2	V3
Significance of BACE over MINIC	30	7.0	0.08	0.03	0.10	6.7	0.07	0.05	0.08
	50	11.0	0.12	0.07	0.14	11.7	0.13	0.09	0.13
	100	14.7	0.15	0.20	0.09	14.0	0.12	0.20	0.10
	300	20.7	0.18	0.23	0.21	23.0	0.21	0.26	0.22
	1000	15.0	0.17	0.15	0.13	18.0	0.19	0.18	0.17
Significance of MINIC over BACE	30	8.7	0.07	0.10	0.09	8.0	0.08	0.08	0.08
	50	5.0	0.04	0.09	0.02	8.7	0.09	0.12	0.05
	100	4.3	0.02	0.04	0.07	2.7	0.02	0.04	0.02
	300	2.3	0.04	0.01	0.02	3.0	0.04	0.02	0.03
	1000	2.7	0.03	0.04	0.01	1.3	0.02	0.01	0.01
Difference (BACE-MINIC)	30	-1.7	0.01	-0.07	0.01	-1.3	-0.01	-0.03	0.00
	50	6.0	0.08	-0.02	0.12	3.0	0.04	-0.03	0.08
	100	10.3	0.13	0.16	0.02	11.3	0.10	0.16	0.08
	300	18.3	0.14	0.22	0.19	20.0	0.17	0.24	0.19
	1000	12.3	0.14	0.11	0.12	16.7	0.17	0.17	0.16

Table A.2.1

Alpha: 0.05; Number of Iterations: 100

Case: No Omitted Variable; No MA Terms

Model	Sample Size	Absolute Loss Function				Squared Loss Function			
		Average (%)	V1	V2	V3	Average (%)	V1	V2	V3
Significance of BACE over MINIC	30	10.0	0.11	0.08	0.11	11.0	0.11	0.10	0.12
	50	9.3	0.10	0.10	0.08	8.7	0.08	0.09	0.09
	100	6.3	0.06	0.09	0.04	7.3	0.08	0.11	0.03
	300	2.0	0.04	0.02	0.00	4.0	0.06	0.04	0.02
	1000	1.0	0.00	0.00	0.03	2.0	0.02	0.01	0.03
Significance of MINIC over BACE	30	8.0	0.08	0.08	0.08	7.7	0.09	0.08	0.06
	50	2.7	0.04	0.02	0.02	3.7	0.04	0.04	0.03
	100	5.0	0.04	0.07	0.04	6.7	0.05	0.08	0.07
	300	16.3	0.17	0.17	0.15	16.0	0.19	0.14	0.15
	1000	21.0	0.23	0.19	0.21	21.0	0.19	0.22	0.22
Difference (BACE-MINIC)	30	2.0	0.03	0.00	0.03	3.3	0.02	0.02	0.06
	50	6.7	0.06	0.08	0.06	5.0	0.04	0.05	0.06
	100	1.3	0.02	0.02	0.00	0.7	0.03	0.03	-0.04
	300	-14.3	-0.13	-0.15	-0.15	-12.0	-0.13	-0.10	-0.13
	1000	-20.0	-0.23	-0.19	-0.18	-19.0	-0.17	-0.21	-0.19

Table A.2.2

Alpha: 0.05; Number of Iterations: 100

Case: No Omitted Variable; With MA Terms

Model	Sample Size	Absolute Loss Function				Squared Loss Function			
		Average (%)	V1	V2	V3	Average (%)	V1	V2	V3
Significance of BACE over MINIC	30	9.7	0.07	0.08	0.14	12.0	0.05	0.18	0.13
	50	9.0	0.07	0.09	0.11	10.3	0.10	0.08	0.13
	100	9.7	0.10	0.13	0.06	9.0	0.10	0.12	0.05
	300	3.0	0.03	0.04	0.02	3.7	0.04	0.01	0.06
	1000	3.7	0.05	0.02	0.04	3.0	0.05	0.02	0.02
Significance of MINIC over BACE	30	9.0	0.12	0.07	0.08	9.7	0.13	0.07	0.09
	50	7.7	0.07	0.09	0.07	7.7	0.07	0.09	0.07
	100	4.3	0.08	0.02	0.03	3.3	0.07	0.02	0.01
	300	12.0	0.11	0.15	0.10	12.3	0.14	0.13	0.10
	1000	8.0	0.08	0.08	0.08	7.7	0.09	0.07	0.07
Difference (BACE-MINIC)	30	0.7	-0.05	0.01	0.06	2.3	-0.08	0.11	0.04
	50	1.3	0.00	0.00	0.04	2.7	0.03	-0.01	0.06
	100	5.3	0.02	0.11	0.03	5.7	0.03	0.10	0.04
	300	-9.0	-0.08	-0.11	-0.08	-8.7	-0.10	-0.12	-0.04
	1000	-4.3	-0.03	-0.06	-0.04	-4.7	-0.04	-0.05	-0.05

Table A.2.3

Alpha: 0.05; Number of Iterations: 100

Case: With Omitted Variable; No MA Terms

Model	Sample Size	Absolute Loss Function				Squared Loss Function			
		Average (%)	V1	V2	V3	Average (%)	V1	V2	V3
Significance of BACE over MINIC	30	3.0	0.07	0.01	0.01	3.7	0.09	0.01	0.01
	50	8.0	0.05	0.09	0.10	6.3	0.05	0.08	0.06
	100	3.7	0.02	0.05	0.04	5.3	0.06	0.04	0.06
	300	9.3	0.08	0.11	0.09	9.3	0.07	0.12	0.09
	1000	11.0	0.14	0.10	0.09	9.3	0.11	0.09	0.08
Significance of MINIC over BACE	30	7.7	0.07	0.07	0.09	6.0	0.05	0.07	0.06
	50	2.3	0.04	0.03	0.00	3.0	0.05	0.02	0.02
	100	3.3	0.06	0.01	0.03	3.0	0.04	0.02	0.03
	300	1.7	0.02	0.01	0.02	2.0	0.03	0.01	0.02
	1000	1.0	0.02	0.00	0.01	0.7	0.00	0.00	0.02
Difference (BACE-MINIC)	30	-4.7	0.00	-0.06	-0.08	-2.3	0.04	-0.06	-0.05
	50	5.7	0.01	0.06	0.10	3.3	0.00	0.06	0.04
	100	0.3	-0.04	0.04	0.01	2.3	0.02	0.02	0.03
	300	7.7	0.06	0.10	0.07	7.3	0.04	0.11	0.07
	1000	10.0	0.12	0.10	0.08	8.7	0.11	0.09	0.06

Table A.2.4

Alpha: 0.05; Number of Iterations: 100

Case: With Omitted Variable; With MA Terms

Model	Sample Size	Absolute Loss Function				Squared Loss Function			
		Average (%)	V1	V2	V3	Average (%)	V1	V2	V3
Significance of BACE over MINIC	30	5.0	0.05	0.03	0.07	4.0	0.05	0.02	0.05
	50	7.3	0.08	0.05	0.09	7.7	0.10	0.04	0.09
	100	7.7	0.09	0.09	0.05	10.0	0.11	0.14	0.05
	300	12.3	0.11	0.13	0.13	12.0	0.07	0.14	0.15
	1000	9.0	0.13	0.07	0.07	12.3	0.13	0.12	0.12
Significance of MINIC over BACE	30	4.7	0.04	0.05	0.05	5.3	0.05	0.05	0.06
	50	2.7	0.02	0.05	0.01	4.0	0.04	0.05	0.03
	100	2.0	0.00	0.03	0.03	1.0	0.00	0.03	0.00
	300	1.3	0.03	0.01	0.00	1.7	0.03	0.01	0.01
	1000	0.0	0.00	0.00	0.00	0.7	0.01	0.00	0.01
Difference (BACE-MINIC)	30	0.3	0.01	-0.02	0.02	-1.3	0.00	-0.03	-0.01
	50	4.7	0.06	0.00	0.08	3.7	0.06	-0.01	0.06
	100	5.7	0.09	0.06	0.02	9.0	0.11	0.11	0.05
	300	11.0	0.08	0.12	0.13	10.3	0.04	0.13	0.14
	1000	9.0	0.13	0.07	0.07	11.7	0.12	0.12	0.11

Table A.3.1

Alpha: 0.01; Number of Iterations: 100
Case: No Omitted Variable; No MA Terms

Model	Sample Size	Absolute Loss Function				Squared Loss Function			
		Average (%)	V1	V2	V3	Average (%)	V1	V2	V3
Significance of BACE over MINIC	30	4.7	0.06	0.03	0.05	5.7	0.06	0.04	0.07
	50	4.0	0.06	0.03	0.03	4.0	0.06	0.03	0.03
	100	1.3	0.01	0.02	0.01	2.0	0.02	0.03	0.01
	300	0.7	0.02	0.00	0.00	1.0	0.02	0.01	0.00
	1000	0.0	0.00	0.00	0.00	0.0	0.00	0.00	0.00
Significance of MINIC over BACE	30	4.7	0.03	0.05	0.06	4.3	0.04	0.03	0.06
	50	1.0	0.01	0.01	0.01	1.0	0.02	0.01	0.00
	100	2.7	0.04	0.02	0.02	2.0	0.04	0.02	0.00
	300	8.0	0.10	0.07	0.07	7.0	0.08	0.07	0.06
	1000	7.3	0.10	0.04	0.08	5.0	0.06	0.04	0.05
Difference (BACE-MINIC)	30	0.0	0.03	-0.02	-0.01	1.3	0.02	0.01	0.01
	50	3.0	0.05	0.02	0.02	3.0	0.04	0.02	0.03
	100	-1.3	-0.03	0.00	-0.01	0.0	-0.02	0.01	0.01
	300	-7.3	-0.08	-0.07	-0.07	-6.0	-0.06	-0.06	-0.06
	1000	-7.3	-0.10	-0.04	-0.08	-5.0	-0.06	-0.04	-0.05

Table A.3.2

Alpha: 0.01; Number of Iterations: 100
Case: No Omitted Variable; With MA Terms

Model	Sample Size	Absolute Loss Function				Squared Loss Function			
		Average (%)	V1	V2	V3	Average (%)	V1	V2	V3
Significance of BACE over MINIC	30	4.7	0.02	0.06	0.06	6.0	0.04	0.07	0.07
	50	4.7	0.05	0.04	0.05	4.0	0.03	0.03	0.06
	100	2.3	0.02	0.04	0.01	2.3	0.04	0.02	0.01
	300	1.0	0.01	0.02	0.00	1.7	0.01	0.01	0.03
	1000	1.0	0.01	0.00	0.02	1.3	0.03	0.00	0.01
Significance of MINIC over BACE	30	5.7	0.07	0.04	0.06	6.0	0.08	0.04	0.06
	50	5.7	0.05	0.07	0.05	4.3	0.04	0.06	0.03
	100	0.7	0.02	0.00	0.00	1.0	0.01	0.01	0.01
	300	4.3	0.07	0.05	0.01	4.3	0.06	0.05	0.02
	1000	2.3	0.00	0.03	0.04	2.3	0.03	0.02	0.02
Difference (BACE-MINIC)	30	-1.0	-0.05	0.02	0.00	0.0	-0.04	0.03	0.01
	50	-1.0	0.00	-0.03	0.00	-0.3	-0.01	-0.03	0.03
	100	1.7	0.00	0.04	0.01	1.3	0.03	0.01	0.00
	300	-3.3	-0.06	-0.03	-0.01	-2.7	-0.05	-0.04	0.01
	1000	-1.3	0.01	-0.03	-0.02	-1.0	0.00	-0.02	-0.01

Table A.3.3

Alpha: 0.01; Number of Iterations: 100

Case: With Omitted Variable; No MA Terms

Model	Sample Size	Absolute Loss Function				Squared Loss Function			
		Average (%)	V1	V2	V3	Average (%)	V1	V2	V3
Significance of BACE over MINIC	30	1.0	0.02	0.00	0.01	0.7	0.01	0.00	0.01
	50	2.7	0.02	0.03	0.03	2.3	0.02	0.03	0.02
	100	1.3	0.01	0.01	0.02	0.7	0.00	0.01	0.01
	300	3.7	0.01	0.06	0.04	1.7	0.02	0.03	0.00
	1000	3.0	0.04	0.03	0.02	3.0	0.05	0.03	0.01
Significance of MINIC over BACE	30	2.3	0.01	0.02	0.04	3.0	0.02	0.04	0.03
	50	0.0	0.00	0.00	0.00	0.0	0.00	0.00	0.00
	100	1.7	0.03	0.00	0.02	1.7	0.03	0.01	0.01
	300	0.7	0.01	0.01	0.00	0.0	0.00	0.00	0.00
	1000	0.0	0.00	0.00	0.00	0.0	0.00	0.00	0.00
Difference (BACE-MINIC)	30	-1.3	0.01	-0.02	-0.03	-2.3	-0.01	-0.04	-0.02
	50	2.7	0.02	0.03	0.03	2.3	0.02	0.03	0.02
	100	-0.3	-0.02	0.01	0.00	-1.0	-0.03	0.00	0.00
	300	3.0	0.00	0.05	0.04	1.7	0.02	0.03	0.00
	1000	3.0	0.04	0.03	0.02	3.0	0.05	0.03	0.01

Table A.3.4

Alpha: 0.01; Number of Iterations: 100

Case: With Omitted Variable; With MA Terms

Model	Sample Size	Absolute Loss Function				Squared Loss Function			
		Average (%)	V1	V2	V3	Average (%)	V1	V2	V3
Significance of BACE over MINIC	30	1.3	0.01	0.01	0.02	1.0	0.01	0.01	0.01
	50	3.0	0.04	0.03	0.02	3.3	0.03	0.03	0.04
	100	2.3	0.03	0.03	0.01	3.0	0.04	0.04	0.01
	300	3.3	0.02	0.04	0.04	3.0	0.03	0.04	0.02
	1000	3.0	0.06	0.02	0.01	2.3	0.05	0.02	0.00
Significance of MINIC over BACE	30	1.3	0.00	0.01	0.03	2.0	0.02	0.02	0.02
	50	1.0	0.01	0.02	0.00	1.7	0.00	0.04	0.01
	100	0.3	0.00	0.01	0.00	0.3	0.00	0.01	0.00
	300	1.0	0.03	0.00	0.00	0.0	0.00	0.00	0.00
	1000	0.0	0.00	0.00	0.00	0.0	0.00	0.00	0.00
Difference (BACE-MINIC)	30	0.0	0.01	0.00	-0.01	-1.0	-0.01	-0.01	-0.01
	50	2.0	0.03	0.01	0.02	1.7	0.03	-0.01	0.03
	100	2.0	0.03	0.02	0.01	2.7	0.04	0.03	0.01
	300	2.3	-0.01	0.04	0.04	3.0	0.03	0.04	0.02
	1000	3.0	0.06	0.02	0.01	2.3	0.05	0.02	0.00

APPENDIX B
Relative MAPE of Forecasts
MAPE(BACE-AVAR)/MAPE(MINIC)

Table B.1

Case: No Omitted Variable; No MA Term

n	V1	V2	V3	Average
30	0.4684	0.6977	0.6924	0.6195
50	0.6220	0.7342	0.8100	0.7221
100	0.6338	1.4573	1.3457	1.1456
300	0.6588	0.5710	0.5329	0.5876
1000	0.6678	0.7614	0.8349	0.7547

Table B.2

Case: No Omitted Variable; With MA Term

n	V1	V2	V3	Average
30	0.6718	0.6824	0.5628	0.6390
50	2.1923	1.0767	0.8456	1.3715
100	0.2722	0.7403	1.1038	0.7054
300	0.3998	0.8795	0.6133	0.6308
1000	2.6022	0.9538	0.9723	1.5094

Table B.3

Case: With Omitted Variable; No MA Term

n	V1	V2	V3	Average
30	0.7150	0.8818	1.4053	1.0007
50	0.6256	0.9643	1.2664	0.9521
100	1.2097	1.3674	1.0622	1.2131
300	0.4643	0.4511	0.5285	0.4813
1000	0.6053	0.5543	0.4691	0.5429

Table B.4

Case: With Omitted Variable; With MA Term

n	V1	V2	V3	Average
30	0.5468	0.5716	1.2224	0.7803
50	1.2342	1.1004	1.0267	1.1204
100	1.4864	1.2833	1.1436	1.3044
300	1.1347	0.3448	0.3331	0.6042
1000	0.6644	0.8525	0.6523	0.7231