



Munich Personal RePEc Archive

Choosing the variables to estimate singular DSGE models: Comment

Iskrev, Nikolay and Ritto, Joao

August 2016

Online at <https://mpa.ub.uni-muenchen.de/72870/>
MPRA Paper No. 72870, posted 06 Aug 2016 07:55 UTC

Choosing the variables to estimate singular DSGE models: Comment

Nikolay Iskrev* João Ritto†

Abstract

In a recent article Canova et al. (2014) study the optimal choice of variables to use in the estimation of a simplified version of the Smets and Wouters (2007) model. In this comment we examine their conclusions by applying a different methodology to the same model. Our results call into question most of Canova et al. (2014) conclusions.

Keywords: DSGE models, Observables, Identification, Information matrix, Cramér-Rao lower bounds

JEL classification: C32, C51, C52, E32

*Banco de Portugal, Av. Almirante Reis 71, 1150-012, Lisboa, Portugal. E-mail: niskrev@bportugal.pt.

†Banco de Portugal, Av. Almirante Reis 71, 1150-012, Lisboa, Portugal. E-mail: jpritto@bportugal.pt.

1 Introduction

DSGE models are usually estimated using only a subset of the variables that are present in them. This is partly due to the fact that some variables, such as capital, are not observed. However, even variables for which data exist are often not utilized. One could explain this with the restriction that the number of included observables should not be greater than the number of shocks in the model. While there are ways to get around this restriction,¹ it is a fact that in the literature there are similar DSGE models estimated with different sets of observables. It is not clear what motivates these different choices, nor what consequences that has on the empirical findings.

In a recent article Canova, Ferroni, and Matthes (2014) (CFM henceforth) seek to provide some guidance on how to select the most informative among several available sets of observables. They propose the use of two criteria which rank different combinations of variables according to measures of identification and information content. The first criterion starts by selecting the sets of variables that satisfy a rank condition for identification of the free model parameters. To pick the best among the selected sets, measures of closeness to a convoluted singular system of all observables are computed in terms of sensitivity of the log-likelihood function to parameters of interest. The one yielding smallest discrepancy is chosen as the most informative. The second criterion is based on Bierens (2007) and uses convolutions of both the singular and non-singular systems with the same non-singular distribution. The combination of variables whose convoluted distribution is closest to the convoluted singular system of all available observables is selected as being the most informative.

CFM apply their selection criteria to a simplified version of the Smets and Wouters (2007) model. The model has 4 shocks and a total of 7 observables, namely output (y_t), consumption (c_t), investment (i_t), wages (w_t), hours (h_t), inflation (π_t), and nominal interest rate (r_t). Thus, 35 combinations of variables are available to use in estimation. Among these, as most informative overall the authors select y_t , c_t , i_t and either w_t or h_t . Furthermore, it is argued that the ranking of different sets of variables does not depend on the value of the parameters at which the model is evaluated, and is robust to increasing the number of shocks as in the original Smets and Wouters (2007) model.

The purpose of this comment is to evaluate these claims, applying a different analytical approach to the same model. As in Iskrev (2010), where the choice of observables is studied with respect to the original Smets and Wouters (2007) model, here we use criteria based on the expected Fisher information matrix (FIM). Using the FIM has several advantages. First, as the name suggests, it is a measure of the amount of information about the parameters available in a sample (see Rothenberg (1971)). It takes the model as it is and does not require convoluting the true data density as the measures CFM use do.² Second, FIM depends on the set of observables and the sample size, but does not depend on actual data. Thus, the information one could expect to have in different sets of observables and in samples of different sizes can be measured and compared prior to estimation. Third, using the FIM one can compute measures of expected estimation uncertainty with respect to each model parameter. In general, there is a trade-off between the amount of information contained in different sets

¹One is to introduce measurement errors in the observed series. Another is the approach in Bierens (2007).

²This approach follows Bierens (2007) where the theoretical model is assumed to be misspecified and the singular distribution it implies is convoluted and compared to a convoluted distribution of an a-theoretical econometric model which is assumed to represent the true data generating process.

of observables with respect to different parameters. Quantifying the amount of information for each parameter provides a clearer understanding of the trade-offs involved in selecting one set of observables over another. The measures CFM use do not provide such information. And fourth, the FIM can be evaluated analytically for linearized Gaussian model such as the one in Smets and Wouters (2007). This is very useful in practice since it allows many possible combinations of variables to be compared quickly for a large number of *a priori* plausible parameter values. Furthermore, the use of analytical derivatives minimizes the risk of reaching wrong conclusions as a result of numerical errors.

2 Analysis

In this section we apply the FIM approach to the model analyzed in CFM. We address three main questions: (1) is the rank condition useful for selecting the set of observables, (2) which is the most informative set of four variables out of the seven variables that are available, and (3) are the results sensitive to changes in the parameter values and the number of shocks.

2.1 Is the rank condition useful?

We start by checking whether the parameters of the simplified SW model are identified if only four of the seven variables are observed. It is well known that four parameters - ξ_w , ξ_p , ϵ_w and ϵ_p , are not separately identifiable in the sense that in the linearized model ξ_w cannot be distinguished from ϵ_w , and ξ_p cannot be distinguished from ϵ_p . As in the original paper, we will assume that ϵ_w and ϵ_p are both known. This leaves 27 free parameters.

A necessary and sufficient condition for local identification is that the FIM has full rank. When evaluated at the parameter values from Table 2 in CFM, the FIM has full rank of 27 for all 35 combinations of four variables. Thus, the rank condition alone provides no useful information regarding the best set of variables to use in estimating the model.

2.2 Which are the best four observables?

Selecting the best combination of variables requires a criterion on the basis of which to compare and rank the alternatives. Which criterion should be used depends on the purpose for which the model is estimated. In any case, the criterion would be a function of the estimated parameters and would rank as better sets of observables that are more informative about the relevant function of the parameters of interest θ .

When the objective is to minimize the estimation uncertainty about θ as a whole, a popular criterion to use is the natural logarithm of the determinant of the inverse of the FIM, i.e. $\ln(\det(\mathcal{I}^{-1}(\theta)))$. This is known in the optimal design literature as D-optimality criterion. The well-known Cramér-Rao (CR) theorem tells us that, depending on whether the asymptotic FIM is used or the finite sample one, its inverse gives either a lower bound on the asymptotic covariance matrix of any consistent estimator of θ , or a lower bound on the covariance matrix of any unbiased estimator θ . Furthermore, the diagonal elements of $\mathcal{I}^{-1}(\theta)$ are lower bounds on the variances of estimators of individual parameters. This can be used to construct a criterion which assigns different weights to the parameters, so as to reflect their

Table 1: Most informative and least informative sets of observables

rank	geometric average			D-optimality
	24 parameters	17 parameters	6 parameters	24 parameters
1	y, c, w, r	y, c, w, π	c, i, π, r	c, i, π, r
2	c, i, π, r	y, c, w, r	y, c, w, π	c, w, π, r
3	y, c, w, h	c, i, π, r	y, c, π, h	y, w, π, h
33	c, i, w, π	y, i, π, r	y, i, w, h	y, i, π, r
34	y, c, i, h	y, c, i, π	y, i, r, h	y, i, r, h
35	y, c, i, π	y, c, i, h	y, c, i, h	y, c, i, h

Note: The table shows the best 3 and the worst 3 sets of observables according to the geometric average and D-optimality criteria. The geometric average criterion is computed for 3 groups of parameters: 24 (all free) parameters; 17 (all except shock) parameters; 6 (only $\lambda, \iota_p, \xi_p, \sigma_l, r_\pi, r_y$) parameters.

relative importance to the researcher. An example of such a criterion is the weighted geometric average of the diagonal elements of $\mathcal{I}^{-1}(\theta)$,

$$\text{geometric average criterion} = \left(\prod_{i=1}^k \text{CRLB}_{\theta_i}^{w_i} \right)^{1/\sum_{i=1}^k w_i} \quad (2.1)$$

where CRLB_{θ_i} is the i -th diagonal element of $\mathcal{I}^{-1}(\theta)$, k is the number of free parameters, and w_i is the weight assigned to θ_i . The geometric average is more appropriate to use than the arithmetic average since parameters typically have different range.

In what follows we use the finite sample FIM in order to take a proper account of the size of the sample, which is set to $T=150$, as in CFM.³ We report three versions of the weighted geometric average criterion with: (1) equal weights on all free parameters; (2) equal weights on the free structural parameters and zero weights on the shock parameters; (3) equal weights on the six parameters emphasized in CFM, namely $\lambda, \iota_p, \xi_p, \sigma_l, r_\pi$, and r_y , and zero weights on all other parameters. To be comparable with CFM, we assume that δ, λ_w and c_g are known. This leaves 24 free parameters, 17 of which are structural and the other 7 are shock parameters.

Table 1 lists the best three and worst three sets of variables according to each criterion. The set containing (c, i, π, r) is selected as most informative by two of the criteria, while the other two rank it among the top three sets. All criteria select sets containing (y, c, i) as least informative, with three of the criteria picking h , and the fourth one selecting π as the worst fourth variable. However, as can be seen in the first quadrant of Figure 1, the difference between (y, c, i, h) and (y, c, i, π) , is very small, when the criterion is the geometric average of all 24 parameter. The figure shows the values associated with the 35 sets of variables, sorted from best to worst according to each criterion. It can be seen that (c, i, π, r) is in fact very close to the optimal sets selected by the first two criteria, which rank it second

³The asymptotic FIM is defined as the limit of the average finite sample FIM, which in turn is the negative expected Hessian of the log-likelihood function. With the asymptotic FIM information accumulates at a constant rate T and therefore the ranking of observables does not depend on the sample size. With the final sample FIM information may accumulate at different and changing rates for different sets of observables. Thus, the ranking may change with the sample size.

and third, respectively. It also shows that there are numerically meaningful differences between the most and least informative sets of variables.




Figure 1: Sorted values of different ranking criteria.

Table 2 reports the values of the individual CRLBs for the most and the least informative sets of variables, as per the results in table 1. In addition, the set (y, c, i, w) is also included as it was selected by CFM as one of the two most informative combinations. According to the criteria we use, this is the most informative combination of variables that includes simultaneously y , c , and i . It is ranked 9-th when the criterion is the geometric average of the CRLBs of the 17 structural parameters. As can be seen from the table, choosing one combination of variables over another usually involves a trade-off in terms of information about different parameters. Even the least informative set (y, c, i, h) is the most informative one, amongst those in the table, for three of the free parameters, ρ_{ga} , φ , and σ_a . The overall best set (c, i, π, r) , yields the lowest (among the six in the table) CRLBs for a half of the free parameters, including three of the six deep parameters CFM focus on. If these are the parameters we are most interested in, the only reason to select (y, c, i, w) over (c, i, π, r) would be if one assigns much larger weights on σ_l and ξ_p than on the other four parameters. In particular, there is much less information about the Taylor rule parameters, due to the absence of both r and π in that set. As can be seen from the last row in panel B, with equal weights (c, i, π, r) is more than twice as informative any of the sets that include y , c , and i .

One of the criteria used by CFM ranks the sets of variables on the basis of the sensitivity of the likelihood to a group of parameters of interest. The measures they use compare the scores of the non-singular and convoluted singular systems, and require simulated data to compute. A simpler and more direct measure of sensitivity to a single parameter θ_i is the expected curvature of log-likelihood function, given by $-\text{E}\left(\frac{\partial^2 \ell_T(\boldsymbol{\theta})}{\partial \theta_i^2}\right)$. Note that this is just the i -th diagonal element of the FIM and does not require data to compute. When there is only one free parameter the sensitivity (also called Fisher information) is equal to the inverse of the CRLB. Hence, in the single parameter case, the most sensitive and most informative combinations of variables coincide. When there are multiple free parameters, however, this

Table 2: Individual and overall parameter uncertainty

param.	(c, i, π, r)	(y, c, w, r)	(y, c, w, π)	(y, c, i, w)	(y, c, i, π)	(y, c, i, h)
	A. CRLBs of individual parameters					
ρ_{ga}	1.754	0.206	0.215	0.245	0.307	0.158
α	0.048	0.049	0.046	0.054	0.075	0.071
ψ	0.102	0.201	0.255	0.131	0.187	0.179
β	0.011	0.016	0.018	0.011	0.015	0.014
φ	2.812	3.852	4.874	6.153	6.770	2.566
σ_c	0.154	0.192	0.279	0.224	0.536	0.430
λ	0.031	0.027	0.035	0.044	0.099	0.151
Φ	0.323	0.123	0.124	0.159	1.482	0.221
ι_w	0.180	0.269	0.072	0.243	1.061	1.359
ξ_w	0.142	0.021	0.018	0.021	0.177	0.257
ι_p	0.044	0.247	0.074	0.208	0.099	1.187
ξ_p	0.155	0.044	0.043	0.048	0.401	0.480
σ_l	0.790	0.163	0.196	0.223	2.000	0.934
r_π	0.300	1.588	1.336	4.476	1.753	3.973
$r_{\Delta y}$	0.048	0.042	0.138	0.166	0.193	0.569
r_y	0.060	0.296	0.236	0.632	0.257	1.057
ρ	0.028	0.050	0.053	0.082	0.091	0.189
ρ_a	0.015	0.018	0.022	0.025	0.029	0.023
ρ_g	0.004	0.011	0.014	0.014	0.015	0.016
ρ_I	0.049	0.064	0.074	0.095	0.100	0.066
σ_a	0.418	0.100	0.098	0.208	0.361	0.046
σ_g	0.145	0.044	0.048	0.043	0.089	0.054
σ_I	0.063	0.197	0.271	0.082	0.086	0.082
σ_τ	0.016	0.017	0.132	0.126	0.255	0.292
	B. Overall (geometric average of CRLBs)					
24 parameters	0.094	0.092	0.103	0.126	0.218	0.217
17 parameters	0.133	0.129	0.126	0.170	0.321	0.384
6 parameters	0.120	0.169	0.138	0.256	0.391	0.834
	C. Overall (D-optimality criterion)					
24 parameters	-183	-158	-148	-139	-114	-108

Note: Panel A shows the values of the Cramér-Rao lower bounds (CRLBs) for sample size $T = 150$. Panel B shows the geometric averages of the bounds for three groups of parameters: 24 (all free) parameters; 17 (all except shock) parameters; 6 (only $\lambda, \iota_p, \xi_p, \sigma_l, r_\pi, r_y$). Panel C shows the values of $\ln(\det(\mathcal{I}^{-1}))$. Lower values always indicate more information.

Table 3: Most sensitive and most informative sets of variables

param.	most sensitive		most informative	
	variables	CRLBs	variables	CRLBs
ρ_{ga}	(y, c, i, h)	0.158	(y, c, i, h)	0.158
α	(c, i, π, r)	0.048	(c, i, r, h)	0.039
ψ	(c, w, π, r)	0.284	(c, i, π, r)	0.102
β	(c, i, π, r)	0.011	(c, i, r, h)	0.009
φ	(c, w, π, h)	2.679	(i, w, r, h)	1.793
σ_c	(c, i, π, r)	0.154	(c, w, r, h)	0.131
λ	(c, w, π, r)	0.028	(y, c, i, r)	0.022
Φ	(y, w, π, h)	0.039	(y, w, π, h)	0.039
ι_w	(y, w, π, h)	0.106	(c, w, π, r)	0.065
ξ_w	(y, w, π, h)	0.052	(y, c, w, h)	0.014
ι_p	(y, w, π, h)	0.042	(y, π, r, h)	0.036
ξ_p	(y, w, π, h)	0.016	(y, w, π, h)	0.016
σ_l	(y, w, π, h)	1.305	(y, c, w, r)	0.163
r_π	(c, w, π, r)	1.282	(c, i, π, r)	0.300
$r_{\Delta y}$	(c, i, π, r)	0.048	(y, c, w, r)	0.042
r_y	(c, w, π, r)	0.217	(c, i, π, r)	0.060
ρ	(c, w, π, r)	0.032	(c, π, r, h)	0.026
ρ_a	(c, w, π, r)	0.019	(c, i, w, r)	0.009
ρ_g	(c, w, π, r)	0.012	(c, i, π, r)	0.004
ρ_I	(c, w, π, r)	0.091	(c, w, π, h)	0.041
σ_a	(y, w, π, h)	0.030	(y, w, π, h)	0.030
σ_g	(y, i, w, r)	0.054	(y, c, i, w)	0.043
σ_I	(c, i, π, r)	0.063	(c, i, π, r)	0.063
σ_r	(c, i, π, r)	0.016	(c, i, π, r)	0.016

Note: The most sensitive set of variables w.r.t. θ_i is the one maximizing the i -th diagonal element of \mathcal{I} . The most informative set is the one minimizing the i -th diagonal element of \mathcal{I}^{-1} .

is not necessarily true. As can be seen in Table 3, the most sensitive and most informative sets coincide only for 6 of the 24 parameters. The table also shows the CRLBs corresponding to each set of variables. In several cases the differences are very large, meaning that the most sensitive selection contains much less information than the most informative one. A case in point is σ_l for which the CRLB with the most sensitive combination (y, w, π, h) is 8 times larger than with the most informative combination (y, c, w, r) .

As explained in greater details in Iskrev (2010), the values of the CRLBs are determined by the interactions of two factors – the sensitivity of the log-likelihood function to changes in individual parameters, and the degree of collinearity among the effects of such changes. A large value of the CRLB indicates that a parameter has only a weak effect on the log-likelihood function, and/or that its effect on the log-likelihood can to a large extent be offset by the effects of other parameters. In the case of σ_l , it is much harder to distinguish its effect on the log-likelihood from the effects of parameters like ι_w , ξ_w and ι_p , when the observables are (y, w, π, h) , compared to when the observables are (y, c, w, r) .

Table 4: Most informative and least informative sets of observables, different parameterizations

rank	geometric average			D-optimality
	24 parameters	17 parameters	6 parameters	24 parameters
A. CFM model, prior mean				
1	c, w, r, h	c, w, r, h	c, w, r, h	c, i, r, h
2	c, i, r, h	c, i, r, h	c, i, r, h	c, w, r, h
3	y, c, w, π	y, c, w, π	c, π, r, h	y, c, i, w
33	y, c, i, h	y, i, π, r	c, i, w, h	y, c, i, r
34	y, c, i, r	y, c, i, h	y, i, w, h	y, i, r, h
35	y, i, π, r	y, c, i, r	y, c, i, h	y, c, i, h
B. SW model, prior mean				
1	y, i, h, r	y, i, h, r	c, w, h, π	y, i, h, r
2	y, c, h, r	y, c, h, r	c, h, π, r	y, c, h, r
3	y, i, h, π	i, h, π, r	y, c, h, π	y, c, i, r
33	y, c, i, w	y, c, i, h	y, i, w, h	c, w, h, π
34	c, i, w, h	y, c, i, w	y, c, i, w	c, i, w, π
35	c, i, w, π	c, i, w, π	y, c, i, h	c, w, π, r
C. SW model, posterior mean				
1	y, i, h, r	y, i, π, r	c, h, π, r	y, i, h, r
2	y, i, π, r	i, w, h, r	c, i, π, r	y, h, π, r
3	y, c, h, r	y, i, h, r	w, h, π, r	y, c, h, r
33	y, w, h, π	c, i, w, π	y, c, i, r	y, c, i, w
34	y, c, i, h	y, c, i, π	y, c, i, w	y, w, h, π
35	c, i, w, π	y, c, i, h	y, c, i, h	y, c, i, h

Note: see note to Table 1.

2.3 Are the results robust to changes in the parameter values and the number of shocks?

The results presented in the last section are conditional on the particular parameter values and the assumptions CFM make regarding the number of shocks and the stationarity of the observables. Here we check whether the optimal selection of observables is robust to changes in the parameter values and the model specification.

We consider three alternatives. First, we change the parameter values keeping the rest of the model as before. Instead of the baseline parametrization, which is similar to the posterior mean in Smets and Wouters (2007), we use the prior mean from that paper. The prior mean is a natural choice since the analysis is supposed to happen prior to estimation. In the other two cases the model is as specified in Smets and Wouters (2007), i.e. with seven shocks and deterministic trend, and the growing variables being observed in terms of growth rates. We refer to this as the SW specification. That model is also evaluated at the means of the prior and the posterior distributions. In order for the results to be comparable with those in the previous section, we maintain the same group of free parameters. Therefore, we assume that the parameters of the three additional shocks as well as the trend parameter

Table 5: Optimal sets of observables, different parameterizations

param.	A. CFM model				B. SW model			
	prior mean		posterior mean		prior mean		posterior mean	
	variables	CRLBs	variables	CRLBs	variables	CRLBs	variables	CRLBs
ρ_{ga}	(y, i, w, h)	0.110	(y, c, i, h)	0.158	(y, c, i, h)	0.119	(y, c, i, h)	0.136
α	(y, c, i, w)	0.005	(c, i, r, h)	0.039	(c, h, π, r)	0.025	(c, h, π, r)	0.167
ψ	(y, c, i, w)	0.051	(c, i, π, r)	0.102	(c, i, h, r)	0.030	(y, c, i, r)	0.031
β	(y, c, i, h)	0.003	(c, i, r, h)	0.009	(y, i, w, h)	0.148	(y, c, i, w)	0.251
φ	(c, i, π, h)	0.076	(i, w, r, h)	1.793	(c, i, h, r)	0.667	(c, i, h, r)	2.544
σ_c	(c, i, r, h)	0.039	(c, w, r, h)	0.131	(c, i, h, r)	0.202	(c, i, h, r)	0.253
λ	(c, i, r, h)	0.004	(y, c, i, r)	0.022	(c, i, h, r)	0.047	(c, i, h, r)	0.079
Φ	(y, w, π, h)	0.060	(y, w, π, h)	0.039	(c, i, w, h)	0.083	(y, i, h, r)	0.195
ι_w	(c, i, w, π)	0.083	(c, w, π, r)	0.065	(c, w, h, π)	0.122	(c, w, h, π)	0.173
ξ_w	(y, w, π, r)	0.015	(y, c, w, h)	0.014	(c, w, h, π)	0.066	(c, w, h, r)	0.055
ι_p	(c, w, π, h)	0.044	(y, π, r, h)	0.036	(y, w, h, π)	0.073	(w, h, π, r)	0.093
ξ_p	(y, w, π, h)	0.033	(y, w, π, h)	0.016	(y, w, h, π)	0.091	(y, w, h, π)	0.067
σ_l	(c, w, r, h)	0.039	(y, c, w, r)	0.163	(c, w, h, π)	1.164	(c, w, h, r)	1.555
r_π	(c, i, r, h)	0.167	(c, i, π, r)	0.300	(y, h, π, r)	0.552	(c, i, π, r)	0.510
$r_{\Delta y}$	(c, w, r, h)	0.024	(y, c, w, r)	0.042	(y, i, π, r)	0.043	(c, i, h, r)	0.072
r_y	(c, i, r, h)	0.073	(c, i, π, r)	0.060	(c, h, π, r)	0.072	(c, i, π, r)	0.057
ρ	(c, π, r, h)	0.021	(c, π, r, h)	0.026	(y, h, π, r)	0.068	(c, h, π, r)	0.047
ρ_a	(c, i, r, h)	0.038	(c, i, w, r)	0.009	(y, i, w, h)	0.073	(y, i, h, r)	0.025
ρ_g	(y, c, i, w)	0.052	(c, i, π, r)	0.004	(y, c, i, h)	0.076	(y, c, h, r)	0.017
ρ_I	(c, i, r, h)	0.044	(c, w, π, h)	0.041	(c, i, h, r)	0.082	(c, i, h, r)	0.075
σ_a	(y, i, w, h)	0.007	(y, w, π, h)	0.030	(y, i, w, h)	0.009	(y, i, w, h)	0.042
σ_g	(y, w, π, r)	0.008	(y, c, i, w)	0.043	(y, c, i, h)	0.008	(y, c, i, h)	0.050
σ_I	(c, i, r, h)	0.008	(c, i, π, r)	0.063	(c, i, h, r)	0.010	(c, i, w, r)	0.057
σ_r	(c, π, r, h)	0.006	(c, i, π, r)	0.016	(c, h, π, r)	0.008	(c, h, π, r)	0.021

Note: see note to Table 3

are known.

Table 4 shows a summary of the results using the same criteria as before. Clearly, while there is considerable consistency in the ranking across different criteria, the optimal combination of variables is not invariant to the parametrization. Also, the two sets, (y, c, i, h) and (y, c, i, w) , recommended by CFM, are consistently ranked among the least informative, especially when the focus is on the six deep parameters.

The optimal combinations of variables for each parameter are shown in Table 5. Panel A reports the results for the CFM specification of the model evaluated at the prior and posterior mean values. In panel B are shown the results for the SW specification. In the case of the CFM model, the optimal combinations of variables are the same for only 3 parameters. For the SW specification there are 11 parameters for which the optimal sets coincide. There is only one parameter for which the most informative combination is the same across both parameterizations and model specifications. In very few cases the optimal combination of variables includes simultaneously y , c , and i .

Table 6: Information gains, posterior mean of the SW model

param.	y	c	i	w	h	π	r
ρ_{ga}	92	32	40	2	72	1	9
β	4	29	15	2	11	46	88
α	39	39	81	3	12	10	63
ψ	41	21	43	18	25	5	22
φ	4	30	54	6	22	5	28
σ_c	8	48	30	5	16	22	44
λ	13	39	29	8	25	10	45
Φ	26	16	33	9	77	6	15
ι_w	2	3	3	84	2	55	5
ξ_w	7	31	12	34	16	14	22
ι_p	4	3	3	38	7	60	6
ξ_p	30	9	21	58	65	36	9
σ_l	11	37	14	19	43	16	26
r_π	3	14	10	4	15	54	45
$r_{\Delta y}$	15	22	21	4	16	20	71
r_y	4	12	8	5	17	50	43
ρ	3	13	8	4	15	53	56
ρ_a	32	38	37	8	39	3	31
ρ_g	43	57	36	4	21	3	23
ρ_I	2	14	63	3	8	5	13
σ_a	88	9	35	3	81	2	7
σ_g	90	53	56	3	37	1	11
σ_I	1	8	86	2	4	5	13
σ_r	3	11	1	1	17	38	89

Note: The efficiency gain $EG_{\theta_i}(x_j)$ measures the reduction in uncertainty about parameter θ_i due to observing variable x_j , expressed as a per cent of the parameter uncertainty when x_j is not observed.

2.4 The role of interest rate and inflation

One of the main conclusions reached by CFM is that neither interest rate nor inflation data should be selected, if one had to choose only four of the seven variables. This is surprising since some of the parameters CFM focus on are the price stickiness and price indexation parameters as well as the inflation and output coefficients in the monetary policy rule. Intuitively, one would expect that inflation and interest rate are very informative about these parameters.

To formally measure the amount of information contributed by each one of the observed variables, we compute parameter efficiency gains defined as the expected reduction in parameter uncertainty due to observing a variable, expressed as a percent of the uncertainty when that variable is not observed. Formally, the efficiency gain of a variable x_j with respect to a parameter θ_i is defined as

$$\text{EG}_{\theta_i}(x_j) = 100 \left(\frac{\text{CRLB}_{\theta_i}(\mathbf{x} \setminus x_j) - \text{CRLB}_{\theta_i}(\mathbf{x})}{\text{CRLB}_{\theta_i}(\mathbf{x} \setminus x_j)} \right) \quad (2.2)$$

where \mathbf{x} is the set of all variables: $\mathbf{x} := \{y, c, i, h, w, \pi, r\}$.

Since we want to know how much information each variable contributes relative to all other variables, we consider the full SW model, evaluated at the posterior mean value of $\boldsymbol{\theta}$. Table 6 shows the efficiency gains with respect to all free parameters. For 8 of them the largest efficiency gains come from either r or π . This includes 4 of the 6 parameters CFM focus of, namely ι_p , r_π , r_y , for which π is the most informative variable, and λ , for which r is the most informative variable. As can be seen from the table, π and r are also very informative about several other parameters, e.g. α , σ_c , ι_w , ξ_p , and σ_l , suggesting that excluding these variables would lead to a substantial loss of information.

2.5 Monte Carlo study

The FIM-based analysis is a simple way of quantifying the information content of the restrictions the DSGE model imposes on the joint probability distribution of the observed variables. This makes it well suited for ranking different sets of observables in terms of the amount of information about the unknown parameters one could *expect* to get from each set. In this section we evaluate the predictions of the FIM approach using Monte Carlo simulations. In particular, we simulate the baseline model with 4 structural shocks to generate 400 artificial samples of 150 observations for each of the seven observable variables. We estimate by maximum likelihood the 24 free parameters using different subsets of four variables. We focus on the six subsets presented in Table 2, which comprise of the most informative ones according to our FIM-based criteria, and the subsets recommended by CFM.

Table 7 reports the simulated root mean squared errors (RMSE) for individual parameters as well as the average RMSEs of the three groups of parameters considered before. As before, we regard the sets of observables yielding lower average RMSE as being relatively more informative with respect to a given group of parameters. Overall, the results confirm the ranking of observables implied by the FIM analysis. In particular, the three subsets containing y, c and i have significantly larger average RMSEs than the other three subsets. As before (c, i, π, r) is ranked as the most informative set of variables with respect to the six structural parameter CFM focus on. In fact, the ranking of variables according to that criterion is exactly the same as the one in Table 2. Also, it is worth noting that

Table 7: Simulated root mean squared errors, $T = 150$

param.	(c, i, π, r)	(y, c, w, r)	(y, c, w, π)	(y, c, i, w)	(y, c, i, π)	(y, c, i, h)
	A. individual parameters					
ρ_{ga}	0.478	0.216	0.219	0.229	0.275	0.173
α	0.050	0.053	0.060	0.050	0.060	0.055
ψ	0.110	0.204	0.251	0.148	0.195	0.187
β	0.011	0.016	0.020	0.011	0.015	0.012
φ	3.120	3.321	3.230	3.849	4.222	3.256
σ_c	0.135	0.214	0.359	0.306	0.405	0.447
λ	0.028	0.034	0.059	0.069	0.098	0.097
Φ	0.389	0.144	0.181	0.209	0.690	0.267
ι_w	0.244	0.299	0.086	0.295	0.418	0.428
ξ_w	0.072	0.029	0.021	0.033	0.221	0.250
ι_p	0.050	0.242	0.078	0.236	0.128	0.450
ξ_p	0.112	0.058	0.058	0.073	0.205	0.283
σ_l	1.186	0.288	0.315	0.347	3.529	2.226
r_π	0.394	0.803	0.816	0.914	0.735	0.824
$r_{\Delta y}$	0.053	0.046	0.151	0.218	0.152	0.395
r_y	0.073	0.224	0.230	0.245	0.177	0.161
ρ	0.025	0.044	0.054	0.117	0.083	0.121
ρ_a	0.008	0.017	0.051	0.052	0.062	0.040
ρ_g	0.005	0.012	0.059	0.026	0.031	0.030
ρ_I	0.049	0.060	0.066	0.070	0.082	0.070
σ_a	0.148	0.096	0.144	0.258	0.241	0.057
σ_g	0.164	0.047	0.064	0.047	0.077	0.056
σ_I	0.058	0.268	0.301	0.114	0.130	0.100
σ_r	0.017	0.018	0.131	0.145	0.182	0.181
	B. Overall					
24 parameters	0.085	0.097	0.125	0.137	0.191	0.176
17 parameters	0.112	0.128	0.142	0.162	0.252	0.237
6 parameters	0.132	0.171	0.158	0.212	0.325	0.392

Note: The root mean squared errors (RMSE) are calculated using 400 Monte Carlo simulations. Panel B shows the geometric averages of the RMSE for three groups of parameters: 24 (all free) parameters; 17 (all except shock) parameters; 6 (only λ , ι_p , ξ_p , σ_l , r_π , r_y).

Table 8: Monte Carlo results and theoretical CRLBs (part I)

param.	True	(c, i, π, r)			(y, c, w, r)			(y, c, w, π)		
		Mean	Std. dev.	CRLB	Mean	Std. dev.	CRLB	Mean	Std. dev.	CRLB
ρ_{ga}	0.510	0.459	0.476	1.754	0.503	0.216	0.206	0.452	0.211	0.215
α	0.20	0.210	0.049	0.048	0.204	0.053	0.049	0.216	0.058	0.046
ψ	0.26	0.258	0.110	0.102	0.272	0.203	0.201	0.277	0.250	0.255
β	0.99	0.988	0.010	0.011	0.983	0.014	0.016	0.980	0.017	0.018
φ	5.48	6.033	3.074	2.812	5.113	3.305	3.852	4.601	3.111	4.874
σ_c	1.39	1.408	0.134	0.154	1.477	0.196	0.192	1.524	0.334	0.279
λ	0.71	0.705	0.028	0.031	0.696	0.031	0.027	0.687	0.054	0.035
Φ	1.61	1.681	0.383	0.323	1.652	0.138	0.123	1.530	0.162	0.124
ι_w	0.59	0.592	0.244	0.180	0.698	0.279	0.269	0.560	0.080	0.072
ξ_w	0.73	0.734	0.072	0.142	0.716	0.026	0.021	0.729	0.021	0.018
ι_p	0.47	0.456	0.048	0.044	0.506	0.239	0.247	0.463	0.078	0.074
ξ_p	0.65	0.615	0.107	0.155	0.640	0.057	0.044	0.659	0.057	0.043
σ_l	1.92	2.235	1.145	0.790	1.969	0.285	0.163	1.933	0.315	0.196
r_π	2.03	2.129	0.382	0.300	1.907	0.794	1.588	2.155	0.808	1.336
$r_{\Delta y}$	0.22	0.239	0.049	0.048	0.213	0.046	0.042	0.263	0.144	0.138
r_y	0.08	0.092	0.072	0.060	0.211	0.182	0.296	0.218	0.184	0.236
ρ	0.87	0.871	0.025	0.028	0.874	0.044	0.050	0.883	0.052	0.053
ρ_a	0.95	0.947	0.007	0.015	0.948	0.017	0.018	0.923	0.043	0.022
ρ_g	0.97	0.968	0.004	0.004	0.968	0.012	0.011	0.949	0.055	0.014
ρ_I	0.71	0.708	0.049	0.049	0.712	0.060	0.064	0.720	0.066	0.074
σ_a	0.46	0.421	0.143	0.418	0.442	0.095	0.100	0.504	0.137	0.098
σ_g	0.61	0.619	0.164	0.145	0.605	0.046	0.044	0.598	0.063	0.048
σ_I	0.60	0.609	0.058	0.063	0.758	0.216	0.197	0.810	0.216	0.271
σ_r	0.25	0.254	0.017	0.016	0.249	0.018	0.017	0.276	0.129	0.132

Note: The means and standard deviations of MLE are calculated using 400 Monte Carlo simulations.

the values of the RMSE are generally very similar to the respective values of the CRLBs. This is not something we would necessarily expect for at least two reasons. First, the CRLBs are by definition lower bounds on the standard deviations of unbiased estimators. Hence, even if the estimation bias is small, the actual RMSEs may be significantly larger than the theoretical lower bounds. Second, the CRLBs do not account for any a priori restrictions on the parameter values, such as the restriction that a parameter has to be between 0 and 1. In our ML estimation we imposed such restrictions on a number of parameters, e.g. $\beta, \alpha, \lambda, \xi_w, \xi_p, \iota_w, \iota_p$, as well as the autoregressive coefficients of the shocks.⁴ One consequence of ignoring these restrictions could be that the theoretical bounds on the estimation uncertainty are larger than the actual uncertainty. Such discrepancies occurred in a very few cases in our simulations, as can be seen in Tables 8 and 9 (see for instance the values for ρ_{ga} when (c, i, π, r) is observed). In the vast majority of cases the theoretical bounds are very close to the simulation-based standard errors.

⁴We also impose the restriction that the model has a unique solution, which further restricts the parameter space.

Table 9: Monte Carlo results and theoretical CRLBs (part II)

		(y, c, i, w)			(y, c, i, π)			(y, c, i, h)		
param.	True	Mean	Std. dev.	CRLB	Mean	Std. dev.	CRLB	Mean	Std. dev.	CRLB
ρ_{ga}	0.51	0.436	0.217	0.245	0.629	0.248	0.307	0.488	0.172	0.158
α	0.20	0.210	0.049	0.054	0.206	0.060	0.075	0.209	0.055	0.071
ψ	0.26	0.278	0.147	0.131	0.282	0.194	0.187	0.294	0.184	0.179
β	0.99	0.988	0.010	0.011	0.987	0.015	0.015	0.988	0.012	0.014
φ	5.48	5.887	3.832	6.153	5.809	4.214	6.770	6.019	3.215	2.566
σ_c	1.39	1.443	0.302	0.224	1.435	0.403	0.536	1.529	0.425	0.430
λ	0.71	0.684	0.064	0.044	0.675	0.091	0.099	0.668	0.087	0.151
Φ	1.61	1.562	0.203	0.159	1.497	0.681	1.482	1.636	0.266	0.221
ι_w	0.59	0.726	0.262	0.243	0.691	0.406	1.061	0.638	0.426	1.359
ξ_w	0.73	0.718	0.031	0.021	0.722	0.221	0.177	0.691	0.247	0.257
ι_p	0.47	0.548	0.223	0.208	0.487	0.127	0.099	0.609	0.429	1.187
ξ_p	0.65	0.672	0.069	0.048	0.698	0.200	0.401	0.522	0.252	0.480
σ_l	1.92	1.889	0.346	0.223	3.903	2.924	2.000	2.920	1.991	0.934
r_π	2.03	1.850	0.897	4.476	1.993	0.735	1.753	1.995	0.824	3.973
$r_{\Delta y}$	0.22	0.242	0.217	0.166	0.120	0.150	0.193	0.394	0.355	0.569
r_y	0.08	0.228	0.195	0.632	0.137	0.167	0.257	0.126	0.154	1.057
ρ	0.87	0.838	0.113	0.082	0.851	0.081	0.091	0.822	0.111	0.189
ρ_a	0.95	0.927	0.046	0.025	0.925	0.057	0.029	0.930	0.034	0.023
ρ_g	0.97	0.961	0.024	0.014	0.958	0.028	0.015	0.958	0.028	0.016
ρ_I	0.71	0.701	0.069	0.095	0.718	0.082	0.100	0.705	0.070	0.066
σ_a	0.46	0.580	0.229	0.208	0.479	0.241	0.361	0.454	0.057	0.046
σ_g	0.61	0.597	0.045	0.043	0.565	0.062	0.089	0.591	0.053	0.054
σ_I	0.60	0.654	0.101	0.082	0.653	0.119	0.086	0.635	0.094	0.082
σ_r	0.25	0.260	0.145	0.126	0.274	0.181	0.255	0.294	0.176	0.292

Note: see the note to table 8

3 Concluding remarks

Our results can be summarized as follows: (1) The rank condition for identification is not informative about the optimal choice of observables for the model that was analyzed. In general, such a criterion could be useful when only one, or a very few, of the many possible sets of variables satisfy the identification condition. This seems to rarely be the case in practice. (2) At the baseline parametrization of the model the most informative set of variables includes consumption, investment, interest rate, and inflation. (3) The most informative set of variables is not invariant to the parametrization of the model or the number of shocks.

All of these findings call into question the conclusions of CFM. Perhaps the most important one is (3), which implies that the optimal selection of observables cannot, in general, be based on analysis at a single point in the parameter space. A reasonable approach in a Bayesian context would be to base the choice of variables on the expected value of the preferred criterion, with expectation taken over the prior distribution of the parameters.

References

- BIERENS, H. J. (2007): “Econometric analysis of linearized singular dynamic stochastic general equilibrium models,” *Journal of Econometrics*, 136, 595–627.
- CANOVA, F., F. FERRONI, AND C. MATTHES (2014): “Choosing the variables to estimate singular DSGE models,” *Journal of Applied Econometrics*, 29, 1099–1117.
- ISKREV, N. (2010): “Evaluating the strength of identification in DSGE models. An a priori approach,” Working paper series, Banco de Portugal.
- ROTHENBERG, T. J. (1971): “Identification in Parametric Models,” *Econometrica*, 39, 577–91, available at <http://ideas.repec.org/a/ecm/emetrp/v39y1971i3p577-91.html>.
- SMETS, F. AND R. WOUTERS (2007): “Shocks and Frictions in US Business Cycles: A Bayesian DSGE Approach,” *The American Economic Review*, 97, 586–606.