



Munich Personal RePEc Archive

On the choice of most-preferred alternatives

Nikolai S. Kukushkin

9. November 2006

Online at <http://mpa.ub.uni-muenchen.de/803/>
MPRA Paper No. 803, posted 13. November 2006

On the Choice of Most-Preferred Alternatives*

Nikolai S. Kukushkin[†]

November 9, 2006

Abstract

Maximal elements of a binary relation on compact subsets of a metric space define a choice function. Necessary and sufficient conditions are found for: (1) the choice function to have nonempty values and be path independent; (2) the choice function to have nonempty values provided the underlying relation is an interval order. For interval orders and semiorders, the same properties are characterized in terms of representations in a chain.

Keywords: Maximal element; Path independence; Interval order; Semiorder

JEL classification: D 71.

*Financial support from a Presidential Grant for the State Support of the Leading Scientific Schools (NSh-5379.2006.1), the Mathematics Department of the Russian Academy of Sciences (program of basic research No. 3), the Russian Foundation for Basic Research (grant 05-01-00942), and the Spanish Ministry of Education (project SEJ 2004-00968) is acknowledged.

[†]Russian Academy of Sciences, Dorodnicyn Computing Center, 40, Vavilova, Moscow 119991 Russia
Tel.: +7-495-420-5707; fax: +7-495-135-6159. E-mail: ququns@pochta.ru

1 Introduction

The notion of a choice function plays a central role in the decision theory (Fishburn, 1973; Sen, 1984; Aizerman and Aleskerov, 1995), the most important being the case of a choice function defined by a binary relation. Traditionally, the attention was focused on choice functions on finite sets; connections between properties of such a choice function and properties of the underlying binary relation have been studied in detail. When the number of conceivable alternatives is large enough, a model with an infinite set of alternatives becomes appealing, but then many familiar results become irrelevant and intuitions developed over the finite case may prove misleading.

Let a binary relation be given on a metric space. There is a considerable literature studying conditions for the existence of maximal elements on compact sets (Gillies, 1959; Smith, 1974; Bergstrom, 1975; Mukherji, 1977; Walker, 1977; Kiruta et al., 1980; Danilov and Sotskov, 1985; Campbell and Walker, 1990). However, Kukushkin (2006) showed that no “simple” condition could characterize binary relations possessing undominated points in every compact subset.

Here we bypass the impossibility theorem by assuming (or demanding) a certain degree of rationality behind the binary relation (or the choice function it defines). A similar path was taken before by Smith (1974), who found a condition necessary and sufficient for an *ordering* (weak order) to admit a maximum on every compact subset of its domain. Theorem 1 below gives a condition for the choice function to be nonempty-valued and path independent on all compact subsets. Theorem 2, for the choice function to be nonempty-valued on all compact subsets provided the underlying relation is an interval order. Both conditions are equivalent (and equivalent to Smith’s condition) for semiorders (Theorem 3).

The “internal” characterization results are supplemented with “external” ones. As is well known, interval orders and semiorders can be represented with the help of mappings to a chain (Fishburn, 1985). Most popular are numeric representations (which need not exist generally), but, e.g., lexicographic scales may be preferable in certain cases. Theorems 4–6 characterize representations of interval orders ensuring the non-emptiness or path independence of choice; after minimal modifications, the theorems remain valid for semiorders.

The next section contains basic definitions. Section 3 establishes equivalence between “ ω -transitivity” and path independence; Section 4, between “ ω -acyclicity” and the existence of maximal elements for interval orders. Sections 5 and 6 contain characterization results in terms of representations in a chain for interval orders and semiorders, respectively.

2 Basic Notions

A *binary relation* on a set A is a Boolean function on $A \times A$; as usual, we write $y \succ x$ when the relation \succ is true on a pair (y, x) and $y \not\succeq x$ when it is false. An *interval order* is an irreflexive relation \succ such that

$$[y \succ x \ \& \ a \succ b] \Rightarrow [y \succ b \ \text{or} \ a \succ x]; \quad (1)$$

every interval order is transitive. An interval order is called a *semiorder* if

$$z \succ y \succ x \Rightarrow \forall a \in A [z \succ a \ \text{or} \ a \succ x]. \quad (2)$$

A relation \succ is called a *strict ordering* if it is irreflexive, transitive and *negatively transitive*, i.e., $z \not\succeq y \ \& \ y \not\succeq x \Rightarrow z \not\succeq x$. It is easy to see that every strict ordering is a semiorder.

Typically, we consider binary relations on a metric space A (a first countable Hausdorff topological space would do as well: what we actually need is that the topology on A be adequately described by convergent sequences). We denote \mathfrak{B} the lattice of all subsets of A and $\mathfrak{C} \subset \mathfrak{B}$ the set of all nonempty compact subsets of A (we never consider different sets A simultaneously).

Let $X \in \mathfrak{B}$; a point $x \in X$ is a *maximizer* for \succ on X if $y \not\succeq x$ for any $y \in X$. The set of all maximizers for \succ on X is denoted $M_\succ(X)$; we omit the subscript when the relation is clear from the context. A relation \succ has the *NM-property* on $X \subseteq A$ if for every $x \in X \setminus M_\succ(X)$ there is $y \in M_\succ(X)$ such that $y \succ x$. The property means that $M_\succ(X)$ is a von Neumann–Morgenstern solution on X .

$M_\succ(\cdot)$ defines a *choice function* on \mathfrak{B} ; we are mostly interested in properties of this function on \mathfrak{C} . First, whether all its values $M_\succ(X)$ are not empty (i.e., a choice can be made); second, whether the choice function is *path independent* on \mathfrak{B} (or \mathfrak{C}):

$$M(X \cup Y) = M(M(X) \cup Y) \tag{3}$$

for all $X, Y \in \mathfrak{B}$ (or whenever $X \cup Y \in \mathfrak{C}$).

Remark. This equality is usually perceived as a minimal rationality requirement. Plott’s (1973) original definition was a bit different from (3), but both are equivalent.

Given a binary relation \succ , an *improvement path* is a (finite or infinite) sequence $\langle x^k \rangle_{k=0,1,\dots}$ such that $x^{k+1} \succ x^k$ whenever both sides are defined. A relation \succ is *acyclic* if it admits no *finite improvement cycle*, i.e., no improvement path such that $x^m = x^0$ for an $m > 0$. Clearly, a relation is acyclic if and only if its transitive closure is irreflexive. A relation is *strongly acyclic* if it admits no infinite improvement path.

These two statements are well known (and easy to check anyway): a relation \succ is acyclic if and only if $M_\succ(X) \neq \emptyset$ for every finite $X \in \mathfrak{B} \setminus \{\emptyset\}$; a relation \succ is strongly acyclic if and only if $M_\succ(X) \neq \emptyset$ for every $X \in \mathfrak{B} \setminus \{\emptyset\}$.

A binary relation \succ on a metric space is called *ω -transitive* if it is transitive and, whenever $\langle x^k \rangle_{k=0,1,\dots}$ is an infinite improvement path and $x^k \rightarrow x^\omega$, there holds $x^\omega \succ x^0$. It is worth noting that $x^\omega \succ x^k$ is valid for all $k = 0, 1, \dots$ in this situation, once \succ is ω -transitive.

The notion seems to have been first considered by Gillies (1959), who proved its sufficiency for the existence of maximizers on compact sets. Smith (1974) gave it the name of “ σ -transitivity” and proved that it is *necessary* and sufficient for the existence of maximizers on all compact subsets provided \succ is a strict ordering. However, the prefix “ σ ” traditionally refers to the cardinal concept of a countable set whereas what matters here is the order type of the chain of natural numbers, usually referred to as ω .

A binary relation \succ is called *ω -acyclic* if it is acyclic and, whenever $\langle x^k \rangle_{k=0,1,\dots}$ is an infinite improvement path and $x^k \rightarrow x^\omega$, there holds $x^\omega \neq x^0$. It is worth noting that $x^k \succ x^\omega$ is impossible in this situation for any k once \succ is ω -acyclic. The property was considered by Mukherji (1977) as “Condition (A5).”

Let L be a chain and $C \subseteq L$; an *upper bound* for C is $u \in L$ such that $u \geq v$ for every $v \in C$. A *least upper bound* for C ($\sup C$) is an upper bound u such that $u \leq v$ for every upper bound v for C . If $\sup C$ exists at all, it is unique. Note that $\sup \emptyset$ is, by definition, the

least point of L . It is well known (Birkhoff, 1967) that a chain contains a least upper bound for every subset if and only if it is compact in its intrinsic topology; henceforth, we call such chains just *compact*.

Let A be a metric space and B be a partially ordered set. A mapping $\varphi : A \rightarrow B$ is called *upper ω -semicontinuous* if $\varphi(x^\omega) > \varphi(x^0)$ whenever $x^k \rightarrow x^\omega$ and $\varphi(x^{k+1}) > \varphi(x^k)$ for all $k = 0, 1, \dots$

3 Transitivity

Theorem 1. *Let \succ be a binary relation on a metric space. Then the following statements are equivalent.*

1.1. \succ is irreflexive and ω -transitive.

1.2. \succ has the NM property on every $X \in \mathfrak{C}$.

1.3. The choice function M_\succ is nonempty-valued and path independent on \mathfrak{C} .

Proof. Let us prove the implication [1.1] \Rightarrow [1.2] first. For each $x \in X$, we denote $G(x) = \{y \in X \mid y \succ x\}$. Once x^* is not a maximizer, we have $G(x^*) \neq \emptyset$. If we show that $M_\succ(G(x^*)) \neq \emptyset$, then we are obviously home.

To apply Zorn's Lemma (see, e.g., Kuratowski, 1966, p. 27), we have to consider an arbitrary chain $C \subseteq G(x^*)$ and show the existence of $y \in X$ such that $y \succ x$ or $y = x$ for each $x \in C$ (hence $y \in G(x^*)$). If C contains a greatest element, there is nothing to prove; otherwise $G(x) \neq \emptyset$ for each $x \in C$. We denote $F(x) = \text{cl } G(x)$ and $F = \bigcap_{x \in C} F(x)$. Since C is a chain, all the sets $G(x)$ ($x \in C$), hence $F(x)$ too, contain each other; therefore, every finite intersection of $F(x)$ is not empty. Since X is compact, $F \neq \emptyset$.

Let us prove that $y \succ x$ for each $y \in F$ and $x \in C$. Supposing the contrary, we pick $y \in F$ and $x^0 \in C$ for which $y \succ x^0$ does not hold and define a sequence $x^k \in G(x^0)$ ($k = 1, 2, \dots$) inductively so that $x^{k+1} \succ x^k$ and $x^k \rightarrow y$; then the ω -transitivity of \succ will imply that $y \succ x^0$, i.e., a contradiction. Having x^k already defined, we notice that $y \in F(x^k) \setminus G(x^k)$; therefore, we can pick $x^{k+1} \in G(x^k) \subseteq G(x^0)$ such that $0 < \rho(y, x^{k+1}) < \rho(y, x^k)/2$. Obviously, $x^{k+1} \succ x^k$ and $x^k \rightarrow y$.

Now let us turn to [1.2] \Rightarrow [1.3]. First, we notice that $M(X) \neq \emptyset$ for every $X \in \mathfrak{C}$ by the definition of the NM property. As to path independence, the inclusion $M(X \cup Y) \subseteq M(M(X) \cup Y)$ holds for maximizers of every binary relation and for all sets X and Y . Let $x \in (X \cup Y) \setminus M(X \cup Y)$ and $X \cup Y \in \mathfrak{C}$; by [1.2], there is $z \in M(X \cup Y) \subseteq M(M(X) \cup Y)$ such that $z \succ x$. Therefore, even if $x \in M(X) \cup Y$, $x \notin M(M(X) \cup Y)$.

Finally, let us prove [1.3] \Rightarrow [1.1]. If $x \succ x$, then $M(\{x\}) = \emptyset$. If $z \succ y \succ x$, then $\{z\} = M(\{x, y, z\}) = M(\{y, z\} \cup \{x\}) = M(\{x, z\})$, hence $z \succ x$. Let $x^k \rightarrow x^\omega$ and $x^{k+1} \succ x^k$ for each k . Denoting $X = \{x^\omega\} \cup \{x^k\}_{k=0,1,\dots}$ and $X' = X \setminus \{x^0\}$, we have $x^k \notin M(X)$ for each k , hence $M(X) = \{x^\omega\}$; similarly, $M(X') = \{x^\omega\}$. Now $\{x^\omega\} = M(X' \cup \{x^0\}) = M(M(X') \cup \{x^0\}) = M(\{x^\omega, x^0\})$, hence $x^\omega \succ x^0$. \square

Theorem 1 gives us a sufficient condition for the existence of maximizers. A *potential* for \succ is an irreflexive and ω -transitive relation \succsim finer than \succ , i.e., satisfying $y \succ x \Rightarrow y \succsim x$ for all $y, x \in A$. The notion was first introduced in Kukushkin (1999).

Corollary. *If \succ admits a potential, then $M_\succ(X) \neq \emptyset$ for each $X \in \mathfrak{C}$.*

Proof. Obviously, $M_{\succ}(X) \subseteq M_{\succ}(X)$ for any potential \succ for \succ . □

The Corollary often helps to establish the existence of maximizers.

The following statement is proven with arguments similar to the proof of Theorem 1 (and similar to the standard proof of the finite analogue).

Proposition 3.1. *Let \succ be a binary relation on a metric space. Then the following statements are equivalent.*

3.1.1. *\succ is transitive and strongly acyclic.*

3.1.2. *\succ has the NM property on every $X \in \mathfrak{B}$.*

3.1.3. *The choice function M_\succ is nonempty-valued and path independent on \mathfrak{B} .*

4 Acyclicity

Proposition 4.1. *If $M_\succ(X) \neq \emptyset$ for each $X \in \mathfrak{C}$, then \succ is ω -acyclic on A .*

Proof. If $\langle x^k \rangle_{k \in \mathbb{N}}$ is an improvement path converging to x^0 , then it is a compact subset without maximizers itself. (This argument was present in the proof of Corollary 3 from Mukherji, 1977, although the formulation was different.) □

The condition is by no means sufficient.

Example 4.1. Let A be a circle parameterized with real numbers from $[-\pi, \pi[$ (with “ $-\pi = \pi$ ”). We define \succ by

$$y \succ x \iff [\pi > y > x \geq 0 \text{ or } 0 > y > x \geq -\pi].$$

It is easily checked that \succ is transitive and ω -acyclic, but not ω -transitive. A itself is compact, but there is no maximizer on A .

Actually, the relation in Example 4.1 admits a cycle a bit different from those prohibited by the definition of ω -acyclicity. Let $x^k = (1 - 1/(k + 1))\pi$ and $y^k = -1/(k + 1)\pi$ ($k \in \mathbb{N}$); clearly, $x^{k+1} \succ x^k$ and $y^{k+1} \succ y^k$ for all k , while $x^k \rightarrow y^0$ and $y^k \rightarrow x^0$. As in Proposition 4.1, the absence of such cycles is also necessary for the nonemptiness of $M_\succ(X)$ for all $X \in \mathfrak{C}$. Similarly, one can define transfinite cycles of a greater length and again find that they must be prohibited too. However, a sufficient condition will never be obtained on this way.

Example 4.2. Let us consider a circle represented as the set of complex numbers with $|z| = 1$; formally, $A = \{e^{it} \mid t \in \mathbb{R}\}$. We define a binary relation $y \succ x \iff y = e^i \cdot x$. The relation is acyclic because 1 is incommensurable with 2π . It is ω -acyclic because no infinite improvement path, i.e., a sequence $\langle x^k \rangle_{k \in \mathbb{N}}$ such that $x^k = e^{ki} \cdot x^0$, can converge. Cycles similar to that from Example 4.1, or of a greater length, are impossible for the same reason. On the other hand, A itself is compact, but $M_\succ(A) = \emptyset$.

Here we have cycles in a different sense: every infinite improvement path is dense in A (the Jacobi theorem, see, e.g., Billingsley, 1965), hence its origin is among its limit points. The absence of *such* cycles, however, is not necessary for $M_{\succ}(X) \neq \emptyset$ for all $X \in \mathfrak{C}$.

With any binary relation \succ on A , we associate these two relations:

$$y \succ^{\omega} x \iff \exists \langle x^k \rangle_{k \in \mathbb{N}} [x^0 = x \ \& \ \forall k \in \mathbb{N} [x^{k+1} \succ x^k] \ \& \ x^k \rightarrow y]; \quad (4a)$$

$$y \succ x \iff [y \succ x \ \text{or} \ y \succ^{\omega} x]. \quad (4b)$$

Proposition 4.2. *If \succ is an ω -acyclic interval order, then \succ defined by (4) is irreflexive and ω -transitive.*

Proof. Irreflexivity of \succ immediately follows from the ω -acyclicity of \succ .

Lemma 4.2.1. *If $y \succ^{\omega} x$ and $z \succ y$, then $z \succ x$.*

Proof. Let $\langle x^k \rangle_{k \in \mathbb{N}}$ be an appropriate improvement path. From $z \succ y$ and $x^1 \succ x$, we derive that either $z \succ x$ or $x^1 \succ y$. The latter relation would contradict the ω -acyclicity of \succ (consider the sequence y, x^1, x^2, \dots). \square

Let $z \succ y \succ x$. If $y \succ x$, then obviously $z \succ x$; let $y \succ^{\omega} x$. If $z \succ y$, then Lemma 4.2.1 applies immediately. Let $z \succ^{\omega} y$ and $\langle y^k \rangle_{k \in \mathbb{N}}$ be an appropriate sequence. Lemma 4.2.1 implies that $y^1 \succ x$; therefore, $z \succ^{\omega} x$. Thus, \succ is transitive.

Let $x^k \rightarrow x^{\omega}$ and $x^{k+1} \succ x^k$ for all k . If $x^{k+1} \succ x^k$ for all k except for a finite number, a straightforward backward induction based on Lemma 4.2.1 shows that $x^{\omega} \succ^{\omega} x^0$. Otherwise, we may assume that $x^{k+1} \succ^{\omega} x^k$ for all k . Let $\langle x_k^h \rangle_{h \in \mathbb{N}}$ denote an appropriate sequence “between x^k and x^{k+1} ” (in particular, $x_k^0 = x^k$). Lemma 4.2.1 implies that $x_k^{h'} \succ x_k^h$ whenever $k' > k$ and $h' > 0$. We denote $y^0 = x^0$, pick an arbitrary sequence $r_k \rightarrow 0$, and, for each $k > 0$, pick $h(k) > 0$ such that $\rho(x_k^{h(k)}, x^{k+1}) < r_k$ and denote $y^k = x_k^{h(k)}$. Now we have $y^k \rightarrow x^{\omega}$ and $y^{k+1} \succ y^k$ for all k ; therefore, $x^{\omega} \succ^{\omega} x^0$ and \succ is ω -transitive. \square

Theorem 2. *Let \succ be an interval order on a metric space A . Then $M_{\succ}(X) \neq \emptyset$ for every $X \in \mathfrak{C}$ if and only if \succ is ω -acyclic.*

Proof. The necessity immediately follows from Proposition 4.1; the sufficiency, from Proposition 4.2 and Corollary to Theorem 1. \square

Campbell and Walker (1990) called a relation \succ “weak lower continuous” if $y \succ x$ implies the existence of an open neighborhood U of x such that $z \not\succeq y$ for every $z \in U$. Obviously, the weak lower continuity of \succ implies its ω -acyclicity; therefore, Theorem 1 of Campbell and Walker (when restricted to metric spaces) immediately follows from our Theorem 2. Weak lower continuity is not necessary for an interval order to admit a maximizer on every $X \in \mathfrak{C}$: consider a lexicographic order on a plane with fixed coordinates.

Theorem 3. *Let \succ be a semiorder; then the following statements are equivalent.*

3.1. $M_{\succ}(X) \neq \emptyset$ for every $X \in \mathfrak{C}$.

3.2. $M_{\succ}(X) \neq \emptyset$ for every $X \in \mathfrak{C}$ and $M_{\succ}(\cdot)$ is path independent on \mathfrak{C} .

3.3. \succ is ω -transitive.

3.4. \succ is ω -acyclic.

Proof. The implications [3.2] \Rightarrow [3.1] and [3.3] \Rightarrow [3.4] are obvious; [3.1] \Rightarrow [3.4] follows from Proposition 4.1; [3.2] \iff [3.3], from Theorem 1. Thus, it is sufficient to prove [3.4] \Rightarrow [3.3]. Let \succ be an ω -acyclic semiorder, $x^k \rightarrow x^\omega$, and $x^{k+1} \succ x^k$ for all k . Then $x^2 \succ x^1 \succ x^0$ implies that either $x^\omega \succ x^0$ or $x^0 \succ x^\omega$. The latter would contradict the ω -acyclicity (with the sequence $x^\omega, x^0, x^1, \dots$). \square

The restriction of the equivalence [3.3] \iff [3.1] to strict orderings renders the main theorem of Smith (1974, Theorem 4.1).

Example 4.3. Let $A = [0, 1]$ and $y \succ x \iff 1 > y > x$ for all $y, x \in A$. Then \succ is an interval order, ω -acyclic but not ω -transitive. Thus, Theorem 3 does not hold for interval orders. (Actually, there is no NM property on A itself, which is compact: any $x \in [0, 1[$ is neither a maximizer nor dominated by a maximizer.)

Every ω -acyclic interval order has an “ ε -version” of the NM property on every $X \in \mathfrak{C}$.

Proposition 4.3. *Let \succ be an ω -acyclic interval order on a compact metric space X and $x^* \in X \setminus M_\succ(X)$. Then there is $z \in M_\succ(X)$ for which either $z \succ x^*$ or there is an infinite improvement path $\langle z^k \rangle_{k \in \mathbb{N}}$ such that $z^0 = x^*$ and $z^k \rightarrow z$.*

Proof. Applying Theorem 1 to \succ defined by (4), we obtain $z \in M_\succ(X)$ such that $z \succ x^*$. A reference to (4b) completes the proof. \square

Remark. If A is finite, then every interval order generates a path independent choice function with nonempty values, so there is no analogue of the distinction between ω -transitivity and ω -acyclicity.

5 Representation of Interval Orders

Neither Proposition 5.1, nor its proof can claim any originality. However, both are needed in Theorems 4–6 and their corollaries.

Proposition 5.1. *Let \succ be a binary relation on a set A . Then \succ is an interval order if and only if there are a chain L and two mappings $\varphi^+, \varphi^- : A \rightarrow L$ such that, for all $x, y \in A$,*

$$\varphi^+(x) \geq \varphi^-(x); \tag{5a}$$

$$y \succ x \iff \varphi^-(y) > \varphi^+(x). \tag{5b}$$

Proof. Let \succ be defined by (5b), $y \succ x$, and $b \succ a$. Since L is a chain, we may, without restricting generality, assume $\varphi^-(y) \geq \varphi^-(b)$; then $\varphi^-(y) > \varphi^+(a)$, hence $y \succ a$. Thus, (5) imply (1).

Let \succ be an interval order. We consider \mathfrak{B} with set inclusion as a partially ordered set and define two mappings $\varphi^+, \varphi^- : A \rightarrow \mathfrak{B}$ by $\varphi^-(y) = \{x \in A \mid y \succ x\}$ and $\varphi^+(y) = \bigcup_{z \not\succ y} \varphi^-(z)$. By definition, $\varphi^-(y) \subseteq \varphi^+(y)$ for every $y \in A$; we denote $L = \varphi^-(A) \cup \varphi^+(A) \subseteq \mathfrak{B}$.

Let us show that L is a chain. If $\varphi^-(y_1) \setminus \varphi^-(y_2) \neq \emptyset \neq \varphi^-(y_2) \setminus \varphi^-(y_1)$ for some $y_1, y_2 \in A$, then there are $x_1, x_2 \in A$ such that $y_i \succ x_i$, but $y_i \not\succ x_j$ ($i, j \in \{1, 2\}$, $i \neq j$); clearly, this

contradicts (1). If $\varphi^+(y_1) \setminus \varphi^+(y_2) \neq \emptyset \neq \varphi^+(y_2) \setminus \varphi^+(y_1)$ for some $y_1, y_2 \in A$, there are $x_1, x_2, z_1, z_2 \in A$ such that $z_i \succ x_i$, $z_i \not\succeq y_i$, and $z_i \succ y_j$ ($i, j \in \{1, 2\}$, $i \neq j$); the configuration formed by z_1, z_2, y_1, y_2 violates (1). Comparing $\varphi^-(y_1)$ with $\varphi^+(y_2)$, we have two alternatives: if $\varphi^-(y_1) \subset \varphi^-(z)$ for some $z \not\succeq y_2$, then $\varphi^-(y_1) \subset \varphi^+(y_2)$; if $\varphi^-(y_1) \supseteq \varphi^-(z)$ for every $z \not\succeq y_2$, then $\varphi^-(y_1) \supseteq \varphi^+(y_2)$.

Let us check (5b). If $y \succ x$, then $\varphi^-(y) \ni x \notin \varphi^+(x)$; since L is a chain, there must be $\varphi^+(x) \subset \varphi^-(y)$. If $\varphi^+(x) \subset \varphi^-(y)$, we pick $z \in \varphi^-(y) \setminus \varphi^+(x)$. If $y \not\succeq x$, then $z \in \varphi^+(x)$ by definition; therefore, $y \succ x$. \square

Let \succ be an interval order represented in the sense of (5) with a compact chain L . For every $v \in L$, we define $\Xi(v) = \{x \in A \mid \varphi^+(x) \leq v\}$ and $\lambda(v) = \sup \varphi^-(\Xi(v))$; obviously, $v \geq \lambda(v)$ for every $v \in L$. A representation in the sense of (5) is called *regular* if L is compact and $\varphi^+(x) > \lambda \circ \varphi^+(x)$ for every $x \in A$.

Proposition 5.2. *Every interval order admits a regular representation.*

Proof. Let \succ be an interval order; by Proposition 5.1, there is a representation in the sense of (5). We define $\mathcal{L} = \{V \subseteq L \mid [v \in V \ \& \ v > w] \Rightarrow w \in V\}$ and $\iota : L \rightarrow \mathcal{L}$ by $\iota(v) = \{w \in L \mid v > w\}$. \mathcal{L} with set inclusion is a compact chain – $\sup C$ for $C \subseteq \mathcal{L}$ is the set union of C – while ι is an order-preserving embedding. Then we define $\mathcal{L}^* = \mathcal{L} \times \{0, 1\}$ with a lexicographic order: $(V', \vartheta') > (V, \vartheta) \iff [V' \supset V \text{ or } [V' = V \ \& \ \vartheta' > \vartheta]]$ for all $V', V \in \mathcal{L}$ and $\vartheta', \vartheta \in \{0, 1\}$; clearly, \mathcal{L}^* is a compact chain as well. Finally, we define $\psi^+, \psi^- : A \rightarrow \mathcal{L}^*$ by $\psi^+(x) = (\iota \circ \varphi^+(x), 1)$ and $\psi^-(x) = (\iota \circ \varphi^-(x), 0)$. Clearly, $\psi^-(y) > \psi^+(x) \iff \varphi^-(y) > \varphi^+(x)$ for all $y, x \in A$, hence ψ^- and ψ^+ provide a representation of \succ in the sense of (5). Besides, we have $\lambda \circ \psi^+(x) \leq (\iota \circ \varphi^+(x), 0) < \psi^+(x)$ for every $x \in A$. \square

Given a regular representation of an interval order \succ , we define subsets $S, T^-, T^+ \subseteq L$: $S = \{v \in L \mid v > \lambda(v)\} \cup \{\min L\}$ (the regularity implies $\varphi^+(A) \subseteq S$); $v \in T^-$ if and only if, whenever $x^k \rightarrow x^\omega$, $\varphi^+(x^{k+1}) > \varphi^+(x^k)$ and $\varphi^-(x^{k+1}) > \varphi^-(x^k)$ for all $k = 0, 1, \dots$, and $v = \sup_k \varphi^+(x^k) = \sup_k \varphi^-(x^k)$, there holds $\varphi^-(x^\omega) \geq v$; $v \in T^+$ if and only if, whenever $x^k \rightarrow x^\omega$, $\varphi^+(x^{k+1}) > \varphi^+(x^k)$ and $\varphi^-(x^{k+1}) > \varphi^-(x^k)$ for all $k = 0, 1, \dots$, and $v = \sup_k \varphi^+(x^k) = \sup_k \varphi^-(x^k)$, there holds $\varphi^+(x^\omega) \geq v$. Obviously, $T^- \subseteq T^+$; if $v \in S$, then $v \in T^-$ by default.

Theorem 4. *Let \succ be an interval order; then the following statements are equivalent.*

4.1. $S = L$ for every regular representation of \succ .

4.2. There is a regular representation of \succ with $S = L$.

4.3. \succ is strongly acyclic.

Proof. The implication [4.1] \Rightarrow [4.2] immediately follows from Proposition 5.2.

[4.2] \Rightarrow [4.3]: Suppose there is an infinite improvement path x^0, x^1, \dots ; then $\varphi^+(x^{k+1}) > \varphi^+(x^k)$ and $\varphi^-(x^{k+1}) > \varphi^-(x^k)$ for all k . We denote $v^+ = \sup_k \varphi^+(x^k)$ and $v^- = \sup_k \varphi^-(x^k)$; since $v^+ \in S$, we have $v^+ > \lambda(v^+) \geq v^-$. Therefore, there is k such that $\varphi^+(x^k) > v^- \geq \varphi^-(x^{k+1})$, but this contradicts the supposed $x^{k+1} \succ x^k$.

[4.3] \Rightarrow [4.1]: Let there be a regular representation of \succ and $v \in L \setminus S \subseteq L \setminus \varphi^+(A)$. We pick $x^0 \in \Xi(v)$ arbitrarily, and then define an infinite sequence of $x^k \in \Xi(v)$ inductively, in the following way. Since $\varphi^+(x^k) < v$ and $\lambda(v) = v$, there must be $x^{k+1} \in \Xi(v)$ such that $\varphi^-(x^{k+1}) > \varphi^+(x^k)$. Now we have $x^{k+1} \succ x^k$, i.e., a contradiction with the strong acyclicity of \succ . \square

The conditions of the theorem are satisfied if, e.g., \succ is represented by two numeric functions $\varphi^+, \varphi^- : A \rightarrow \mathbb{R}$ such that φ^+ is bounded above and $\varphi^+(x) \geq \varphi^-(x) + \varepsilon$ for all $x \in A$ with $\varepsilon > 0$.

Theorem 5. *Let \succ be an interval order on a metric space A ; then the following statements are equivalent.*

5.1. $T^- = L$ for every regular representation of \succ .

5.2. There is a regular representation of \succ with $T^- = L$.

5.3. \succ is ω -transitive.

Proof. [5.2] \Rightarrow [5.3]: Let x^0, x^1, \dots be an improvement path such that $x^k \rightarrow x^\omega$. Denoting $v = \sup_k \varphi^+(x^k)$, we immediately see that $v = \sup_k \varphi^-(x^k)$ as well; by [5.2], $v \in T^-$. Therefore, $\varphi^-(x^\omega) \geq v > \varphi^+(x^0)$, hence $x^\omega \succ x^0$.

[5.3] \Rightarrow [5.1]: Let there be a regular representation of \succ and $v \in L \setminus T^- \subseteq L \setminus \varphi^+(A)$. By definition, there is a sequence $x^k \rightarrow x^\omega$ such that $\varphi^+(x^{k+1}) > \varphi^+(x^k)$ and $\varphi^-(x^{k+1}) > \varphi^-(x^k)$ for all k , and $v = \sup_k \varphi^+(x^k) = \sup_k \varphi^-(x^k)$, but $\varphi^-(x^\omega) < v$; therefore, $\varphi^-(x^\omega) < \varphi^+(x^h)$ for all $h \in \mathbb{N}$ large enough. Since $\sup_k \varphi^+(x^k) = \sup_k \varphi^-(x^k)$, we may, deleting superfluous members from the sequence if needed, assume that $x^{k+1} \succ x^k$ for all k . Now the improvement path x^h, x^{h+1}, \dots violates the supposed ω -transitivity. \square

Proposition 5.3. *Let \succ be an interval order on a metric space A , represented in the sense of (5) by two mappings $\varphi^+, \varphi^- : A \rightarrow L$. Then \succ is ω -transitive if φ^- is upper ω -semicontinuous.*

Proof. If x^0, x^1, \dots is an improvement path such that $x^k \rightarrow x^\omega$, then $\varphi^-(x^{k+1}) > \varphi^-(x^k)$ for all k , and x^1, x^2, \dots also converges to x^ω . Therefore, $\varphi^-(x^\omega) > \varphi^-(x^1) > \varphi^+(x^0)$, hence $x^\omega \succ x^0$. \square

Example 5.1. Let $A = [0, 2]$ and $y \succ x \iff 2 > y > x + 1$. Clearly, \succ is transitive and strongly acyclic, hence ω -transitive by default; it can be represented in the sense of (5) by functions $\varphi^+(x) = x + 1$ for $0 \leq x < 2$, $\varphi^+(2) = 2$, and $\varphi^-(x) = \varphi^+(x) - 1$ for all $x \in A$, hence is an interval order. The representation is by no means unique, but it is easy to see that both φ^+ and φ^- must strictly increase for $x < 2$ and then jump down. In other words, the relation cannot be represented with an upper ω -semicontinuous function φ^- (or φ^+ for that matter).

Theorem 6. *Let \succ be an interval order on a metric space A ; then the following statements are equivalent.*

6.1. $T^+ = L$ for every regular representation of \succ .

6.2. There is a regular representation of \succ with $T^+ = L$.

6.3. \succ is ω -acyclic.

Proof. [6.2] \Rightarrow [6.3]: Let x^0, x^1, \dots be an improvement path such that $x^k \rightarrow x^\omega$. Arguing as in the proof of [5.2] \Rightarrow [5.3] above, we obtain $\varphi^+(x^\omega) > \varphi^+(x^0)$, hence $x^\omega \neq x^0$.

[6.3] \Rightarrow [6.1]: Let there be a regular representation of \succ and $v \in L \setminus T^+$. Arguing as in the proof of [5.3] \Rightarrow [5.1] above, we obtain an improvement path $x^k \rightarrow x^\omega$ such that $\varphi^+(x^\omega) < \varphi^+(x^h) < \varphi^-(x^{h+1})$, hence $x^{h+1} \succ x^\omega$. Now x^ω, x^{h+1}, \dots is an improvement path violating the supposed ω -acyclicity. \square

Proposition 5.4. *Let \succ be an interval order on a metric space A , represented in the sense of (5) by two mappings $\varphi^+, \varphi^- : A \rightarrow L$. Then \succ is ω -acyclic if φ^+ is upper ω -semicontinuous.*

The proof is similar to that of Proposition 5.3. Example 5.1 shows that Proposition 5.4 also cannot be reversed.

An upper semicontinuous numeric function $\varphi : A \rightarrow \mathbb{R}$ gives an obvious example of an upper ω -semicontinuous mapping to a chain. A bit more sophisticated examples are provided by lexicographic orders. Let L_1, \dots, L_n be chains and $L = \prod_{i=1}^n L_i$. The *lexicographic order* on L is defined in a usual way:

$$(v'_1, \dots, v'_n) > (v_1, \dots, v_n) \iff \exists m [\forall i < m [v'_i = v_i] \& v'_m > v_m].$$

Clearly, L with the order is a chain. A list of mappings $\varphi_i : A \rightarrow L_i$ ($i = 1, \dots, n$) defines a mapping $\varphi : A \rightarrow L$, their Cartesian product.

Proposition 5.5. *Let A be a metric space and each $\varphi_i : A \rightarrow \mathbb{R}$ ($i = 1, \dots, n$) be an upper semicontinuous function. Then the Cartesian product $\varphi : A \rightarrow \mathbb{R}^n$ (with the lexicographic order on \mathbb{R}^n) is upper ω -semicontinuous.*

Proof. Let $x^k \rightarrow x^\omega$ and $\varphi(x^{k+1}) > \varphi(x^k)$ for all $k = 0, 1, \dots$. For each k we denote μ_k ($1 \leq \mu_k \leq n$) the corresponding m from the definition of the lexicographic order. Without restricting generality, we may assume that $\mu_k = m$ for all k – we only have to delete a finite number of initial steps and then all steps with $\mu_k > m$. Therefore $\varphi_i(x^k)$ ($i < m$) does not depend on k , while $\varphi_m(x^{k+1}) > \varphi_m(x^k)$. From the upper semicontinuity of φ_m , we obtain $\varphi_m(x^\omega) > \varphi_m(x^0)$; from the upper semicontinuity of φ_i for $i < m$, $\varphi_i(x^\omega) \geq \varphi_i(x^0)$. Clearly, $\varphi(x^\omega) > \varphi(x^0)$. \square

6 Representation of Semiorders

The comment preceding Proposition 5.1 is relevant here as well.

Proposition 6.1. *Let \succ be a binary relation on a set A . Then \succ is a semiorder if and only if there are a chain L and two mappings $\varphi : A \rightarrow L$ and $\lambda : \varphi(A) \rightarrow L$ such that, for all $x, y \in A$:*

$$\varphi(y) > \varphi(x) \Rightarrow \lambda \circ \varphi(y) \geq \lambda \circ \varphi(x); \tag{6a}$$

$$\varphi(x) \geq \lambda \circ \varphi(x); \tag{6b}$$

$$y \succ x \iff \lambda \circ \varphi(y) > \varphi(x). \tag{6c}$$

Proof. Let (6) hold; we define two mappings $\varphi^+, \varphi^- : A \rightarrow L$ by $\varphi^+(x) = \varphi(x)$ and $\varphi^-(x) = \lambda \circ \varphi(x)$. Conditions (5) immediately follow from (6), hence \succ is an interval order by Proposition 5.1. Let $z \succ y \succ x$ and $a \in A$. If $\varphi(a) < \lambda \circ \varphi(z)$, then $z \succ a$; otherwise, we have $\varphi(a) \geq \lambda \circ \varphi(z) > \varphi(y)$, hence $\lambda \circ \varphi(a) \geq \lambda \circ \varphi(y) > \varphi(x)$, hence $a \succ x$.

Let \succ be a semiorder on A . Since \succ is an interval order, the necessity part of Proposition 5.1 applies. Let L, φ^+ and φ^- be as in the proof of the latter. We define $\mathcal{L} = L \times \{0, 1\} \times L$ with the lexicographic order, $\varphi : A \rightarrow \mathcal{L}$ by $\varphi(x) = (\varphi^+(x), 1, \varphi^-(x))$, and $\lambda : \varphi(A) \rightarrow \mathcal{L}$ by $\lambda(\varphi^+(x), 1, \varphi^-(x)) = (\varphi^-(x), 0, \varphi^-(x))$. By definition, (6b) holds as a strict inequality. To check (6c), it is enough to note that $\lambda \circ \varphi(y) > \varphi(x) \iff \varphi^-(y) > \varphi^+(x)$. To check (6a),

let us show that the inequalities $\varphi^+(y) > \varphi^+(x)$ and $\varphi^-(x) > \varphi^-(y)$ are incompatible for any $x, y \in A$. Supposing the contrary, we obtain the existence of $z \in A$ such that $x \succ z$ and $y \not\succeq z$ ($z \in \varphi^-(x) \setminus \varphi^-(y)$), and $a, b \in A$ such that $b \succ a$, $b \succ x$, and $b \not\succeq y$ ($a \in \varphi^+(y) \setminus \varphi^+(x)$). Now we have $b \succ x \succ z$ and $b \not\succeq y \not\succeq z$, i.e., a configuration violating (2). \square

A representation in the sense of (6) is called *regular* if L is compact and (6b) holds as a strict inequality.

Proposition 6.2. *Every semiorder admits a regular representation.*

Proof. The representation defined in the necessity proof of Proposition 6.1 satisfies (6b) as a strict inequality. Embedding into a compact chain can be performed exactly as in the proof of Proposition 5.2. \square

Given a regular representation of a semiorder \succ , we define $\Xi(v) = \{x \in A \mid \varphi(x) \leq v\}$ for every $v \in L$, and extend λ to $L \setminus \varphi(A)$ by $\lambda(v) = \sup \lambda \circ \varphi(\Xi(v))$; note that λ satisfies the same equality on $\varphi(A)$ as well. Now λ is increasing in the sense of $v' > v \Rightarrow \lambda(v') \geq \lambda(v)$ and satisfies $v \geq \lambda(v)$ on the whole L . We define subsets $S, T \subseteq L$: $S = \{v \in L \mid v > \lambda(v)\} \cup \{\min L\}$ (the regularity implies $\varphi^+(A) \subseteq S$); $v \in T$ if and only if, whenever $x^k \rightarrow x^\omega$, $\varphi(x^{k+1}) > \varphi(x^k)$ for all k , and $v = \sup_k \varphi(x^k) = \sup_k \lambda \circ \varphi(x^k)$, there holds $\varphi(x^\omega) \geq v$. If $v \in S$, then $v \in T$ by default.

Corollary to Theorem 4. *Let \succ be a semiorder; then the following statements are equivalent.*

1. $S = L$ for every regular representation of \succ .
2. There is a regular representation of \succ with $S = L$.
3. \succ is strongly acyclic.

Let $\varphi : A \rightarrow \mathbb{R}$ be bounded above and $\varepsilon > 0$; consider a relation defined by $y \succ x \iff \varphi(y) > \varphi(x) + \varepsilon$. A regular representation of \succ with $S = L$ is obtained if we define $L = [-\infty, u]$, where u is an upper bound for φ , and $\lambda(v) = v - \varepsilon$. Maximizers for \succ are exactly ε -maxima of φ .

Example 6.1. Let $x^k = k/(k+1)$ and $A = \{x^k\}_{k=0,1,\dots} \subset \mathbb{R}$. Clearly, the standard order on A is an ω -transitive semiorder, which is not strongly acyclic. A representation in the sense of (6) is given, e.g., by $L = A$, the identity mapping as φ , and $\lambda(x^{k+1}) = x^k$ for all k while $\lambda(x^0) = x^0$. We have $S = L$, but L is not compact. If we redefine $L = A \cup \{1\}$, we will have the compactness, but it will be impossible to extend λ to the new L keeping a strict inequality in (6b).

Corollary to Theorem 5. *Let \succ be a semiorder; then the following statements are equivalent.*

1. $T = L$ for every regular representation of \succ .
2. There is a regular representation of \succ with $T = L$.
3. \succ is ω -transitive.

Proposition 6.3. *Let \succ be a semiorder on a metric space A , represented in the sense of (6) by two mappings $\varphi : A \rightarrow L$ and $\lambda : \varphi(A) \rightarrow L$. Then \succ is ω -transitive if φ is upper ω -semicontinuous.*

The relation in Example 5.1 is an ω -transitive semiorder which admits no representation with an upper ω -semicontinuous function φ .

References

- Aizerman, M.A., and F.T. Aleskerov, 1995. *Theory of Choice*. North-Holland, Amsterdam.
- Bergstrom, T.C., 1975. Maximal elements of acyclic relations on compact sets. *Journal of Economic Theory* 10, 403–404.
- Billingsley, P., 1965. *Ergodic Theory and Information*. J. Wiley & Sons, N.Y.–London–Sydney.
- Birkhoff, G., 1967. *Lattice Theory*. American Mathematical Society, Providence.
- Campbell, D.E., and M. Walker, 1990. Maximal elements of weakly continuous relations. *Journal of Economic Theory* 50, 459–464.
- Danilov, V.I., and A.I. Sotskov, 1985. Rational choice and convex preferences. *Izvestiya Akademii Nauk SSSR, Tekhnicheskaya Kibernetika*, 2, 14–23. (in Russian)
- Fishburn, P.C., 1973. *The Theory of Social Choice*. Princeton University Press, Princeton.
- Fishburn, P.C., 1985. *Interval Orders and Interval Graphs: A Study of Partially Ordered Sets*. Wiley, N.Y.
- Gillies, D.B., 1959. Solutions to general non-zero-sum games. In: Tucker, A.W., and R.D. Luce (Eds.) *Contributions to the Theory of Games 4* (*Annals of Mathematical Studies* 40). Princeton University Press, Princeton, 47–85.
- Kiruta, A.Ya., A.M. Rubinov, and E.B. Yanovskaya, 1980. *Optimal Choice of Allocations in Complex Social-Economic Problems*. Nauka, Leningrad. (in Russian)
- Kukushkin, N.S., 1999. Potential games: A purely ordinal approach. *Economics Letters* 64, 279–283.
- Kukushkin, N.S., 2006. On the existence of maximal elements: An impossibility theorem. Available from <http://www.ccas.ru/mmes/mmeda/ququ/Max6.pdf>
- Kuratowski, K., 1966. *Topology*, Vol. 1. Academic Press, N.Y.–London.
- Mukherji, A., 1977. The existence of choice functions. *Econometrica* 45, 889–894.
- Plott, C.R., 1973. Path independence, rationality, and social choice. *Econometrica* 41, 1075–1091.
- Sen, A.K., 1984. *Collective Choice and Social Welfare*. North-Holland, Amsterdam.
- Smith, T.E., 1974. On the existence of most-preferred alternatives. *International Economic Review* 15, 184–194.
- Walker, M., 1977. On the existence of maximal elements. *Journal of Economic Theory* 16, 470–474.