



Munich Personal RePEc Archive

Comparing the Aitchison distance and the angular distance for use as inequality or disproportionality measures for votes and seats

Colignatus, Thomas

Thomas Cool Consultancy Econometrics

18 January 2018

Online at <https://mpra.ub.uni-muenchen.de/84334/>
MPRA Paper No. 84334, posted 04 Feb 2018 08:12 UTC

Comparing the Aitchison distance and the angular distance for use as inequality or disproportionality measures for votes and seats

Didactics and routines using *Mathematica*

Thomas Colignatus

<http://thomascool.eu>

January 18 & February 3 2018

Abstract

Votes and seats satisfy only two of seven criteria for application of the Aitchison distance. Vectors of votes and seats, say for elections for political parties the House of Representatives, can be normalised to 1 or 100%, and then have the outward appearance of compositional data. The Aitchison geometry and distance for compositional data then might be considered for votes and seats too. However, there is an essential zero when a party gets votes but doesn't gain a seat, and a zero gives an undefined logratio. In geology, changing from weights to volumes affects the percentages but not the Aitchison distance. For votes and seats there are no different scales or densities per party component however, and thus reportioning (perturbation) would be improper. Another key issue is subcompositional dominance. For votes {10, 20, 70} and seats {20, 10, 70} it is essential that we consider three parties. For a disproportionality measure we would value it positively that there is a match on 70. The Aitchison distance looks at the ratio $\{10, 20, 70\} / \{20, 10, 70\} = \{1/2, 2, 1\}$ and then neglects a ratio equal to 1. In this case it essentially compares the subcompositions, i.e. votes {10, 20} and seats {20, 10}, rescales to $\{1/3, 2/3\}$ and $\{2/3, 1/3\}$, and finds high disproportionality. This means that it essentially looks at a two party outcome instead of a three party outcome. It follows that votes and seats are better served by another distance measure. Suggested is the angular distance and the *Sine-Diagonal Inequality / Disproportionality* (SDID) measure based upon this. Users may of course apply both the angular and the Aitchison measures while being aware of the crucial differences in properties.

Keywords

Votes, Seats, Electoral System, Distance, Disproportionality, Aitchison Geometry, Angular Distance, Sine-Diagonal Inequality / Disproportionality, Loosemore-Hanby, Gallagher, Descriptive Statistics, Education, Reportion

MSC2010

00A69 General applied mathematics

28A75 Measure and integration. Length, area, volume, other geometric measure theory

62J20 Statistics. Diagnostics

97M70 Mathematics education. Behavioral and social sciences

JEL

A100 General Economics: General

D710 Social Choice; Clubs; Committees; Associations,

D720 Political Processes: Rent-seeking, Lobbying, Elections, Legislatures, and Voting Behavior

D630 Equity, Justice, Inequality, and Other Normative Criteria and Measurement

Cloud

This paper is also available at (updated version with link unchanged):

<https://www.wolframcloud.com/objects/thomas-cool/Voting/2018-01-18-Aitchison.nb>

I thank Stephen Wolfram for making this analysis possible in the Wolfram Cloud. I thank professors Vera Pawlowsky and Josep Antoni Martín-Fernández of UdG for essential comments. All errors remain mine.

Contents

1. Introduction
 2. Euclidean improduct, norm, distance and angle
 3. Theoretical requirements for a norm
 4. Disproportionality measures of votes and seats
 5. Compositional data
 6. Log Closure Plus One
 7. The Aitchison geometry
 8. Aitchison geometry and voting data
 9. Scoring parties on a policy scale
 10. Conclusion
- Appendix A. Formal properties of the Aitchison distance
- Appendix B. Approximation to the angle
- Appendix C. Votes and seats in Catalunya December 2017
- Appendix D. Aitchison distance and Webster / Sainte-Laguë
- Links
- References

Start (evaluate this subsection for the initialisation packages)

1. Introduction

This paper solves an issue on content and employs some didactics to do so.

On content: Votes and seats satisfy only two out of seven criteria for application of the Aitchison distance. It follows that votes and seats are better served by another distance measure, notably the angular distance and its transforms. Nevertheless, the Aitchison geometry might be a useful environment for the analysis of compositional data, and thus we should be aware of the properties.

On didactics: Mathematics provides clear definitions of **improduct**, **norm** and **distance**, and we are familiar with them for Euclidean space. When we apply these notions to different spaces then we find that understanding becomes still complicated. Especially education is challenged on those concepts for different spaces. It appears that *Mathematica* allows us to better grasp the issues. *Mathematica* is a system for doing mathematics on the computer. It allows the combination of

texts, symbols, numbers, graphs, patterns, motion, sound, etcetera. The following application to the Aitchison geometry again shows the power of using this integrated system.

We consider three spaces:

- (1) the original Euclidean space with vectors (of votes and seats) $x \geq 0$
- (2) the space with *shifted* logratio $\text{Log}[(x + 1) / (y + 1)]$
- (3) the space with *centered* logratio $\text{Log}[x / g[x]]$, with $y = g[x]$ the geometric mean.

We first review the notions of improduct, norm and distance for Euclidean space.

Subsequently we introduce the notion of **compositional data**: vectors $x > 0$ such that $1'x = 1$, or that the sum of the components add up to 1. These vectors are in the unit simplex, or, the endpoints of the vectors are onto the plane that characterises the unit simplex. Potentially some components might be zero, and these might be neglected unless there are *essential zeros*.

Our target application concerns *votes* and *seats*. For example, the outcomes for Parties *A* and *B* might be *votes* = {49, 51} and *seats* = {51, 49}, both in percentages. Those votes and seats are compositional data, at least in outward appearance. We would be interested in a measure of distance between these two outcomes. Political science on electoral systems has provided various measures of distance, of which we will mention the two main ones: The Loosemore-Hanby measure is based upon the *absolute difference* (ALHID) and the Gallagher measure is based upon the *Euclidean distance* (EGID), both with corrections to get an outcome in [0, 1], [0, 10] or [0, 100] depending upon the preferences of the researcher. It appears that these measures have some drawbacks. My suggestion from econometrics and political economy is to use the angle between the vectors, which in particular gives the transformation into the *Sine-Diagonal Inequality / Disproportionality* (SDID), see Colignatus (2017ac). For this present notebook it suffices to use a simpler indicator. Since the maximum angle between nonnegative vectors is 90 degrees, we can interpret the angle also as share of 90 degrees, henceforth *AngularID*.

The motivation for this notebook is that we should not overlook the suggestion of the Aitchison distance for compositional data. This notebook first discusses this geometry so that we understand its key properties. The two penultimate sections discuss the relevance of the Aitchison geometry for votes and seats. The **conclusion** is that the Aitchison geometry is less suited for a disproportionality measure for votes and seats. Obviously we may use both measures as long as we understand the properties.

For the literature on compositional data, it would be useful to distinguish between cases where the Aitchison geometry applies and those where it doesn't apply. Perhaps it might suffice to say that the criterion of being on the unit simplex is not enough to count as proper compositional data, so that votes and seats are only pseudo-compositional data.

For compositional data, the ratios x_i / x_j are regarded as generating more relevant information than the absolute values. Aitchison (1926-2016) proposed the "log-ratio" approach, using $\text{Log}[x_i / x_j] = \text{Log}[x_i] - \text{Log}[x_j]$ as the relevant measurement, corrected for the geometric means. The *centered logratio* approach essentially switches to the log of the *ratio to the geometric average*, and then applies the distance notions of the Euclidean space to those. To do so, however, still requires some essential understanding of geometry. Researchers in a particular field of interest obviously must understand what the transformations imply for the field of interest.

For proper compositional data, it appears that the *centered logratio* transformation is more important than the phenomenon that the data can be seen as being on the unit simplex. The conclusion of this present discussion may also be summarized as that the centered logratio transformation is inadequate for votes and seats.

For didactic reasons, we look first at an intermediate model that allows us to better grasp what Aitchison does. This approach uses the transform $\text{Log}[x + 1] \approx x$, which approximation is fairly good for values of $x \approx 0$. The addition of 1 moves the vector upwards but the logarithm moves it back towards the origin. We then look at the revised ratio $\text{Log}[(x_i + 1) / (x_j + 1)] = \text{Log}[x_i + 1] - \text{Log}[x_j + 1] \approx x_i - x_j$. This has *both* a notion of a ratio *and* outcomes similar to the absolute values. For example, comparing 10% to 5%, then $\text{Log}[10/5] = \text{Log}[2] \approx 70\%$ while $\text{Log}[1.1 / 1.05] \approx 5\%$. This didactic setup allows us to get used to the notion of a ratio and the log transform, and repeats the triad of *improduct, norm and distance*, so that we get a better feel for that triad. This approach however fails for votes and seats because there is no real departure from the absolute levels, and thus it suffers the same drawbacks as the Loosemore-Hanby and Gallagher measures. Comparing this $\text{Log}[x + 1]$ transform with the Aitchison approach shows that the latter deals much better with the ratios, so that we are more motivated to understand what the Aitchison geometry is.

This notebook uses v for votes, s for seats, $V = 1'v$ for the sum of votes, $S = 1's$ for the sum of seats, $w = v / V$, $z = s / S$ for the shares. The plots tend to use opposite formulas, as $z = 1 - w$, using $z = \{t, 1 - t\}$, with t for the seats of the first party. For votes and seats we prefer a normalisation to $[0, 10]$, given that $[0, 1]$ causes leading zeros and $[0, 100]$ suggests precision that is often lacking.

PM. This is the second version of the paper. The main change is that the Appendix E in the original text has been split off as an independent text Colignatus (2018). There is now also a stricter scaling of votes and seats into the $[0, \mathfrak{D}]$ range, default $\mathfrak{D} = 10$ (see Section 4).

NB. This discussion forms part of a larger framework given most recently by Colignatus (2017b): “*One woman, one vote. Though not in the USA, UK and France*”. My diagnosis is that “political science on electoral systems” is still in the Humanities and pre-science, notably by relying more upon common language instead of sharp definitions that are relevant for empirics. On the other hand, there are also mathematicians who deal with their definitions abstractly, without a proper grounding in empirical research. My invitation to empirical researchers is to help make a difference, notably in re-engineering the theory on electoral systems.

- <https://mpira.ub.uni-muenchen.de/82513/>

Routines in this notebook

? Cool`Statistics`Aitchison` *

▼ Cool`Statistics`Aitchison`

AitchisonDista- nce	AitchisonNorm	DistanceLCPO	LogClosurePlu- sOne	Stretch
AitchisonExpla- in	CenteredLogR- atio	ImprodLCPO	NormLCPO	UnAit
AitchisonImpr- oduct	Closure	InverseCLR	Reportion	

? Cool`Voting`Angular` *

▼ Cool`Voting`Angular`

AbsLoosemoreHandbyID	AngularPolicyDistancePlot	IndepAngPolicyDistance	UnitD
AllZeroButOneQ	AngularPolicyMeasure	MaxAngularPolicies	WAvAngPolicyMeasure
AngleProxy	AngularPolicyTerm	PlusAngPolicyDistance	WAvPlot
AngPolMax	CombinedPolicyDistance	PolicyNorm	WAvPolicyDistance
AngularID	D	SineDiagonalID	WAvSqrtPolicyDistance
AngularPolicyDistance	EuclidGallagherID	SingleMajority	WebsterSainteLaguereID

2. Euclidean improduct, norm, distance and angle

Pythagoras, length and size

In *Mathematica*, `Dimensions[x]` gives the list of dimensions of x . For a vector x , `Length[x]` gives the number of components.

```
{Dimensions[{a, b, c}], Length[{a, b, c}]}
{{3}, 3}
```

The Pythagorean Theorem allows us to find the *size length* of a vector. Another name is the **norm** $\|x\|$. *Mathematica* assumes complex vectors and thus uses `Abs[x]` for the size of the components. Since we use real vectors, we may eliminate this `Abs`.

? Norm

`Norm[expr]` gives the norm of a number, vector, or matrix.
`Norm[expr, p]` gives the p -norm. >>

```
Norm[{a, b, c}]
```

$$\sqrt{\text{Abs}[a]^2 + \text{Abs}[b]^2 + \text{Abs}[c]^2}$$

```
% /. Abs -> Identity
```

$$\sqrt{a^2 + b^2 + c^2}$$

The triad of improduct, norm and distance

Consider the steps: **improduct** \rightarrow **norm** \rightarrow **distance**.

In Euclidean space, the **improduct** is given as the Dot product $x \cdot y$. In linear algebra we write $x'y$.

? Dot

$a.b.c$ or `Dot[a, b, c]` gives products of vectors, matrices, and tensors. >>

$\{a, b, c\} \cdot \{x, y, z\}$

$a x + b y + c z$

For variables in real space, the norm of x amounts to application of the improduct to x and itself, and then taking the square root.

`Norm[{a, b, c}] == Sqrt[{a, b, c} . {a, b, c}]`

$$\sqrt{\text{Abs}[a]^2 + \text{Abs}[b]^2 + \text{Abs}[c]^2} = \sqrt{a^2 + b^2 + c^2}$$

The **distance** between vectors x and y is given as the norm of their difference. We look at real vectors, and eliminate `Abs`.

`Norm[{a, b, c} - {x, y, z}] /. Abs -> Identity`

$$\sqrt{(a-x)^2 + (b-y)^2 + (c-z)^2}$$


`EuclideanDistance[{a, b, c}, {x, y, z}] /. Abs -> Identity`

$$\sqrt{(a-x)^2 + (b-y)^2 + (c-z)^2}$$

Invariance to translation

A translation is an additive relocation by some z . The distance between x and y is unaffected by a translation to $x+z$ and $y+z$. This is easy to prove: $\text{Norm}[(x+z) - (y+z)] = \text{Norm}[x-y]$. The following shows the translation of two (black) vectors by $z = \{2, 1\}$ (red) giving two translated (blue) arrows.

```
org = {0, 0}; pa = {0, 3}; pb = {1, 0}; pat = pa + {2, 1}; pbt = pb + {2, 1};
lin1 = Line[{pa, pat, pbt, pb, pa}]; lin2 = Line[{pat, {2, 1}, pbt}];
arpa = Arrow[{org, pa}]; arpat = Arrow[{org, pat}];
arpb = Arrow[{org, pb}]; arpbt = Arrow[{org, pbt}];
arpz = Arrow[{org, {2, 1}}];
Graphics[{lin1, lin2, Thick, Arrowheads[.1], arpa, arpb, Blue,
  arpat, arpbt, Red, arpz}, Axes -> True, AspectRatio -> Automatic]
```



Subspace dominance

The size of vectors is always larger than their parts in subspaces. This is called subspace dominance.

Formally, $dist[s_1, s_2] \leq dist[x_1, x_2]$ when s_1 and s_2 are in the same subspaces of x_1 and x_2 .

Consider:

```
x1 = {.1, .2, .7}; s1 = Drop[x1, -1];
x2 = {.2, .1, .7}; s2 = Drop[x2, -1];
```

The final elements in x_1 and x_2 are the same, and this causes that x_1 and x_2 are as far apart as s_1 and s_2 . When the final elements in x_1 and x_2 would differ, this would only increase the Euclidean distance.

```
{EuclideanDistance[x1, x2], EuclideanDistance[s1, s2]}
{0.141421, 0.141421}
```

When we use the Norm then we see a zero where the vectors have the same values. Adding a zero size length obviously does not affect the outcome of the Norm.

```
{Hold[Norm][x1 - x2], Hold[Norm][s1 - s2]}
{Hold[Norm][{-0.1, 0.1, 0.}], Hold[Norm][{-0.1, 0.1}]}
```

```
% // ReleaseHold
{0.141421, 0.141421}
```

Metric

Given the properties of the Euclidean norm, the Euclidean distance d is a **metric**.

A distance d is a metric iff it satisfies these three properties:

- (1) $d[x, y] \geq 0$, and $d[x, y] = 0$ iff $x = y$
- (2) Symmetry: $d[x, y] = d[y, x]$
- (3) The triangular inequality: $d[x, z] \leq d[x, y] + d[y, z]$ for y other than x or z .

Angle and its estimate

Two vectors always form a plane. Improduct and norm allow us to define the cosine of an angle between vectors.

$$\text{Cos}[\theta] = \text{Cos}[x, y] = x \cdot y / (||x|| ||y||) = \text{Improduct}[x, y] / (\text{Norm}[x] \text{Norm}[y]).$$

The angle itself can be found by the inverse of the cosine, as $\theta = \text{ArcCos}[\text{Cos}[\theta]]$.

? VectorAngle

VectorAngle[u, v] gives the angle between the vectors u and v . >>

The angle distance is a metric. The “cosine distance” $1 - \text{Cos}[x, y]$ is not a metric because it does not satisfy the triangular inequality, see Van Dongen & Enright (2012).

- 1 - cos is a.k.a. the versed sine: <https://en.wikipedia.org/wiki/Versine>

Mathematica assumes complex variables, and thus its internal definition of the vector angle uses `Abs` and `Conjugate`. We can eliminate this however. We also look at nonnegative vectors, for which the maximum angle is 90 degrees. We get a relative measure of angular distance when we divide the angle by this maximum. We already showed this in the *Introduction* in numbers but let us repeat it here in symbols.

`VectorAngle[{a, b, c}, {d, e, f}] /. {Abs -> Identity, Conjugate -> Identity}`

$$\text{ArcCos} \left[\frac{a d + b e + c f}{\sqrt{a^2 + b^2 + c^2} \sqrt{d^2 + e^2 + f^2}} \right]$$

The AngularID (AID) scales to $[0, \mathfrak{D}]$, default $\mathfrak{D} = 10$. Thus, an AID outcome h stands for $90 h / 10 = 9 h$ degrees of the angle between the vectors.

? AngularID

AngularID[x, y] gives the angular inequality / disproportionality. This is the angle between real vectors x and y, divided by $\text{Pi}/2$ radian or 90 Degree, times \mathfrak{D} . Output is in $[0, \mathfrak{D}]$

The angular distance is independent from any positive scalar multiple per vector.

Simplify[

`AngularID[λ {a, b, c}, μ {d, e, f}] == AngularID[{a, b, c}, {d, e, f}]`
`], {λ > 0, μ > 0}]`

True

Appendix B shows an approximation of the angle $\theta \approx \|x / \|x\| - y / \|y\|\|$, which works good for small angles.

3. Theoretical requirements for a norm

It is useful to state the theoretical requirements of a norm, so that we can use these also for the Aitchison geometry.

A **norm** requires a **vector space**.

Definition of a vector space

A **vector space** has both **vector addition** and **scalar multiplication**.

(A) **Addition** has:

- (1) Symmetry / commutativity: $x + y = y + x$
- (2) Associativity: $x + (y + z) = (x + y) + z$
- (3) There is a neutral or identity element 0, such that $x + 0 = x$
- (4) Each element x has an inverse, denoted $-x$, such that $x + (-x) = 0$

(B) For **scalar multiples** λ and μ :

- (1) Associativity: $\lambda (\mu x) = (\lambda \mu) x$

(2) There is a neutral or identity element $\lambda = 1$, such that $\lambda x = x$

(3) Distributivity 1: $\lambda (x + y) = \lambda x + \lambda y$

(4) Distributivity 2: $(\lambda + \mu) x = \lambda x + \mu x$

Definition of a norm

When there is a vector space V , then the definition of a **norm** $norm: V \rightarrow Reals$ is:

(1) Homogeneity or absolutely scalable: $norm[\lambda x] = Abs[\lambda] norm[x]$

(2) Subadditive or triangular inequality: $norm[x + y] \leq norm[x] + norm[y]$

(3) Nonnegative: $norm[x] \geq 0$

(4) Definite: $norm[x] = 0$ iff $x = 0$

When distance $d[x, y] = norm[x - y]$ then the homogeneity of the norm gives $d[\lambda x, \lambda y] = Abs[\lambda] d[x, y]$

- wikipedia is a portal and no source
- https://en.wikipedia.org/wiki/Vector_space#Definition
- https://en.wikipedia.org/wiki/Distance#General_metric
- [https://en.wikipedia.org/wiki/Norm_\(mathematics\)#Euclidean_norm](https://en.wikipedia.org/wiki/Norm_(mathematics)#Euclidean_norm)

4. Inequality / disproportionality measures of votes and seats

This section is essentially copied from Colignatus (2018).

Normalisation to 10 rather than 1 or 100

For political science there is the choice between training students to grow sensitive to small numbers or create more sensitive measures that also allow direct communication with the larger public. For the Richter scale for earthquakes, the approach was to use logarithms, so that smaller values could be compared easier with larger ones. For our purposes the square root is adequate, with an easy change from 10 (grade) to 100 (percentage).

For this subject of votes and seats it is better to use the **grades** in $[0, 10]$, like $votes = \{4.9, 5.1\}$. The reason is that variables in $[0, 1]$ causes leading zeros, while percentages in the range $[0, 100]$ suggest a degree of precision that is often overdone.

For the scale we will use the symbol \mathfrak{D} (capital eth). For example $w = \mathfrak{D} v / V$. The default value in this paper is $\mathfrak{D} = 10$.

? Unit \mathfrak{D}

Unit $\mathfrak{D}[x, f:\mathfrak{D}]$ gives $f x / (\text{Plus } @@ \text{ Abs}[x])$

Unit $\mathfrak{D}[\text{Clear}]$ unprotects and clears \mathfrak{D}

Unit $\mathfrak{D}[\text{Set}, x]$ sets \mathfrak{D} to the value x and protects

The default protected value of \mathfrak{D} is 10

? D

The default value is 10, but it might be reset, see UnitD.

Compare the use of % as having the value 1/100. Write [ESC]D-[ESC] or "[CapitalEth]". Pronounce here as "deka" instead of "eth", and 1 / D as "decim"

UnitD[{49, 51}] // N

{4.9, 5.1}

1 / D == 10 Percent

$\frac{1}{10} == 10 \text{ Percent}$

Data of the US House of Representatives elections of 2016

It will be useful to have some example. At the US (half) elections of 2016 for the House of Representatives, the distributions for the votes and seats were, using the order {Democrats, Republicans, Other}:

votes = {dem = 61776554, rep = 63173815, 128627010 - rep - dem};

seats = {194, 241, 0};

- https://en.wikipedia.org/wiki/United_States_House_of_Representatives_elections,_2016#Results_summary

The operation of "closure" divides a vector by its sum total.

? Closure

Closure[x] gives $x / \text{Total}[x]$, i.e. divides x by


its sum. This assumes that x is a vector of positive real numbers

For votes we prefer to use UnitD.

{vts = UnitD[votes], sts = UnitD[seats]} // N

{ {4.80277, 4.9114, 0.285837}, {4.45977, 5.54023, 0.} }

```
Show[ListLinePlot[{{0, 0}, 6 {1, 1}},
  ListPlot[MapThread[Labeled, {Transpose[{vts, sts}], {Dem, Rep, Other}}],
  PlotStyle -> PointSize[Large]],
  AxesLabel -> {"Votes (D)", "Seats (D)"}, AspectRatio -> 1]
```



The USA have *district representation* which differs from *equal or proportional representation* (like in Holland). In a US district, a candidate is elected with the plurality rule. The votes for other candidates than the winner are discarded. That the Democrats and Republicans are still relatively close to the 45 Degree line of equality or proportionality is the effect of geography, and perhaps the median voter theorem. The word “election” is over-used when it refers to such different meanings for either district or proportional representation. Given the discarding of votes in the system of district representation, we rather should speak about “half elections”.

In district representation, there will also be more strategic voting, in which voters will not vote for a smaller party for fear that their vote will be discarded. This effect *cannot* be measured by the common inequality or disproportionality measures, since we only have the recorded votes and not the true preferences.

Disproportionality measures

The common notion is *distance* but for votes and seats we would like to see *equal proportions*, whence we also speak about an *inequality / disproportionality* (ID) measure.

The following measures are used in the literature on votes and seats, with the sine-diagonal measure a new suggestion from 2017. The measures are all symmetric, except Webster / Sainte-Laguë.


The literature uses values of ALHID and EGID in the [0, 100] range. We use the [0, 10] range here.

AbsLoosemoreHanbyID (ALHID)

? AbsLoosemoreHanbyID

AbsLoosemoreHanbyID[v, s] takes the absolute value of the differences of UnitD[v] and UnitD[s], divided by 2 to correct for double counting: $D * Total[Abs[v / v'1 - s / s'1]] / 2$. Outcomes are in the range [0, D]

```
Plot[AbsLoosemoreHanbyID[{D - t, t}, {t, D - t}], {t, 0, D}]
```



The LHID measure has the useful interpretation that it gives the share of displaced seats between parties for a House of D seats (corrected for double counting). In the US, 0.6 seats of a House of 10 seats are displaced, or 6 in a House of 100 seats.

```
AbsLoosemoreHanbyID[votes, seats] // N
```

```
0.628834
```

The ALHID distance is insensitive to the location of a 1 grade difference, say at $\{4, 6\}$ or $\{9, 1\}$.

```
{ad1 = AbsLoosemoreHanbyID[{4, 6}, {5, 5}],
  ad2 = AbsLoosemoreHanbyID[{9, 1}, {10, 0}], ad1 / ad2} // N
{1., 1., 1.}
```


- https://en.wikipedia.org/wiki/Loosemore%E2%80%93Hanby_index

EuclidGallagherID (EGID)

? EuclidGallagherID

EuclidGallagherID[v, s] takes the Euclided distance of Unit D [v] and Unit D [s], divided by Sqrt[2] to correct for double counting. Outcomes are in the range $[0, D]$. For length 2 the outcome is the same as AbsLoosemoreHanbyID

```
Plot[EuclidGallagherID[{D - t, t}, {t, D - t}], {t, 0, D}]
```



The 2016 US House data have 3 parties, and thus the EGID can have a different outcome than the ALHID. The US EGID is a bit larger than a half grade or 5%.

```
EuclidGallagherID[votes, seats] // N
```

```
0.545336
```

For two parties, the insensitivity is the same for EGID as ALHID.

```
{ad1 = EuclidGallagherID[{4, 6}, {5, 5}],
  ad2 = EuclidGallagherID[{9, 1}, {10, 0}], ad1 / ad2} // N
{1., 1., 1.}
```


- https://en.wikipedia.org/wiki/Gallagher_index

Angular inequality / disproportionality (AID)

? AngularID

AngularID[x, y] gives the angular inequality / disproportionality. This is the angle between real vectors x and y, divided by Pi/2 radian or 90 Degree, times D. Output is in [0, D]

```
Plot[AngularID[{D - t, t}, {t, D - t}], {t, 0, D}]
```



The 2016 US AID would suggest a disproportionality in the USA of 0.7 grades or 7%.

```
AngularID[votes, seats] // N
```

```
0.66848
```

The angular distance is sensitive to the location of 1 grade difference. It shows a halving or doubling of the angular distance, depending where one starts.

```
{ad1 = AngularID[{4, 6}, {5, 5}],
 ad2 = AngularID[{9, 1}, {10, 0}], ad1 / ad2} // N
{1.25666, 0.704466, 1.78385}
```

Because of this greater sensitivity the angular distance is a better measure for disproportionality than ALHID and EGID.

- https://en.wikipedia.org/wiki/Angular_distance


Sine-Diagonal (SDID)

? SineDiagonalID

SineDiagonalID[v, s] gives the Sqrt (greater sensitivity) of the Sine of the angle between the vectors, multiplied by 10 to get rid of leading zeros, and with the sign of Covariance[v, s] < -1 for majority switches. Thus it ranges in [-10, 10]


If we do not look at input of opposites.

```
Plot[SineDiagonalID[{50, 50}, {t, D - t}], {t, 0, D}]
```



The input of opposites generates negative outcomes, indicating majority switches between votes and seats.

```
Plot[SineDiagonalID[{D - t, t}, {t, D - t}], {t, 0, D}]
```



The SDID measure is designed to be sensitive to disproportionality. For the US it gives 3.2 on a scale of 10, thus 32%. The SDID uses a magnifying glass like the scale of Richter on earthquakes does. (Smaller earthquakes are made comparable with the bigger ones.)

```
SineDiagonalID[votes, seats] // N
```

```
3.23746
```

The SDID is steeper than the AID, which makes the ratio of inner versus outer range values for the displacement a bit lower.

```
{ad1 = SineDiagonalID[{4, 6}, {5, 5}],  
  ad2 = SineDiagonalID[{9, 1}, {10, 0}], ad1 / ad2} // N
```

```
{4.4285, 3.32312, 1.33263}
```

- <https://mpra.ub.uni-muenchen.de/81389/>

Webster / Sainte-Laguë (WSLID)


The measure is sensitive to ratios and without bound, so there is no real criterion to judge what an outcome really means.

? WebsterSainteLagueID

WebsterSainteLagueID[v, s] unitises $w = v / \text{Total}[v]$ and

$z = s / \text{Total}[s]$, and then gives $D w \cdot ((z/w - 1)^2) = D \text{Total}[(z - w)^2 / w]$



```
Plot[WebsterSainteLagueID[{D - t, t}, {t, D - t}], {t, 0, D}]
```



```
WebsterSainteLagueID[votes, seats] // N
0.390846
```

When a distance d has no upper value then it may always be transformed into $D d / (1 + d)$ to get a value in $[0, D]$ if needed.

```
Plot[(d = WebsterSainteLagueID[{D - t, t}, {t, D - t}] / D; D d / (1 + d)), {t, 0, D}]
```



- https://en.wikipedia.org/wiki/Webster/Sainte-Lagu%C3%AB_method


The Aitchison distance

Though we haven't defined the Aitchison distance here yet, we may already calculate and plot it with the same input as above.

? AitchisonDistance

AitchisonDistance[x, y] gives Norm[CenteredLogRatio[x] - CenteredLogRatio[y]] = Norm[CenteredLogRatio[x/y]] which is AitchisonNorm[x / y] which is also AitchisonNorm[Reportion[x, Stretch[y, -1]]]

```
Plot[AitchisonDistance[{D - t, t}, {t, D - t}], {t, 0, D}]
```




It is difficult to judge an outcome when there is no frame of reference. When a distance d has no upper value then it may always be transformed into $D d / (1 + d)$ to get a value in $[0, D]$ if needed. It itself this plot of the Aitchison distance suggests that it might be used, but below we will look at various properties that are awkward for votes and seats.

```
Limit[AitchisonDistance[{1 - t, t}, {t, 1 - t}], t -> 0]
```

∞

```
Plot[(d = AitchisonDistance[{D - t, t}, {t, D - t}]; D d / (1 + d)), {t, 0, D}]
```



For the US data, the Aitchison distance is undefined when there is a zero. Let us drop that case.

```
AitchisonDistance[Drop[votes, -1], Drop[seats, -1]] // N
```

0.137584

Appendix D discusses the relation between the Aitchison distance and the Webster / Sainte-Laguë inequality / disproportionality measure.

5. Compositional data

The unit simplex, closure and the neutral element

Compositional data are positive and their sum adds up to 1. Zeros would be eliminated, and all are divided by a constant.

The operation of “closure” divides a vector by its sum value.

? Closure

Closure[x] gives $x / \text{Total}[x]$, i.e. divides x by its sum. This assumes that x is a vector of positive real numbers

```
Closure[{x, y, z}]
```

$$\left\{ \frac{x}{x+y+z}, \frac{y}{x+y+z}, \frac{z}{x+y+z} \right\}$$

This means that all vectors are projected onto the unit simplex.

? Simplex

Simplex $\{p_1, \dots, p_k\}$ represents the simplex spanned by points p_i . >>

Since we will be looking at logarithms and $\text{Log}[1] = 0$, there is a special role for the unit vector $\{1, 1, \dots, 1\}$.

The projection of $\{1, 1, \dots, 1\}$ onto the unit simplex is called "UnAit" here.

? UnAit

UnAit[n] gives $\{1/n, \dots, 1/n\}$ for n positions, for integer n.

UnAit[] uses the integer from Options[UnAit]

UnAit[x] for vector x sets the integer option of UnAit to the length of x

Options [UnAit]


{Integer → 3}

UnAit []

$$\left\{ \frac{1}{3}, \frac{1}{3}, \frac{1}{3} \right\}$$

The following is an example from 3D, in which we project $\{1, 1, 1\}$ onto the unit simplex and find $\{1, 1, 1\}/3$.

```
zero = Table[0, {3}];
ones = Table[1, {3}];
simp = Simplex[{{1, 0, 0}, {0, 1, 0}, {0, 0, 1}}];
lin = Line[{zero, ones}];
pnt = Point[Closure[ones]];
Graphics3D[{simp, lin, Red, PointSize[Large], pnt}]
```



The “ternary plot” is again a special transformation of the unit simplex for 3D.

- For 3D, the ternary plot reworks the simplex: <http://mathworld.wolfram.com/TernaryDiagram.html>

Closure and angle

The angular distance is independent from closure, since this holds for any positive scalar multiple per vector.

As said we use the votes and seats of the US elections of 2016.

```

votes = {dem = 61776554, rep = 63173815, 128627010 - rep - dem};
seats = {194, 241, 0};
{vts = Closure[votes], sts = Closure[seats]} // N
{{0.480277, 0.49114, 0.0285837}, {0.445977, 0.554023, 0.}}

```

Closure is relevant for human understanding but not for the calculation of the angular distance, as this distance is unaffected by scalar multiples per vector.

```

{VectorAngle[votes, seats], VectorAngle[vts, sts]} / (90 Degree) // N
{0.066848, 0.066848}

```

The AngularID (AID) has been defined so that it scales into $[0, \mathbb{D}]$.

```

{AngularID[votes, seats], AngularID[vts, sts]} // N
{0.66848, 0.66848}

```

By comparison, to be discussed below, the Aitchison distance for American politics is, (i) depending upon our method to get rid of the zero, and (ii) without a reference like 100%:

```

AitchisonDistance[votes, seats + {0, -1, 1}] // N
2.07952

```

```

AitchisonDistance[Drop[votes, -1], Drop[seats, -1]] // N
0.137584

```

Practical cases and the essential zero

The notion of “Closure” may affect our perception of the data. We not only lose a degree of freedom. Aitchison has a strong argument that relative data are different from absolute data. The latter argument though seems more related to the use of the geometric mean than to closure, see below.

Examples of compositional data are:

- (i) geology: soil, e.g. sand, biological materials, some water
- (ii) biology: milk, e.g. fat, proteins, minerals, water (e.g. for different species like cows, cats, elephants, camels, ...)
- (iii) politics: votes and seats for various political parties.

There are some distinctions between these kinds of compositional data however. On the following three aspects, votes and seats differ from the first two examples.

(1) For soil and milk, one can take many samples from various sources, but for elections we essentially compare a single outcome of votes with a single outcome of seats. It makes little sense to take an average of votes over various countries when the parties are quite different. Even for the same country and same parties, comparison over years causes us to compare votes and seats in each year apart, and it makes little sense to average the votes and seats separately, and then take the disproportionality of those constructs.

(2) For soil and milk, there might be no “essential zeros” while for seats there is the essential phenomenon that there can be votes for parties who gain no seats (collected in the category “other”). If we switch to logarithms, then $\text{Log}[0]$ is undefined. NB. Parties that receive no votes may be

neglected, as they would neither get seats.

(3) The underlying issue for votes and seats derives most meaning from *apportionment* in which the number of available seats S is important. With apportionment function Ap , we determine: $seats = Ap[votes, S]$. Thus S has a key role. Votes and seats have the outward appearance of compositional data but this should not cause us to infer that they share the other properties.

- <https://en.wikipedia.org/wiki/Apportionment>

Aitchison's four criteria for distance

We can continue the list of aspects with four criteria on the distance measure (giving a total of seven criteria).

(4) Aitchison (undated), "Concise guide", p35, requires four properties of a compositional distance measure:

(a) *scale invariance*

(b) *permutation invariance*

(c) *perturbation invariance* (reportioning, componentwise rescaling)

(d) *subcompositional dominance*.

See also Martin-Fernandez et al. (undated) p2. The latter p3 show that the angular distance satisfies the first two properties but not the latter two. Thus, of the seven points considered here, votes and seats satisfy only two.

- http://ima.udg.edu/activitats/codawork05/A_concise_guide_to_compositional_data_analysis.pdf
- http://ima.udg.edu/~barcelo/index_archivos/Measures_of_difference__Clustering.pdf

Comments on these criteria are:

(4a) *Permutation invariance* means that the political parties can be sorted in whatever manner.

(4b) For *scaling*: Votes and seats have a natural scale, namely the natural numbers, and the invariance to scale must only hold for the vectors as a whole: dividing the votes by turnout and dividing the seats by the number of seats in the House.

(4c) For *reportioning (perturbation)*: This concerns different scales per component. For soil and milk, the components might be measured by weight or volume. If the components $\{x, y, z\}$ are in volumes and the components have different densities $\{a, b, c\}$, or weights per volume, then $\{ax, by, cz\}$ would be the composition in weights. These formats would give different vectors by closure. For multiplicative factors, rescaling each component by its own factor (*reportion* or *perturbation*) is the equivalent of the (linear) translation in Euclidean geometry (see the diagram above for translation). For soil and milk it would be relevant that the outcome of the analysis does not depend upon weight or volume. For votes and seats there are no different scales per party component, and thus this does not apply. We discuss this further below.

(4d) We will look at *subcompositional dominance* below.

Spurious correlation

Karl Pearson (1857-1936) coined the term "spurious correlation" by referring to a particular case of compositional data. It appears that his example is somewhat convoluted, see Colignatus (2017d).

The question is whether we correlate samples (rows) or components (columns). The term “spurious correlation” still is useful for when we have the wrong model and then find a correlation that does not reflect the true model. In the literature on compositional data, the reference to Pearson’s discussion is somewhat popular, using a similarly convoluted example. If you consider soil samples, and compare the compositions of the original samples with the compositions of those that have been dried (with removal of the water), then one might get different correlations. It must be doubted however whether this particular setup for the correlations is so relevant, when there is a confusion of samples (rows) and components (columns). Overall, my impression is that neither the reference to Pearson nor this particular example of “spurious correlation” of soil compositions are so enlightening. It is one distraction that one has to look into though, when the literature refers to it.

- <https://boycottholland.wordpress.com/2017/11/19/karl-pearsons-curious-construction-of-spurious-correlation/>

6. Log Closure Plus One

Rebasing to $\text{Log}[\text{Closure}[x] + 1]$

This section shows how a log transform can create something similar to a distance that is sensitive to some ratio. The transforms cause properties that deviate from both Euclid and Aitchison. This is intended as a didactic step between Euclidean and Aitchison distance measures.

? `LogClosurePlusOne`

`LogClosurePlusOne[x]` gives $\text{Log}[\text{Closure}[x] + 1] / \text{Log}[2]$. This assumes a nonnegative vector x with at least one positive value. Division by $\text{Log}[2]$ keeps corners like $\{1, 0, 0\}$ in position. Note: for small values $x \sim \text{Log}[x + 1]$. (Not implemented: All zeros may be translated in all zeros again.)

`LogClosurePlusOne[{x, y, z}]`

$$\left\{ \frac{\text{Log}\left[1 + \frac{x}{x+y+z}\right]}{\text{Log}[2]}, \frac{\text{Log}\left[1 + \frac{y}{x+y+z}\right]}{\text{Log}[2]}, \frac{\text{Log}\left[1 + \frac{z}{x+y+z}\right]}{\text{Log}[2]} \right\}$$

`LogClosurePlusOne[{0, 1, 0}]`


`{0, 1, 0}`

The transformation $x + 1$ maps x first with 1 to some “upper Simplex” beyond $\{1, 1, \dots, 1\}$, and then Log projects back again. The following example uses co-ordinates $cs = \{.10, .35, .55\}$ on the unit simplex. The blue line connects the origin to cs , to $\text{Closure}[cs] + 1 = cs + 1$, to $\text{Log}[cs + 1]$, to $\text{Log}[cs + 1] / \text{Log}[2]$. See also the numerical output. The transformation will generate outcomes close to the unit simplex.

```

zero = Table[0, {3}];
ones = Table[1, {3}];
simp = Simplex[{{1, 0, 0}, {0, 1, 0}, {0, 0, 1}}];
lin = Line[{zero, ones}];
pnt = Point[Closure[ones]];
pnt1 = Point[{cs = {.10, .35, .55}, cls = Closure[cs], cpo = cls + 1,
  lcpottrue = Log[cpo], lcpo = lcpottrue / Log[2]}];
lin1 = Line[{zero, cs, cls, cpo, lcpottrue, lcpo}];
Graphics3D[{simp, lin, Red, PointSize[Large], pnt, Blue, pnt1, lin1}]]
Point[{{0.1, 0.35, 0.55}, {0.1, 0.35, 0.55}, {1.1, 1.35, 1.55},
  {0.0953102, 0.300105, 0.438255}, {0.137504, 0.432959, 0.632268}}]]

```



The triad of improduct, norm and distance

Following the model of Euclidean space, we need an improduct, norm and distance. We can employ the Euclidean definitions to the transformed data since these are still within Euclidean space.

? ImprodLCPO

ImprodLCPO[x, y] applies the Euclidean improduct to the two Log[Closure[...]] + 1

ImprodLCPO[{a, b, c}, {d, e, f}]

$$\frac{\text{Log}\left[1 + \frac{a}{a+b+c}\right] \text{Log}\left[1 + \frac{d}{d+e+f}\right]}{\text{Log}[2]^2} + \frac{\text{Log}\left[1 + \frac{b}{a+b+c}\right] \text{Log}\left[1 + \frac{e}{d+e+f}\right]}{\text{Log}[2]^2} + \frac{\text{Log}\left[1 + \frac{c}{a+b+c}\right] \text{Log}\left[1 + \frac{f}{d+e+f}\right]}{\text{Log}[2]^2}$$

? NormLCPO

NormLCPO[x] applies the Euclidean Norm to Log[Closure[x] + 1]

NormLCPO[{a, b, c}]

$$\sqrt{\frac{\text{Log}\left[1 + \frac{a}{a+b+c}\right]^2}{\text{Log}[2]^2} + \frac{\text{Log}\left[1 + \frac{b}{a+b+c}\right]^2}{\text{Log}[2]^2} + \frac{\text{Log}\left[1 + \frac{c}{a+b+c}\right]^2}{\text{Log}[2]^2}}$$

? DistanceLCPO

DistanceLCPO[x, y] gives the Euclidean distance between the two Log[Closure[...] + 1]

DistanceLCPO[{a, b, c}, {d, e, f}]

$$\sqrt{\left(\frac{\text{Log}\left[1 + \frac{a}{a+b+c}\right]}{\text{Log}[2]} - \frac{\text{Log}\left[1 + \frac{d}{d+e+f}\right]}{\text{Log}[2]}\right)^2 + \left(\frac{\text{Log}\left[1 + \frac{b}{a+b+c}\right]}{\text{Log}[2]} - \frac{\text{Log}\left[1 + \frac{e}{d+e+f}\right]}{\text{Log}[2]}\right)^2 + \left(\frac{\text{Log}\left[1 + \frac{c}{a+b+c}\right]}{\text{Log}[2]} - \frac{\text{Log}\left[1 + \frac{f}{d+e+f}\right]}{\text{Log}[2]}\right)^2}$$

DistanceLCPO[x, y] gives Norm[LCPO[x] - LCPO[y]]. The subtraction means that we use something similar to the logratio: Log[(Closure[x] + 1) / (Closure[y] + 1)].

Evaluation of this didactic approach for votes and seats

Applying the routine to above votes and seats generates a distance of 0.08, which we should not interpret as 8% because it is not quite defined what 100% would be.

DistanceLCPO[seats, votes] // N

0.0796768

In voting theory the Loosemore-Hanby measure takes the sum of the absolute differences in the shares, and divides by 2 to correct for double counting. In a House of 100 seats, this gives the number of seats that are relocated between parties. In this case the Republicans got 6% of the seats more than their share of votes, taking the seats from the Democrats and the others.

Total[Abs[Closure[seats] - Closure[votes]]] / 2 // N

0.0628834

Our routine scales to [0, ∞].

AbsLoosemoreHanbyID[votes, seats] // N

0.628834

- https://en.wikipedia.org/wiki/Loosemore%E2%80%93Hanby_index

The result of the LCPO distance is close to the result of Loosemore-Hanby. We see it confirmed that there is no real difference between using LCPO and an absolute measure. Thus we have a stronger interest in finding out more about the Aitchison approach that focuses on the relative data.

7. The Aitchison geometry

Rebase to the geometric mean

We repeat the trick of transforming the data, and then using the Euclidean definitions on the transformed data.

In this case, we will rebase the original data to their geometric mean, and then take the log. This gives the *centered log-ratio* transform, normally abbreviated as (function) *clr*. The neutral element `UnAit[]` will transform to zero.

? CenteredLogRatio

`CenteredLogRatio[x]` gives $\text{Log}[x / \text{GeometricMean}[x]]$. This assumes a positive real vector x

```
CenteredLogRatio[UnAit[7]]
```

```
{0, 0, 0, 0, 0, 0, 0}
```

The geometric mean is only relevant for positive vectors. Since our vector of seats has an essential zero, we will flip one seat from a Republican to Other.

```
votes2 = Closure[votes] // N;
seats2 = Closure[seats + {0, -1, 1}] // N;
```

```
N[CenteredLogRatio /@ {seats2, votes2}]
{{1.68503, 1.89781, -3.58283}, {0.933053, 0.955419, -1.88847}}
```

The sum of centered logratio data is always 0.

```
Total /@ % // Chop
{0, 0}
```

Actually, using closure is not needed before we transform to the centered logratio. The latter transformation is also invariant for scalar multiples of the vectors. (A more complicated way of saying this is that it generates an equivalence class.)

```
CenteredLogRatio[{x, y, z}] == CenteredLogRatio[λ {x, y, z}]
```

$$\left\{ \text{Log}\left[\frac{x}{(xyz)^{1/3}}\right], \text{Log}\left[\frac{y}{(xyz)^{1/3}}\right], \text{Log}\left[\frac{z}{(xyz)^{1/3}}\right] \right\} ==$$

$$\left\{ \text{Log}\left[\frac{x\lambda}{(xyz\lambda^3)^{1/3}}\right], \text{Log}\left[\frac{y\lambda}{(xyz\lambda^3)^{1/3}}\right], \text{Log}\left[\frac{z\lambda}{(xyz\lambda^3)^{1/3}}\right] \right\}$$

```
% /. Log[x_] => Log[PowerExpand[x]]
True
```

When we use *votes* instead of *votes2* then the result does not change from the above.

```
N[CenteredLogRatio /@ {seats2, votes}]
{{1.68503, 1.89781, -3.58283}, {0.933053, 0.955419, -1.88847}}
```

Thus the story about the unit simplex is less relevant here than the story about the clr-transform.


Graphs

The following two graphs show the transformed data. We assume original co-ordinates that satisfy closure, with $x + y + z = 1$, and transform into new co-ordinates $\{x, y, z\} / (xyz)^{1/3}$ and their logarithms.


Perhaps a 2D graph is just as clear but we want to relate to the unit simplex above.

The first graph give the direct transform, the second graph gives the Log of this, and check that the latter is a plane (though *Mathematica* still draws contour lines because of minute numerical approximations).

```
ParametricPlot3D[z = 1 - x - y; If[z > 0, {x, y, z} / (x y z)^(1/3)],
  {x, 0, 1}, {y, 0, 1}, AspectRatio -> 1, ViewPoint -> Right]
```



```
ParametricPlot3D[z = 1 - x - y; If[z > 0, Log[{x, y, z} / (x y z)^(1/3)]],
  {x, 0, 1}, {y, 0, 1}, AspectRatio -> 1, PlotRange -> All]
```



The triad of improduct, norm and distance

For the *improduct*, *norm* and *distance* it suffices to use the centered logratio transform. We do not have to assume closure.

? AitchisonImproduct

AitchisonImproduct[x, y] gives CenteredLogRatio[x] . CenteredLogRatio[y]

AitchisonImproduct[{a, b, c}, {d, e, f}]

$$\begin{aligned} & \text{Log}\left[\frac{a}{(abc)^{1/3}}\right] \text{Log}\left[\frac{d}{(def)^{1/3}}\right] + \\ & \text{Log}\left[\frac{b}{(abc)^{1/3}}\right] \text{Log}\left[\frac{e}{(def)^{1/3}}\right] + \text{Log}\left[\frac{c}{(abc)^{1/3}}\right] \text{Log}\left[\frac{f}{(def)^{1/3}}\right] \end{aligned}$$

AitchisonImproduct[seats2, votes2] // N

10.1515

As said, we do not have to assume that the vectors satisfy closure.

AitchisonImproduct[seats2, votes] // N

10.1515

? AitchisonNorm

AitchisonNorm[x] gives Sqrt[AitchisonImproduct[x, x]]

AitchisonNorm[UnAit[]]

0

AitchisonNorm[{a, b, c}]

$$\sqrt{\text{Log}\left[\frac{a}{(abc)^{1/3}}\right]^2 + \text{Log}\left[\frac{b}{(abc)^{1/3}}\right]^2 + \text{Log}\left[\frac{c}{(abc)^{1/3}}\right]^2}$$

AitchisonNorm/@ {seats2, votes2} // N

{4.39063, 2.31295}

The Aitchison distance is the Euclidean distance for the rebased vectors.

If we do not use Norm but above AitchisonNorm then we divide instead of subtract, with the rationale that *division for real data is subtraction for logarithms*.

? AitchisonDistance

AitchisonDistance[x, y] gives Norm[CenteredLogRatio[x]

– CenteredLogRatio[y]] = Norm[CenteredLogRatio[x/y]] which is

AitchisonNorm[x / y] which is also AitchisonNorm[Reportion[x, Stretch[y, -1]]]

AitchisonDistance[{a, b, c}, {d, e, f}]

$$\sqrt{\text{Log}\left[\frac{a}{d \left(\frac{abc}{def}\right)^{1/3}}\right]^2 + \text{Log}\left[\frac{b}{e \left(\frac{abc}{def}\right)^{1/3}}\right]^2 + \text{Log}\left[\frac{c}{\left(\frac{abc}{def}\right)^{1/3} f}\right]^2}$$

```
AitchisonDistance[{a, b, c} / {d, e, f}, {1, 1, 1}]
```

$$\sqrt{\text{Log}\left[\frac{a}{d \left(\frac{abc}{def}\right)^{1/3}}\right]^2 + \text{Log}\left[\frac{b}{e \left(\frac{abc}{def}\right)^{1/3}}\right]^2 + \text{Log}\left[\frac{c}{\left(\frac{abc}{def}\right)^{1/3} f}\right]^2}$$

```
AitchisonDistance[seats2, votes2] // N
```

```
2.07952
```

The outcome 2.1 differs quite a bit from the earlier finding of the Loosemore-Hanby outcome of 6%, yet, we have no standard to judge that value 2.1.

PM 1. In the theory of votes and seats, researchers already considered using the $\text{Log}[\text{seats} / \text{votes}]$ transform. See **Appendix D** for its use in the Webster / Sainte-Laguë measure.

```
AitchisonDistance[seats2 / votes2, {1, 1, 1}] // N
```

```
2.07952
```

PM 2. Check again that it is not required to use closure data. (But for seats2 we flipped a seat.)

```
AitchisonDistance[seats2, votes] // N
```

```
2.07952
```

Underlying the Aitchison distance: Reportion and Stretch

For compositional data, Aitchison might have sufficed to argue that one should use the centered logratio transform, and then use the Euclidean inner product, norm and distance on those transformed vectors.

Instead, he *also* showed that this might be interpreted as a particular geometry on the unit simplex. The emphasis on "closure" and the unit simplex in the literature on the Aitchison distance only derives from this additional purpose. Since our present discussion aspires at a fair comparison of the methods, we now look at this additional purpose. Essentially, though, it is a distraction from the centered logratio transform.

Consider these two operations on the vectors in the unit simplex, which turn this into a *vector space*:

- (1) **Reportion**, a.k.a. perturbation, which has the role of "*addition*" for the vector space.
- (2) **Stretch**, a.k.a. power, which has the role of "*scalar multiplication*" for the vector space.

The routines assume that the input already is in closure.

Reportioning is when you have data in volumes and multiply these data with the densities to get data in weights.

? Reportion

Reportion[x, y] gives Closure[x y]. A.k.a. perturbation
or Log-translation. This assumes that x and y are vectors in Closure

```
Reportion[{a, b, c}, {d, e, f}]
```

$$\left\{ \frac{a d}{a d + b e + c f}, \frac{b e}{a d + b e + c f}, \frac{c f}{a d + b e + c f} \right\}$$

Stretching is the application of a uniform power to all elements.

? Stretch

Stretch[y, p] gives Closure[y^p]. A.k.a. power.

This assumes that p is a real number and that y is a vector in Closure

Stretch[{a, b, c}, p]

$$\left\{ \frac{a^p}{a^p + b^p + c^p}, \frac{b^p}{a^p + b^p + c^p}, \frac{c^p}{a^p + b^p + c^p} \right\}$$

Stretch[{a, b, c}, -1]

$$\left\{ \frac{1}{a \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right)}, \frac{1}{b \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right)}, \frac{1}{\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) c} \right\}$$

For example, when x is on the unit simplex, with $1^T x = 1$, then (i) $y = \sqrt{x}$ will be on the unit hypersphere, with $\|y\| = 1$, (ii) but Stretch[x, 1/2] will be on the unit simplex again.

`nc = Norm[Sqrt[Closure[{a, b, c}]]] /. Abs -> Identity // Simplify`

1

A vector space on the unit simplex with reportion and stretch

Reportioning and stretching generate a **vector space** for the variables under closure.

The Aitchison norm and distance satisfy the requirements for norm and distance, with the UnAit[] as the neutral element for addition. See the referred papers for the math. **Appendix A** shows this formally in *Mathematica* as well. For numbers it is more straightforward.

`Reportion[votes2, Stretch[votes2, -1]] == UnAit[3]`

True

For pairs x, there is the special property that the inverse operation $\text{Closure}[x^{(-1)}] = 1 - x$.

Stretch[{1/3, 2/3}, -1]

$$\left\{ \frac{2}{3}, \frac{1}{3} \right\}$$

Stretch[{x, 1-x}, -1] // Simplify

{1-x, x}

There seems to be a small tension between (a) the vector space under closure and (b) the use of Euclidean improduct, norm and distance for centered log-ratio vectors. However, the vector space under closure has its own definitions of improduct, norm and distance. Thus, when AitchisonNorm uses x/y in Euclidean terms, it actually uses Reportion[x, Stretch[y, -1]] for the variables under closure, whence we have a theoretically closed system.

? AitchisonDistance

AitchisonDistance[x, y] gives Norm[CenteredLogRatio[x] – CenteredLogRatio[y]] which is AitchisonNorm[x / y] which is also AitchisonNorm[Reportion[x, Stretch[y, –1]]]

AitchisonDistance[seats2, votes2]

2.07952

AitchisonNorm[Reportion[seats2, Stretch[votes2, –1]]]

2.07952

Still a question on conceptual consistency

While we have consistency in terms of logic, this still leaves a question on *conceptual* consistency.

(i) For the Aitchison distance, the *centered logratio* transform is more important than closure, since this distance is invariant to scalar multiples per vector (and thus also to the scalars generated by the closure operation).

(ii) Applying closure *afterwards* to clr-transformed data generates zeros. Thus clr-data are not on the unit simplex.

(iii) Thus it is true that multiplication and power (Reportion and Stretch) generate a vector space on the unit simplex, but this is rather irrelevant for *centered logratio* data, which are not on that simplex. The clr-data are in Euclidean space and we can still use the Euclidean improduct, norm and distance (which generates the Aitchison distance for the original data).

Thus, my take on this part of the literature is:

(1) For compositional data for which the ratios are important, one would use the *centered logratio* transform, and then use the normal Euclidean improduct, norm and distance on those transformed data, including regressions and such (minding the loss of 1 degree of freedom). Forecasts can be interpreted again using Exp or the InverseCLR transformation.

? InverseCLR

InverseCLR[x] gives Closure[Exp[x]], the inverse of the CenteredLogRatio

(2) The invariance of the Aitchison distance to *component-wise multiplication before the transform* (e.g. changing volumes in weights) depends upon the properties of the clr-transform. Idem dito for scalar multiples and the power relationship. Closure is not relevant here but only the clr-transform. Empirical researchers would diagnose that clr-transforms are the proper data to work with. They would still discuss whether to use weights or volumes (sensitive to temperature) though, since these would generate different clr-transforms (though with the same distances).

(3) The Aitchison geometry, with the interpretation of multiplication and power as creating a geometry on the unit simplex, is a mathematical exercise that is not relevant for empirical researchers. There is a connection between (1) & (2) and (3), in that logarithms change multiplication into addition, but that is all. The mathematical content is not large either, and perhaps suitable for a first year course on normed spaces. The discussion about this geometry is distractive for empirical researchers, who do not deal with the unit simplex but with the clr-transforms. (For

example, the creation of “lines” and “angles” on the unit simplex using this geometry is a mathematical exercise but conceptually at a distance for the clr-transforms that are not on the simplex.) It is true that one might interpret the camel as the “ship of the desert” but one should not be carried away by this interpretation.

(4) Thus the literature on “analysis on compositional data” can be cleaned up greatly but using these distinctions: (i) The explanation to empirical researchers on how to deal with compositional data that either satisfy the clr-transform (soil, milk) or not (votes and seats). (ii) A succinct text on that geometry for a first year course on normed spaces, not really relevant for researchers.

(5) I don’t expect but am still open to the possibility that the geometry might generate surprise insights for research.

8. Aitchison geometry and voting data

Judging the application of the Aitchison geometry requires knowledge about the particular field of application. As an econometrician and researcher in didactics of mathematics, I cannot judge upon applications for issues of geology or biology and such. The use of the voting data fall under “political science of electoral systems” and this area overlaps with “political economy” and “public choice”. See the reference to my paper on this.

- Use of the angular distance measure and its transform for voting data: <https://mpra.ub.uni-muenchen.de/81389/>

We are looking for properties that are essential for our understanding of proportionality between votes and seats: (1) properties that apply to both the angular and Aitchison distance but for which the first is simpler, (2) properties that apply to the angular distance but not the Aitchison distance, (3) properties that apply to only the Aitchison distance and not the angular distance. (Potentially there is also: (4) properties that apply to neither.)

We already mentioned some drawbacks for the application of the Aitchison geometry to voting data: (1) the pairwise comparison of votes and seats, instead of having lots of samples on the same issue like for soil, (2) the essential zeros for parties who get votes and not seats, (3) the importance of the total available seats S for $seats = Ap[votes, S]$, (4c) no adjustment of scales per party component.

Let us look further at (4c) Reportioning, (4d) subcompositional dominance.

Judging a distance by some reference

The Aitchison distance between the votes and seats of this basic example is, with an outcome close to 0:

AitchisonDistance [{49, 51}, {51, 49}]

$$\sqrt{2} \operatorname{Log} \left[\frac{51}{49} \right]$$

The Aitchison distance is invariant to reportioning:

`AitchisonDistance[{a, b} {49, 51}, {a, b} {51, 49}]`

$$\sqrt{2} \operatorname{Log} \left[\frac{51}{49} \right]$$

The Aitchison distance has a factor $\sqrt{2}$. We might divide standardly by this, like we do for the Euclid / Gallagher index to correct for double counting of relocated seats.

We still lack a yardstick to interpret what remains. A point of reference is perhaps this case that we would consider to be close to a maximal difference (or find a House with more than 1000 seats):

`AitchisonDistance[{999, 1}, {1, 999}]`

$$\sqrt{2} \operatorname{Log}[999]$$

Thus a relative distance might be:

`AitchisonDistance[{49, 51}, {51, 49}] / AitchisonDistance[{999, 1}, {1, 999}] // N`
 0.0057922

Section 4 already showed the transformation is $d / (1 + d)$, though we must beware that it becomes rather insensitive at the extremes.

Reporting of voting data 1: assuming perfection in apportionment

The question in this subsection is whether apportionment has the property $Ap[x v, S] = x Ap[v, S]$ for arbitrary nonnegative vectors x .

We already have $d[w, z] = d[w, Ap[w, S]]$. Let us not assume just now that $1'(x w) = 1$ too.

Above property gives $d[x w, Ap[x w, S]] = d[x w, x Ap[w, S]] = d[x w, x z]$. If apportionment causes an outcome $d[w, z]$ then with above property apportionment also causes $d[x w, x z]$ for any x . When this property would apply, then apportionment can be reduced to merely updating the seats with the change in votes.

In general, apportionment exists precisely since there is no such property that can be used like this. Perhaps for a rough estimate only.

If apportionment would be perfect then $z = w$, and $d[w, z] = d[w, w] = 0$. With this perfection, also $d[x w, Ap[x w, S]] = 0$. If above property would hold then $d[x w, x z] = 0$ too. Thus perfection and above property give $d[w, z] = d[x w, x z] = 0$ for any x . Unfortunately apportionment exists because there is no such perfection.

The following argues the same, now using explicitly two apportionments of seats given votes: $s1 = Ap[v1, S]$ and $s2 = Ap[v2, S]$.

When $\{v1, s1\}$ would be the results of one election, and when a new election generates $v2$, then an estimate would be $s1 * (v2 / v1)$ but we would rather await the process of apportionment as $s2 = Ap[v2, S]$.

Define $t = v2 / v1$ and $r = s2 / s1$. We have $s2 = Ap[t v1, S]$ and $s2 = r s1 = r Ap[v1, S]$.

Thus $Ap[t v1, S] = r Ap[v1, S]$. Can we really assume that $t \sim r$? In that case $Ap[x v, S] = x Ap[v, S]$.

In practice we will have a relation that $r = Ap[v t, S] / Ap[v, S] = r[v, t, S]$.

Suppose that we have a disproportionality measure such that $disp[v1, s1] \sim 0 \sim disp[v2, s2]$.

Obviously $disp[v_2, s_2] = disp[t v_1, r s_1] \sim disp[v_1, s_1]$, namely both close to zero.

Can this be seen as invariance of reportioning by $t \sim r$ or $v_2 / v_1 \sim s_2 / s_1$?

If apportionment Ap is truly proportional, so that $s_1 / v_1 = \lambda$ and $s_2 / v_2 = \mu$, then $r = s_2 / s_1 = (\mu v_2) / (\lambda v_1) = \mu / \lambda t$. Thus $r[v, t, S] = \mu / \lambda t$, in which the complexity has been hidden in λ and μ . Real numbers allow perfect apportionment, giving $z = w$, switching to unitised variables now. For perfection on these unitised variables, $\lambda = \mu = 1$. This is the case from above again, with $d[w, z] = 0$ and $d[x w, x z] = 0$, with the added restriction that $1'(x w) = 1$. For seats we have rounding however. This rounding should perhaps not affect the choice of the distance measure. But rounding may still imply that we cannot rely upon an invariance to reportioning.

In practice there will be few cases such that $t \sim r$ or $v_2 / v_1 \sim s_2 / s_1$, so that we may observe some invariance to reportioning.

It is more likely that there will be imperfections due to rounding. Invariance to reportioning would go against rounding.

The above assumes apportionment under a system of proportional representation. In district representation, the transformation of votes into seats is more complex, and there is no link to the supposed perfection of apportionment.

The comparison of votes and seats likely would do better by choosing an inequality / disproportionality measure that doesn't hinge upon invariance to reportioning.

The discussion whether such invariance $Ap[x v, S] = x Ap[v, S]$ is relevant or practically possible distracts from the very purpose of apportionment.

Reportioning of voting data 2: differential growth of party sizes

For v_1, v_2, s_1 and s_2 , let us suppose that Party A drops by about 40% of its votes and seats, and that Party B rises by about 40% of its votes and seats. These are only approximate effects, and we would get proper values from closure (and apportionment).

```
factor = {.6, 1.4}; {factor {4.9, 5.1}, factor {5.1, 4.9}}
{{2.94, 7.14}, {3.06, 6.86}}
```

In that case, the Aitchison distance remains the same. Let us use integers to generate the exact result.

```
factor = {6, 14}; {factor {49, 51}, factor {51, 49}}
{{294, 714}, {306, 686}}
```

```
factor = {6, 14}; AitchisonDistance[ factor {49, 51}, factor {51, 49}]
```

$$\sqrt{2} \operatorname{Log} \left[\frac{51}{49} \right]$$

```
% // N
```

```
0.0565761
```

By comparison, the angular distance is slightly reduced from 0.25 grade to 0.18 grade, or from 2.5% to 1.8%, since the resulting vectors are closer together indeed.

```
factor = {6, 14}; AngularID[factor {49, 51}, factor {51, 49}] // N
0.184422
```

An extreme effect uses $\{1/1000, 1000\}$. The Aitchison distance does not change, as this depends upon the ratio x_i / y_i while the angular distance becomes close to zero because votes and seats get close to $\{0, 1\}$.

```
factor = {1 / 1000, 1000}; AitchisonDistance[factor {49, 51}, factor {51, 49}]
 $\sqrt{2} \operatorname{Log}\left[\frac{51}{49}\right]$ 
```

```
factor = {1 / 1000, 1000}; AngularID[factor {49, 51}, factor {51, 49}] // N
 $5.10857 \times 10^{-7}$ 
```

PM. Closure has no effect on the outcome again, but can be shown anyway.

```
factor = {1 / 1000, 1000};
vts = Closure[factor {49, 51}];
sts = Closure[factor {51, 49}];
{vts, sts}
{ {  $\frac{49}{51000049}, \frac{51000000}{51000049}$  }, {  $\frac{51}{49000051}, \frac{49000000}{49000051}$  } }
{AitchisonDistance[vts, sts], AngularID[vts, sts] // N}
{  $\sqrt{2} \operatorname{Log}\left[\frac{51}{49}\right], 5.10857 \times 10^{-7}$  }
```

In sum, the differential growth of party sizes suggests that invariance in reportioning is counterproductive.

Reportioning of voting data 3: policy distance

The next main section looks at policy distance, and contains a role for the $\text{AngularID}[p\ v, p\ s]$, in which p are policy positions on a left-to-right scale $[0, 10]$. The measure used there cannot be based upon $\text{AitchisonDistance}[p\ v, p\ s]$ since this is equal to $\text{AitchisonDistance}[v, s]$.

Subcompositional dominance

For compositional data on minerals, a common dimension reduction is by dropping components in the vector, e.g. when the water content is removed, and the remainder components are rebased with “closure”. There would be spurious correlation in the original data when *columns on components* are used instead of *rows on samples*.

An example in the literature

Martin-Fernandez et al. (undated, p2) contain an example on subcompositional dominance.

- http://ima.udg.edu/~barcelo/index_archivos/Measures_of_difference__Clustering.pdf

Consider (using “s” for “sub” and not “seats”):

```
x1 = {.1, .2, .7}; s1 = Drop[x1, -1];
x2 = {.2, .1, .7}; s2 = Drop[x2, -1];
```

Subcompositional dominance would require that $dist[s1, s2] \leq dist[x1, x2]$.

The formal definition is provided in Aitchison (undated) “Concise guide”, p35.

- http://ima.udg.edu/activitats/codawork05/A_concise_guide_to_compositional_data_analysis.pdf

Let us first look at the angular distances and then the Aitchison distance.

The angular distances are, using radians to link up with table 3 on p4 in Martin-Fernandez et al..

```
{VectorAngle[x1, x2], VectorAngle[s1, s2], VectorAngle[Closure[s1], Closure[s2]]}
{0.192748, 0.643501, 0.643501}
```

For voting, these outcomes are logical. We should not confuse a comparison for three parties with a comparison for two parties. The participation of a third party may affect the evaluation of the total vectors of votes and seats.

- (i) The $x1$ and $x2$ are closer together because the third party has been valued accurately.
- (ii) When we consider the situation with only two parties, then $s1$ and $s2$ are further apart, as votes $\{1/3, 2/3\}$ have opposite seats $\{2/3, 1/3\}$.
- (iii) Thus, subcompositional dominance does not seem to be so useful for votes $x1$ and seats $x2$.

Degrees of freedom, dropping one element

This issue relates to the point that compositional data lose one degree of freedom. For the angular distance between votes and seats there is little reason to drop e.g. the last observation (potentially the essential zero) and then look at the angle between the remaining independent party shares. Namely, suppose that the last dependent share in each vector is the same, and that the other elements differ, like indeed is the case for above $x1$ and $x2$ and their subcompositions $s1$ and $s2$. Then the full vectors have a smaller angle, and we would want to see this reflected in the disproportionality measure (and thus we would not drop one element from the vector).

Ergo

Thus for voting we can properly infer that the situation with $x1$ and $x2$ with three parties is more proportional and that the situation with $s1$ and $s2$ with two parties is more disproportional.

The Aitchison distance however generates the same outcomes. The perfect match .7 for the third party is neglected, and the measure concentrates on where there are differences. We can also understand this outcome by reportioning as $x1 / x2 = \{1/2, 2, 1\}$, as another way to show that only the positions are used for which the ratios differ from 1.

```
{AitchisonDistance[x1, x2],
 AitchisonDistance[x1 / x2, {1, 1, 1}],
 AitchisonDistance[s1, s2],
 AitchisonDistance[Closure[s1], Closure[s2]]}
{0.980258, 0.980258, 0.980258, 0.980258}
```

Martin-Fernandez et al. (undated) conclude that the angular measure has no “compositional coherent behavior”. If votes and seats are regarded as compositional data, then there is a distinc-

tion between votes and seats on the one hand and soil and milk on the other hand.

PM. A bit more about the application of the angular measure in geology

Martin-Fernandez et al. refer to a paper by Watson & Philip 1989 who propose the use of the angular measure for geological compositions. Unfortunately that paper is behind a paywall. Aitchison, “Concise guide” p17 refers to this approach as being inappropriate for analysis of compositional data, referring to various other papers behind paywalls too.

- http://ima.udg.edu/~barcelo/index_archivos/Measures_of_difference__Clustering.pdf
- <https://link.springer.com/article/10.1007/BF00893217>

Wang et. al 2007 project a vector of dimension n into a vector of dimension $n - 1$ consisting of angles between components, yet, this might be dependent upon permutation, as one would get different angles for different permutations.

- <https://pdfs.semanticscholar.org/67e5/fa4a7cc9c833109722eed64fc3611c4e15f4.pdf>

See **Appendix B** for the relation between closure and the unit sphere.

9. Scoring parties on a policy scale

The parties can also be scored on a policy scale, say a vector p with elements in $[0, 10]$. A common term for such a scale is “left-to-right” even though those terms are rather out-dated. With unitised votes and seats w and z , a weighted average of the difference between House and electorate is $a = p \cdot (z - w)$. A basic reference is to Golder, M. & Stramski, J. (2010), “Ideological congruence and electoral institutions”, *American Journal of Political Science*, 54(1), 90-106. For example, they state (p95): “(...) substantively representative legislatures [for us now: $a = 0$] increase things like perceived levels of democratic legitimacy and responsiveness, satisfaction with democracy, political participation, or personal efficacy and trust in the political process.”

- <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.606.8475&rep=rep1&type=pdf>

As a political economist, I regard it as “revealed preference” (a term coined by Paul Samuelson) what parties are voted for and what policies those parties enact. When researchers try to determine what the “policy stance” p of the parties is then this seems operationally dubious and needlessly complicated, since we already have the distinctions between parties with their votes and seats, including the angular distance $d[v, s]$ between them. Remarkably, some researchers who prefer a policy distance tend to regard the distance between votes and seats as devoid of such meaning, and they thus do not value the interpretation in terms of revealed preference. The present discussion thus only occurs because there is a section in the literature that tries to look into such a policy distance between House and electorate.

The discussion on a policy distance measure has been split off as a separate paper, Colignatus (2018), since it distracts from the purpose of the present discussion, namely to compare with the Aitchison geometry.

In an earlier text (2017b, p45 and its Appendix J) I suggested that the angular distance $d[p v, p s]$ might be a policy distance itself. I retract this suggestion. This is only true for the perspective of $f[v, s; p] = d[p v, p s]$ for a given p . There is a different perspective for $g[p; v, s] = d[p v, p s]$ for given w

and z , which appears to be policy congruence and not incongruence. A widening policy gap between parties does not *increase* but *reduce* the angle of vectors $p v$ and $p s$, since the components actually come closer to each other. We already saw this above for differential growth of party sizes. A clear example is using $p = \{0, 10\}$ that causes an outcome of 0. Thus $d[p v, p s]$ is a measure of disproportionality given p and a measure of congruence given v and s . This point has now be included in Colignatus (2018), while (2017b) needs an update.

- <https://www.wolframcloud.com/objects/thomas-cool/Voting/2018-02-02-PolicyDistance.nbor> MPRA 84324
- Discussion of electoral systems: <https://mpira.ub.uni-muenchen.de/82513/>

Thus, Colignatus (2018) gives the revised suggestion for an angular policy distance $pd[v, s, p]$ that uses a transform of angular $d[p v, p s]$. Such a proposal would not work for the Aitchison distance, in which $dait[p v, p s] = dait[v, s]$ because of reportioning.

10. Conclusion

Vectors of votes and seats, for (half) elections for political parties for say the House of Representatives, can be normalised to 1 or 10 / \mathbb{D} or 100%, and then can be regarded as compositional data. The Aitchison geometry and distance for compositional data might also be applied to votes and seats. However, of seven criteria, votes and seats only satisfy two.

There is an essential zero when a party gets votes but doesn't gain seats. A zero gives an undefined logratio.

Votes and seats have a natural scale, namely the natural numbers, and the invariance to scale must only hold for the vectors as a whole: dividing the votes by turnout and dividing the seats by the number of seats in the House. The Aitchison distance is invariant to scale per component, so that e.g. volumes can be rescaled to weights. The latter property does not apply to votes and seats. One might argue that using the Aitchison distance for this case would not harm the analysis of the inequality / disproportionality of votes and seats. However, the issues of differential growth and the policy distance show that it does matter.

A key point is subcompositional dominance. For votes $\{1, 2, 7\}$ and seats $\{2, 1, 7\}$ it is essential that we consider three parties. For a distance or disproportionality measure we would value it positively that there is a match on 7. The Aitchison distance looks at the ratio $\{1, 2, 7\} / \{2, 1, 7\} = \{1/2, 2, 1\}$ and neglects a ratio equal to 1. It effectively compares the subcompositions, i.e. votes $\{1, 2\}$ and seats $\{2, 1\}$, rescales to $\{1/3, 2/3\}$ and $\{2/3, 1/3\}$, and finds high disproportionality. This means that it essentially looks at a two party outcome instead of a three party outcome.

It follows that votes and seats have a better measure in the angular distance (AID) and/or the *Sine-Diagonal Inequality / Disproportionality* (SDID) measure based upon this. One may of course use all measures, being aware of the crucial differences in meaning.

John Aitchison actually has two analyses. The literature would do well in making a clear distinction between: (1) the logratio transform and the Aitchison distance that are relevant for empirical research of compositional data, and (2) the Aitchison geometry on closure and the unit simplex that is mainly a mathematical exercise (and the logratio transforms are not on the unit simplex).

Appendix A. Formal properties of the Aitchison distance

The Aitchison distance is symmetric

Let us show $d[x, y] = d[y, x]$ for the Aitchison distance. For the following, application of FullSimplify for the domain of real vectors does not work. Let us get rid of the square root by squaring both sides of the equality sign.

$$\begin{aligned} \text{res} &= \text{AitchisonDistance}[\{a, b, c\}, \{d, e, f\}]^2 = \\ &\quad \text{AitchisonDistance}[\{d, e, f\}, \{a, b, c\}]^2 \\ &= \text{Log}\left[\frac{a}{d \left(\frac{abc}{def}\right)^{1/3}}\right]^2 + \text{Log}\left[\frac{b}{e \left(\frac{abc}{def}\right)^{1/3}}\right]^2 + \text{Log}\left[\frac{c}{\left(\frac{abc}{def}\right)^{1/3} f}\right]^2 = \\ &= \text{Log}\left[\frac{d}{a \left(\frac{def}{abc}\right)^{1/3}}\right]^2 + \text{Log}\left[\frac{e}{b \left(\frac{def}{abc}\right)^{1/3}}\right]^2 + \text{Log}\left[\frac{f}{c \left(\frac{def}{abc}\right)^{1/3}}\right]^2 \end{aligned}$$

It suffices that show equality for the first elements, for the others are similar.

$$\begin{aligned} \text{res}[[1, 1]] &= \text{res}[[2, 1]] \\ \text{Log}\left[\frac{a}{d \left(\frac{abc}{def}\right)^{1/3}}\right]^2 &= \text{Log}\left[\frac{d}{a \left(\frac{def}{abc}\right)^{1/3}}\right]^2 \end{aligned}$$

Only the ratios are relevant here.

$$\begin{aligned} \text{res2} &= \% /. \{d \rightarrow ka, e \rightarrow mb, f \rightarrow nc\} \\ \text{Log}\left[\frac{1}{k \left(\frac{1}{kmn}\right)^{1/3}}\right]^2 &= \text{Log}\left[\frac{k}{(kmn)^{1/3}}\right]^2 \end{aligned}$$

These expressions apparently are too difficult for Mathematica to deal with, but we can observe that the negative power within the logarithm is neutralised by the squares.

`% /. Log[x_] => Log[PowerExpand[x]]`

$$\text{Log}\left[\frac{m^{1/3} n^{1/3}}{k^{2/3}}\right]^2 = \text{Log}\left[\frac{k^{2/3}}{m^{1/3} n^{1/3}}\right]^2$$

`FullSimplify[%, {k, m, n} ∈ Reals]`

$$\text{Log}\left[\frac{k^{2/3}}{m^{1/3} n^{1/3}}\right]^2 = \text{Log}\left[\frac{m^{1/3} n^{1/3}}{k^{2/3}}\right]^2$$

The Aitchison norm has homogeneity or is absolutely scalable

This is $\text{norm}[\lambda x] = \text{Abs}[\lambda] \text{norm}[x]$. This also transfers to the distance.

`na = AitchisonNorm[{a, b, c}]`

$$\sqrt{\text{Log}\left[\frac{a}{(abc)^{1/3}}\right]^2 + \text{Log}\left[\frac{b}{(abc)^{1/3}}\right]^2 + \text{Log}\left[\frac{c}{(abc)^{1/3}}\right]^2}$$

nape = na /. Log[x_] := Log[PowerExpand[x]]

$$\sqrt{\text{Log}\left[\frac{a^{2/3}}{b^{1/3} c^{1/3}}\right]^2 + \text{Log}\left[\frac{b^{2/3}}{a^{1/3} c^{1/3}}\right]^2 + \text{Log}\left[\frac{c^{2/3}}{a^{1/3} b^{1/3}}\right]^2}$$

nap = AitchisonNorm[{a, b, c]^p]

$$\sqrt{\left(\text{Log}\left[\frac{a^p}{(a^p b^p c^p)^{1/3}}\right]^2 + \text{Log}\left[\frac{b^p}{(a^p b^p c^p)^{1/3}}\right]^2 + \text{Log}\left[\frac{c^p}{(a^p b^p c^p)^{1/3}}\right]^2\right)}$$

nape = nap /. Log[x_] := Log[PowerExpand[x]]

$$\sqrt{\left(\text{Log}\left[a^{2p/3} b^{-p/3} c^{-p/3}\right]^2 + \text{Log}\left[a^{-p/3} b^{2p/3} c^{-p/3}\right]^2 + \text{Log}\left[a^{-p/3} b^{-p/3} c^{2p/3}\right]^2\right)}$$

For $p > 0$: $\text{Sqrt}[\text{Log}[x]^2] = p / p \text{Sqrt}[\text{Log}[x]^2] = p \text{Sqrt}[(1/p \text{Log}[x])^2] = p \text{Sqrt}[\text{Log}[x^{(1/p)}]^2]$

napep = nape /. Log[x_] := Log[PowerExpand[x^(1/p)]]

$$\sqrt{\text{Log}\left[\frac{a^{2/3}}{b^{1/3} c^{1/3}}\right]^2 + \text{Log}\left[\frac{b^{2/3}}{a^{1/3} c^{1/3}}\right]^2 + \text{Log}\left[\frac{c^{2/3}}{a^{1/3} b^{1/3}}\right]^2}$$

nape == napep

True

The Aitchison distance remains the same under Reportioning

The Aitchison distance between vectors x and y remains the same when they are Reportioned with some z . Thus $\text{AitchisonDistance}[z x, z y] = \text{AitchisonDistance}[x, y]$. This is similar to the translation for Euclidean vectors.

daf = AitchisonDistance[{a, b, c}, {d, e, f}]

$$\sqrt{\text{Log}\left[\frac{0.0049056 a}{\left(\frac{a b c}{e f}\right)^{1/3}}\right]^2 + \text{Log}\left[\frac{14.2776 b}{e \left(\frac{a b c}{e f}\right)^{1/3}}\right]^2 + \text{Log}\left[\frac{14.2776 c}{\left(\frac{a b c}{e f}\right)^{1/3} f}\right]^2}$$

dafrep = AitchisonDistance[{x, y, z} {a, b, c}, {x, y, z} {d, e, f}]

$$\sqrt{\text{Log}\left[\frac{0.0049056 a}{\left(\frac{a b c}{e f}\right)^{1/3}}\right]^2 + \text{Log}\left[\frac{14.2776 b}{e \left(\frac{a b c}{e f}\right)^{1/3}}\right]^2 + \text{Log}\left[\frac{14.2776 c}{\left(\frac{a b c}{e f}\right)^{1/3} f}\right]^2}$$

daf == dafrep

True

Appendix B. Approximation to the angle

General property w.r.t. the unit circle

The transformation $x / \|x\|$ puts the vector on the unit circle (hypersphere).

A vector under closure has the property that the norm of its square root is 1, which means that this square root lies on the unit hypersphere too. Thus $x^* = x/\|x\|$ and $x^{**} = \text{Sqrt}[\text{Closure}[x]]$ are on the unit hypersphere, though still different.

```
xstar = {a, b, c} / Norm[{a, b, c}] /. Abs -> Identity
```

$$\left\{ \frac{a}{\sqrt{a^2 + b^2 + c^2}}, \frac{b}{\sqrt{a^2 + b^2 + c^2}}, \frac{c}{\sqrt{a^2 + b^2 + c^2}} \right\}$$

```
xdouble = Sqrt[Closure[{a, b, c}]]
```


$$\left\{ \sqrt{\frac{a}{a+b+c}}, \sqrt{\frac{b}{a+b+c}}, \sqrt{\frac{c}{a+b+c}} \right\}$$

```
Norm /@ {xstar, xdouble} /. Abs -> Identity // Simplify
```

```
{1, 1}
```

To show this property for vectors under closure in *Mathematica*, use `ParametricPlot` instead of `Plot`.

```
pp = ParametricPlot[Sqrt[{a, 1 - a}], {a, 0, 1}, AspectRatio -> 1];
pgr = Graphics[Circle[{0, 0}, 1, {0, Pi/2}]];
Show[pp, pgr]
```



Approximation of angle

Small angles between x and y have the interesting approximation given by $\|x/\|x\| - y/\|y\|\|$. The difference gives a chord that approximates the curve. See Maligranda, "Simple norm inequalities," *Amer. Math. Monthly* 113 (3), 256 – 260 (2006).

? `AngleProxy`

`AngleProxy[x, y]` gives $\text{Norm}[x/\text{Norm}[x] - y/\text{Norm}[y]]$ for real vectors (Abs -> Identity). Normalise by division by `Sqrt[2]` when comparing `{1, 0}` and `{0, 1}`

```
AngleProxy[x, y]
```

$$\text{Norm}\left[\frac{x}{\text{Norm}[x]} - \frac{y}{\text{Norm}[y]}\right]$$

For vectors that are close to each other, the approximation is remarkably well.

```
{VectorAngle[ {.49, .51}, {.51, .49}], AngleProxy[ {.49, .51}, {.51, .49} ]}
{0.0399947, 0.039992}
```


For orthogonal vectors, the chord gets length $\sqrt{2}$. For normalisation to 100% we would divide by $\sqrt{2}$ instead of $\pi/2$. However, when the proxy is intended for an angle then division by $\pi/2$ would be better still.


```
AngleProxy[{1 / 2, 0}, {0, 17}]
```

$$\sqrt{2}$$

```
{Sqrt[2], Pi / 2} // N
```

```
{1.41421, 1.5708}
```

```
org = {0, 0}; pa = {0, 3}; pb = {1, 0}; pat = pa + {2, 1}; pbt = pb + {2, 1};
panorm = pat / Norm[pat]; pbnorm = pbt / Norm[pbt];
lin1 = Line[{pat, pbt}]; lin2 = Line[{panorm, pbnorm}];
arpat = Arrow[{org, pat}]; arpanorm = Arrow[{or, panorm}];
arpbt = Arrow[{org, pbt}]; arpbnorm = Arrow[{or, pbnorm}];
unitcircle = Circle[{0, 0}, 1, {0, Pi / 2}];
Graphics[{lin1, lin2, unitcircle, Thick, Arrowheads[.1], Blue, arpat,
  arpbt, Red, arpanorm, arpbnorm}, Axes → True, AspectRatio → Automatic]
```



Application to votes and seats

If we would work with variables x that are transformed to $x^{**} = \text{Sqrt}[\text{Closure}[x]]$ then the Sqrt moves the data from the original ray. The angle between x and y differs from the angle between x^{**} and y^{**} . That the $\text{Norm}[x^{**} - y^{**}]$ would still be an approximation to the angle between x^{**} and y^{**} should not distract us.

Take for example the 2016 (half) elections for the US House of Representatives again.

```
votes = {rep = 63173815, dem = 61776554, 128627010 - rep - dem};
seats = {241, 194, 0};
{vts = Closure[votes], sts = Closure[seats]} // N
{{0.49114, 0.480277, 0.0285837}, {0.554023, 0.445977, 0.}}
```

- https://en.wikipedia.org/wiki/United_States_House_of_Representatives_elections,_2016#Results_summary

The angle between the vectors is (and differs from the angle between their square roots):

```
VectorAngle[votes, seats] // N
0.105005
```

The sqrt transformation to the hypersphere is:

```
{vhyp, shyp} = Sqrt[{vts, sts}] // N
{{0.700814, 0.69302, 0.169067}, {0.744327, 0.667815, 0.}}
```

The approximation by the Norm is remarkably close to the true angle between these variables:

```
{VectorAngle[vhyp, shyp], Norm[vhyp - shyp]}
{0.176617, 0.176387}
```

However, the angle between votes and seats (0.105) differs from the angle of the Sqrt variables (0.177). It is not clear why the latter would be a more useful transform. Merely the property that their angle and Euclidean distance are approximately the same, is no argument on content.

Wang et al. use this square root transformation and the angles, yet, the angles also depend upon the permutation of parties.

- <https://pdfs.semanticscholar.org/67e5/fa4a7cc9c833109722eed64fc3611c4e15f4.pdf>

Appendix C. Votes and seats in Catalunya December 2017

The CoDaWeb research group is established at Girona, Catalunya, Spain.

- Website: <http://www.compositionaldata.com/>

It is somewhat remarkable what is happening in Catalunya in terms of electoral systems. In Autumn 2017, the regional government of Catalunya, Spain, declared independence without implementing it. The Spanish government dissolved this regional government and organised new elections. There are charges of corruption and that parties emphasized independence to distract attention away from this corruption. Because of the electoral system of district representation, the political parties in favour of independence of Catalunya got no majority in votes but still a majority in seats.

- <https://www.irishtimes.com/news/world/europe/corruption-sentence-hits-catalonia-s-dominant-nationalist-party-1.3356372>

Wikipedia is a portal and no source, so in this case I am hoping that they reported correctly.

- January 15 2018: https://en.wikipedia.org/wiki/Catalan_regional_election,_2017#Results

```
data = {{"Names", "Labels", "Votes", "Seats", "1 = Independence"},
  {"Citizens-Party of the Citizenry", Cs, 1109732, 36, 0},
  {"Together for Catalonia", JuntsxCat, 948233, 34, 1},
  {"Republican Left-Catalonia Yes", ERC_CatSí, 935861, 32, 1},
  {"Socialists' Party of Catalonia", PSC_PSOE, 606659, 17, 0},
  {"Catalonia in Common-We Can", CatComú_Podem, 326360, 8, 0},
  {"Popular Unity Candidacy", CUP, 195246, 4, 1},
  {"People's Party", PP, 185670, 4, 0},
  {"Other, Blank, Invalid", Other, 49607 + 19431 + 16092, 0, 0}};
```

```
data // MatrixForm
(
  Names                Labels      Votes   Seats   1 = Independence
Citizens-Party of the Citizenry    Cs      1109732  36      0
  Together for Catalonia             JuntsxCat  948233  34      1
  Republican Left-Catalonia Yes     ERC_CatSí  935861  32      1
  Socialists' Party of Catalonia    PSC_PSOE   606659  17      0
  Catalonia in Common-We Can       CatComú_Podem  326360  8       0
  Popular Unity Candidacy           CUP        195246  4       1
  People's Party                    PP         185670  4       0
  Other, Blank, Invalid             Other      85130   0       0
)
```

```
{names, labels, votes, seats, indep} = Transpose[Rest[data]];
```

Let us use the term $\text{Unit}\mathcal{D}$ with default values in $[0, \mathcal{D}]$, default $\mathcal{D} = 10$, instead of “Closure and 100”.

```
vts = UnitD[votes] // N
```

```
sts = UnitD[seats] // N
```

```
{2.5262, 2.15856, 2.1304, 1.381, 0.742928, 0.444459, 0.42266, 0.19379}
```

```
{2.66667, 2.51852, 2.37037, 1.25926, 0.592593, 0.296296, 0.296296, 0.}
```

```
labels . indep
```

```
CUP + JuntsxCat + ERC_CatSí
```

The independence parties have $4.7/\mathcal{D}$ of the votes but $5.2/\mathcal{D}$ of the seats. Catalunya has district representation instead of equal or proportional representation. See Colignatus (2017b) for the conceptual errors of district representation.

```
{vts . indep, sts . indep}
```

```
{4.73342, 5.18519}
```

The two indicators for electoral disproportionality on a scale of $[0, \mathcal{D}]$ are (with the policy distance also taking variation in the group):

```
{AngularID[vts, sts], AngularPolicyDistance[vts, sts, indep]}
```

```
{0.680163, 1.24041}
```

The Aitchison distance requires us to drop the last zero, while the outcome is hard to judge because we lack some reference like 100%.

```
AitchisonDistance[Drop[vts, -1], Drop[sts, -1]]
```

```
0.552229
```

Appendix D. Aitchison distance and Webster / Sainte-Laguë

Use unitised votes w and seats z . Let us write $z/w = x + 1$, then $\text{Log}[z/w] = \text{Log}[x + 1] \approx x = z/w - 1$.

The Webster / Sainte-Laguë (WSL) disproportionality measure uses $\text{Sum}[w (z/w - 1)^2] = \text{Sum}[(z - w)^2 / w]$.

Thus $WSL \approx \text{Sum}[w \text{Log}[z/w]^2]$. This is a weighted norm of $\text{Log}[z/w]$.

The Aitchison distance $d[w, z] = d[1, z/w] = \text{Norm}[\text{CenteredLogRatio}[z/w]]$, using reporting.

We may also write $\text{Norm}[\text{Log}[z/w] - \text{Log}[\text{GeometricMean}[z/w]]]$.

Thus WSL uses a *weighted norm* of $\text{Log}[z/w]$ while Aitchison uses the *centered* norm of $\text{Log}[z/w]$.

The properties and outcomes differ, of course.

Links

Some relevant links are:

- Use of the angular distance measure for voting data: <https://mpira.ub.uni-muenchen.de/81389/>
- Discussion of electoral systems: <https://mpira.ub.uni-muenchen.de/82513/>
- The policy distance: <https://www.wolframcloud.com/objects/thomas-cool/Voting/2018-02-02-PolicyDistance.nb>
- Aitchison 134 pages: http://ima.udg.edu/activitats/codawork05/A_concise_guide_to_compositional_data_analysis.pdf
- Sheets: <https://laboratoriomatematicas.uniandes.edu.co/cursocoda/04-Vera-geometry.pdf>
- Nutshell: <http://www.sediment.uni-goettingen.de/staff/tolosana/extra/CoDaNutshell.pdf>
- Summary text on closure: http://slon.diamo.cz/hpvt/2011/_Mat/M%2002.pdf
- Different measures (including angular): http://ima.udg.edu/~barcelo/index_archivos/Measures_of_difference_Clustering.pdf
- Use of the geometric mean in the Aitchison improduct: <http://kma.upol.cz/data/xinha/ULOZISTE/ActaMath/0708.pdf>
- Website: <http://www.compositionaldata.com/>
- More involved math of Aitchison's approach to compositional data: <http://dx.doi.org/10.17713/ajs.v45i4.142>
- Wikipedia, a portal and no source: https://en.wikipedia.org/wiki/Compositional_data
- https://en.wikibooks.org/wiki/Why,_and_How,_Should_Geologists_Use_Compositional_Data_Analysis
- Voting but no use of distance: <https://gking.harvard.edu/files/gking/files/multiparty.pdf>
- PM. Aitchison & Brown (1957), The lognormal distribution, CUP

References

Colignatus is the name in science of Thomas Cool, econometrician and teacher of mathematics, Scheveningen, Holland.

This paper is also available at: <https://www.wolframcloud.com/objects/thomas-cool/Voting/2018-01-18-Aitchison.nb> (updated but still the same name).

Aitchison, J., and J.A.C. Brown (1957), "The lognormal distribution", CUP

Aitchison, J. (undated, perhaps latest version 2002), "A concise guide to compositional data analysis", http://ima.udg.edu/activitats/codawork05/A_concise_guide_to_compositional_data_analysis.pdf

Barcelo-Vidal, C., and J.A. Martin-Fernandez (2016), "The mathematics of compositional analysis", Austrian J. of Statistics, Vol 45, No 4, <http://dx.doi.org/10.17713/ajs.v45i4.142>

Colignatus, Th. (2017a), "Comparing votes and seats with cosine, sine and sign, with attention for the slope and enhanced sensitivity to disproportionality", <https://mpira.ub.uni->

muenchen.de/81389/

Colignatus, Th. (2017b), “One woman, one vote. Though not in the USA, UK and France”, <https://mpa-ub.uni-muenchen.de/82513/>

Colignatus, Th. (2017c), “Statistics, slope, cosine, sine, sign, significance and R-squared”, <https://boycottholland.wordpress.com/2017/10/21/statistics-slope-cosine-sine-sign-significance-and-r-squared/>

Colignatus, Th. (2017d), “Karl Pearson’s curious construction of spurious correlation”, <https://boycottholland.wordpress.com/2017/11/19/karl-pearsons-curious-construction-of-spurious-correlation/>

Colignatus, Th. (2018), “Measures of policy distance and inequality / disproportionality of votes and seats”, MPRA_paper_84324.pdf, <https://www.wolframcloud.com/objects/thomas-cool/Voting/2018-02-02-PolicyDistance.nb>

Dongen, S. van & A.J. Enright (2012), “Metric distances derived from cosine similarity and Pearson and Spearman correlations”, <https://arxiv.org/abs/1208.3145>

Golder, M. & Stramski, J. (2010), “Ideological congruence and electoral institutions”, *American Journal of Political Science*, 54(1), 90-106

Hron, K. (2008), “Remark on properties of bases for additive logratio transformations of compositional data”, *Acta Univ. Palacki. Olomuc. Fac. rer. nat., Mathematica* 47, 77-82, <http://kma.upol.cz/data/xinha/ULOZISTE/ActaMath/0708.pdf>

Katz, J.N, and G. King (1999), “A statistical model for multiparty electoral data”, *American political science review*, Vol 93, No 1, p15-32, <https://gking.harvard.edu/files/gking/files/multiparty.pdf>

Maligranda, L. (2006), “Simple norm inequalities”, *Amer. Math. Monthly* 113 (3), 256 – 260

Martin-Fernandez, J.A., and C. Barcelo-Vidal, V. Pawlowsky-Glahn (undated), “Measures of difference for compositional data and hierarchical clustering methods”, http://ima.udg.edu/~barcelo/index_archivos/Measures_of_difference__Clustering.pdf

Pawlowsky-Glahn, V., J.J. Egozcue, and R. Meziat (2007), “The statistical analysis of compositional data: The Aitchison geometry”, (sheets), <https://laboratoriomatematicas.uniandes.edu.co/cursocoda/04-Vera-geometry.pdf>

Pawlowsky-Glahn, V., and J.J. Egozcue (2011), “The closure problem: one hundred years of debate”, http://slon.diamo.cz/hpvt/2011/_Mat/M%2002.pdf

Tolosana, R. (2008), “Compositional Data Analysis in a Nutshell”, <http://www.sediment.uni-goettingen.de/staff/tolosana/extra/CoDaNutshell.pdf>

Vallis, R.A., H. Nunez, and J. C. Martin (2008), “Why, and How, Should Geologists Use Compositional Data Analysis”, https://en.wikibooks.org/wiki/Why,_and_How,_Should_Geologists_Use_Compositional_Data_Analysis

Wang, H., Q. Liu, H.M.K. Mok, L. Fu, W. Man Tse (2007), “A hyperspherical transformation forecasting model for compositional data”, *Eur. J. of Oper. Res.* 179, 459-468, <https://pdfs.semanticscholar.org/67e5/fa4a7cc9c833109722eed64fc3611c4e15f4.pdf>