



Munich Personal RePEc Archive

## **Weighted Shapley hierarchy levels values**

Besner, Manfred

HFT Stuttgart, University of Applied Sciences, Department of  
Geomatics, Computer Science and Mathematics

24 July 2018

Online at <https://mpra.ub.uni-muenchen.de/88160/>  
MPRA Paper No. 88160, posted 24 Jul 2018 11:32 UTC

# Weighted Shapley hierarchy levels values

Manfred Besner\*

July 24, 2018

## Abstract

In this paper we present a new class of values for cooperative games with level structure. We use a multi-step proceeding, suggested first in [Owen \(1977\)](#), applied to the weighted Shapley values. Our first axiomatization is an generalisation of the axiomatization given in [Gómez-Rúa and Vidal-Puga \(2011\)](#), itselfs an extension of a special case of an axiomatization given in [Myerson \(1980\)](#) and [Hart and Mas-Colell \(1989\)](#) respectively by efficiency and weighted balanced contributions. The second axiomatization is completely new and extends the axiomatization of the weighted Shapley values introduced in [Hart and Mas-Colell \(1989\)](#) by weighted standardness for two player games and consistency. As a corollary we obtain a new axiomatization of the Shapley levels value.

**Keywords** Cooperative game · Consistency · Level structure · (Weighted) Shapley (levels) value · Weighted balanced contributions

## 1 Introduction

Presenting his famous value for coalition structures, [Owen \(1977\)](#) suggested an extension of his value that

”... deals with union structure hierarchies, i. e., the possibility that, inside each union, there may be some groups who are closer together then the remaining members of the union. For example, some of the union members may belong to a certain clan, and will therefore make a common front against the other union members.”

[Winter \(1989\)](#) was the first one who putted this idea into action by constructing a model for hierarchical structures, called level structure. Then he defined his value, we call it Shapley levels value, by a set of axioms, which are extensions of axioms of an axiomatization of the Owen value ([Owen, 1977](#)). Another axiomatization of the Shapley levels value

---

\*M. Besner

Department of Geomatics, Computer Science and Mathematics,  
HFT Stuttgart, University of Applied Sciences, Schellingstr. 24  
D-70174 Stuttgart, Germany  
Tel.: +49 (0)711 8926 2791  
E-mail: manfred.besner@hft-stuttgart.de

is presented in [Calvo, Lasaga and Winter \(1996, Theorem 2\)](#). They extended there the axiomatization of the Shapley value in [Myerson \(1980\)](#) and [Hart and Mas-Colell \(1989\)](#) respectively by efficiency and balanced contributions to level structures to determine the Shapley levels value uniquely.

This proceeding in mind [Vidal-Puga \(2012\)](#) defined initially a weighted value for coalition structures and [Gómez-Rúa and Vidal-Puga \(2011\)](#) extended it to level structures. Their value is axiomatized by an extension of an axiomatization of the weighted Shapley values presented in [Myerson \(1980\)](#) and [Hart and Mas-Colell \(1989\)](#) too. So this value is axiomatized by efficiency and an extension of a special case of weighted balanced contributions, called balanced per capita contributions. By this value all coalitions have been assigned weights given by the cardinality of the coalitions. In difference to the Shapley levels value, acting coalitions are only be treated as symmetric if they contain the same amount of players.

But sometimes this approach is not adequate, e.g. if some coalitions, fairly independent from the size of the coalition, have a lack of ability to perform or not the same political power and so on. To be no longer restricted to that limitation we introduce arbitrary exogenously given weights for the coalitions which become prominent if the coalitions act as players. So we can introduce the weighted Shapley hierarchy levels values. The axiomatization of the value from [Gómez-Rúa and Vidal-Puga \(2011\)](#) can be easily transferred to an axiomatization given by efficiency and weighted balanced group contributions, a general extension of weighted balanced contributions in [Myerson \(1980\)](#).

[Hart and Mas-Colell \(1989\)](#) presented another axiomatization of the weighted Shapley values. They used the weighted standardness property, that means that in a two player game the cooperation benefit has to be divided proportional to the weights of the players, and a consistency property. Consistency requires the payout to a player in a reduced game be the same as the payoff to this player in the original game if the reduced game is defined appropriate. As main part of the paper, we extend this axiomatization to level structures that determines the weighted Shapley hierarchy levels values uniquely.

The plan of the paper is as follows. Some preliminaries are given in section 2, in section 3 are introduced the weighted Shapley hierarchy levels values and an axiomatization in the sense of [Calvo, Lasaga and Winter \(1996, Theorem 2\)](#), section 4 presents a new consistency property with an related axiomatization and section 5 gives a short conclusion.

## 2 Preliminaries

In some definitions and notations we will follow with [Besner \(2018a\)](#). Let  $\mathbb{R}$  be the real numbers,  $\mathbb{R}_{++}$  the set of all positive real numbers,  $\mathfrak{U}$  a countably infinite set, the universe of all possible players, and  $\mathcal{N}$  the set of all finite subsets  $N \subseteq \mathfrak{U}$ ,  $N \neq \emptyset$ . A **TU-game** is a pair  $(N, v)$  where  $N \in \mathcal{N}$  and  $v: 2^N \rightarrow \mathbb{R}$ ,  $v(\emptyset) = 0$ ,  $2^N := \{S: S \subseteq N\}$ , is called a **coalition function**. We call  $v(S)$  the **worth** of **coalition**  $S \subseteq N$ , we denote by  $\Omega^S$  the set of all nonempty subsets of  $S$  and the set of all TU-games with player set  $N$  is denoted by  $\mathbb{V}^N$ .  $(S, v)$  is the **restriction** of  $(N, v)$  to  $S \in \Omega^N$  and a player  $i \in N$  is called a **null player** in  $v$  if  $v(S \cup \{i\}) = v(S)$ ,  $S \subseteq N \setminus \{i\}$ .

$\mathcal{B} := \{B_1, \dots, B_m\}$  is said to be a **coalition structure** on  $N$  if  $B_k \cap B_\ell = \emptyset$ ,  $1 \leq k < \ell \leq m$ ,  $B_k \neq \emptyset$  for all  $k$ ,  $1 \leq k \leq m$ , and  $\bigcup_{k=1}^m B_k = N$ . We call each  $B \in \mathcal{B}$  a **component**

and  $\mathcal{B}(i)$  denotes the component that contains the player  $i \in N$ .

A finite sequence  $\underline{\mathcal{B}} := \{\mathcal{B}^0, \dots, \mathcal{B}^{h+1}\}$  of coalition structures  $\mathcal{B}^r$ ,  $0 \leq r \leq h+1$ , on  $N$  is called a **level structure** (Winter, 1989) on  $N$  if  $\mathcal{B}^r$  is a refinement of  $\mathcal{B}^{r+1}$  for each  $r$ ,  $0 \leq r \leq h$ , i.e.  $\mathcal{B}^r(i) \subseteq \mathcal{B}^{r+1}(i)$  for all  $i \in N$ ,  $\mathcal{B}^0 = \{\{i\} : i \in N\}$  and  $\mathcal{B}^{h+1} = \{N\}$ .  $\mathcal{B}^r$  denotes the  $r$ -th **level** of  $\underline{\mathcal{B}}$  and  $\mathcal{B}^r(B^k)$  denotes the component of the  $r$ -th level that contains the component  $B^k \in \mathcal{B}^k$ ,  $0 \leq k \leq r \leq h+1$ . The set of all level structures with player set  $N$  is denoted by  $\mathbb{L}^N$ . A triple  $(N, v, \underline{\mathcal{B}})$  consisting of a TU-game  $(N, v) \in \mathbb{V}^N$  and a level structure  $\underline{\mathcal{B}} \in \mathbb{L}^N$  is called an **LS-game**. The collection of all LS-games on  $N$  is denoted by  $\mathbb{VL}^N$ . Each  $(N, v, \underline{\mathcal{B}}_0)$  with a **trivial level structure**  $\underline{\mathcal{B}}_0 := \{\mathcal{B}^0, \mathcal{B}^1\}$  corresponds to a TU-game  $(N, v)$  and each LS-game  $(N, v, \underline{\mathcal{B}}_1)$ ,  $\underline{\mathcal{B}}_1 := \{\mathcal{B}^0, \mathcal{B}^1, \mathcal{B}^2\}$ , corresponds to a game with coalition structure as introduced in Aumann and Drèze (1974), in (Owen (1977) called "games with a priori unions".

Let  $(N, v, \underline{\mathcal{B}}) \in \mathbb{VL}^N$ ,  $\underline{\mathcal{B}} = \{\mathcal{B}^0, \dots, \mathcal{B}^{h+1}\}$ , and  $T \in \Omega^N$ :

- From a level structure on  $N$  follows a **restricted** level structure on  $T$  by eliminating the players in  $N \setminus T$ . With coalition structures  $\mathcal{B}^r|_T := \{B \cap T : B \in \mathcal{B}^r, B \cap T \neq \emptyset\}$ ,  $0 \leq r \leq h+1$ , the new level structure on  $T$  is given by  $\underline{\mathcal{B}}|_T := \{\mathcal{B}^0|_T, \dots, \mathcal{B}^{h+1}|_T\} \in \mathbb{L}^T$  and  $(T, v, \underline{\mathcal{B}}|_T) \in \mathbb{VL}^T$  is called the **restriction** of  $(N, v, \underline{\mathcal{B}})$  to player set  $T$ .
- We define  $\underline{\mathcal{B}}^r := \{\mathcal{B}^{r^0}, \dots, \mathcal{B}^{r^{h+1-r}}\} \in \mathcal{L}^{\mathcal{B}^r}$ ,  $0 \leq r \leq h$ , as the **induced  $r$ -th level structure** from  $\underline{\mathcal{B}}$  by considering the components  $B \in \mathcal{B}^r$  as players, where  $\mathcal{B}^{r^k} := \{\{B \in \mathcal{B}^r : B \subseteq B'\} : B' \in \mathcal{B}^{r+k}\}$ ,  $0 \leq k \leq h+1-r$ . The **induced  $r$ -th level game**  $(\mathcal{B}^r, v^r, \underline{\mathcal{B}}^r) \in \mathcal{GL}^{\mathcal{B}^r}$  is given by

$$v^r(Q) := v\left(\bigcup_{B \in Q} B\right) \text{ for all } Q \subseteq \mathcal{B}^r.^1 \quad (1)$$

- We define  $\underline{\mathcal{B}}_r := \{\mathcal{B}^0, \dots, \mathcal{B}^r, \{N\}\} \in \mathcal{L}^N$ ,  $0 \leq r \leq h$ , as the  **$r$ -th cut level structure** from  $\underline{\mathcal{B}}$  if we cut out all levels between the  $r$ -th and the  $(h+1)$ -th level.  $(N, v, \underline{\mathcal{B}}_r)$  is called the  **$r$ -th cut** of  $(N, v, \underline{\mathcal{B}})$ . Note that we have for each  $\underline{\mathcal{B}} = \{\mathcal{B}^0, \dots, \mathcal{B}^{h+1}\}$  also  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_h$ .

A **TU-value**  $\phi$  is an operator that assigns a payoff vector  $\phi(N, v) \in \mathbb{R}^N$  to any  $(N, v) \in \mathbb{V}^N$ , an **LS-value**  $\varphi$  is an operator that assigns a payoff vector  $\varphi(N, v, \underline{\mathcal{B}}) \in \mathbb{R}^N$  to any  $(N, v, \underline{\mathcal{B}}) \in \mathbb{VL}^N$ .

We define  $W := \{f : \mathcal{U} \rightarrow \mathbb{R}_{++}\}$  with  $w_i := w(i)$  for all  $w \in W$  and  $i \in \mathcal{U}$  as the set of all positive weight systems on the universe of all players and we define  $\mathcal{W}^{2^N} := \{f : 2^N \setminus \emptyset \rightarrow \mathbb{R}_{++}\}$  with  $w_S := w(S)$  for all  $w \in \mathcal{W}^{2^N}$ , as the set of all positive weight systems on all coalitions  $S \in \Omega^N$ .

The (simply) **weighted Shapley values**<sup>2</sup>  $Sh^w$  (Shapley, 1953a) are defined by

$$Sh_i^w(N, v) := \sum_{S \subseteq N, S \ni i} \frac{w_i}{\sum_{j \in S} w_j} \Delta_v(S) \text{ for all } i \in N \text{ and } w \in W. \quad (2)$$

The **Shapley value**  $Sh$  (Shapley, 1953b) that distributes the dividends  $\Delta_v(S)$  equally among all players of a coalition  $S$  is a special case of a weighted Shapley value, given by

$$Sh_i(N, v) := \sum_{S \subseteq N, S \ni i} \frac{\Delta_v(S)}{|S|} \text{ for all } i \in N.$$

<sup>1</sup>In Owen (1977), where we have the special case  $r = 1$ , such a game is called as quotient game.

<sup>2</sup>We desist from possibly null weights as in Shapley (1953a) or Kalai and Samet (1987).

Let  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$ ,  $\mathcal{B} = \mathcal{B}_h$ ,  $T \in \Omega^N$ ,  $T \ni i$ , and

$$K_T(i) := \prod_{r=0}^h K_T^r(i), \quad \text{where } K_T^r(i) := \frac{1}{|\{B \in \mathcal{B}^r : B \subseteq \mathcal{B}^{r+1}(i), B \cap T \neq \emptyset\}|}.$$

The **Shapley Levels value**<sup>3</sup>  $Sh^L$  (Winter, 1989) is given by

$$Sh_i^L(N, v, \underline{\mathcal{B}}) := \sum_{T \subseteq N, T \ni i} K_T(i) \Delta_v(T) \text{ for all } i \in N.$$

Obviously coincides  $Sh^L$  with  $Sh$  if  $h = 0$ . It is well-known that  $Sh^L$  coincides with the **Owen value** (Owen, 1977) if  $h = 1$ .

We refer to the following axioms for LS-values which are adaptations of standard-axioms for TU-values (with the exception of the last axiom):<sup>4</sup>

**Efficiency, E.** For all  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$ , we have  $\sum_{i \in N} \varphi_i(N, v, \underline{\mathcal{B}}) = v(N)$ .

**Null player, N.** For all  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$  and  $i \in N$  a null player in  $v$ , we have  $\varphi_i(N, v, \underline{\mathcal{B}}) = 0$ .

**Additivity, A.** For all  $(N, v, \underline{\mathcal{B}}), (N, v', \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$ , we have  $\varphi(N, v, \underline{\mathcal{B}}) + \varphi(N, v', \underline{\mathcal{B}}) = \varphi(N, v + v', \underline{\mathcal{B}})$ .

**Balanced group contributions, BGC** (Calvo, Lasaga and Winter, 1996). For all  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$ ,  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_h$ ,  $B_k, B_\ell \in \mathcal{B}^r$ ,  $0 \leq r \leq h$ , such that  $B_\ell \subseteq \mathcal{B}^{r+1}(B_k)$ , we have

$$\sum_{i \in B_k} \varphi_i(N, v, \underline{\mathcal{B}}) - \sum_{i \in B_k} \varphi_i(N \setminus B_\ell, v, \underline{\mathcal{B}}|_{N \setminus B_\ell}) = \sum_{i \in B_\ell} \varphi_i(N, v, \underline{\mathcal{B}}) - \sum_{i \in B_\ell} \varphi_i(N \setminus B_k, v, \underline{\mathcal{B}}|_{N \setminus B_k}).$$

This axiom states that for two components  $B_k, B_\ell$  of the same level which belong to the same component one level higher the contribution of  $B_\ell$  to the total payoff of all players from  $B_k$  equals the contribution of  $B_k$  to the total payoff of all players from  $B_\ell$ .

**Standardness, ST<sup>0</sup>.** (Myerson, 1980) For all  $(N, v) \in \mathbb{V}^N$ ,  $N = \{i, j\}$ ,  $i \neq j$ , we have

$$\varphi_i(N, v) = v(\{i\}) + \frac{1}{2} [v(\{i, j\}) - v(\{i\}) - v(\{j\})].$$

Standardness leads to sharing the surplus by cooperating in two player games that the surplus is divided equally. The following axiom stands for dividing the surplus in two player games proportional to the weights of the players.

**Weighted standardness, WS<sup>0</sup>.** (Myerson, 1980) For all  $(N, v) \in \mathbb{V}^N$ ,  $N = \{i, j\}$ ,  $i \neq j$ , and  $w \in W$ , we have

$$\varphi_i(N, v) = v(\{i\}) + \frac{w_i}{w_i + w_j} [v(\{i, j\}) - v(\{i\}) - v(\{j\})].$$

The next axiom has the meaning that the sum of the payoff to all players in a component coincides with the payoff to this component if this component is regarded as a player in an induced level game.

**Level game property, LG** (Winter, 1989). For all  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$ ,  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_h$ ,  $B \in \mathcal{B}^r$ ,  $0 \leq r \leq h + 1$ , we have

$$\sum_{i \in B} \varphi_i(N, v, \underline{\mathcal{B}}) = \varphi_B(\mathcal{B}^r, v^r, \underline{\mathcal{B}}^r).$$

<sup>3</sup>This formula is presented in Calvo, Lasaga and Winter (1996, eq. (1))

<sup>4</sup>In the case of using a subdomain, we require an axiom to hold when all games belong to this subdomain. If there are used weights for some coalitions, these weights are still valid in the subdomain.

### 3 Weighted Shapley hierarchy levels values

Owen (1977) suggested for his value a two-step approach where the first step consists of using the Shapley value on the quotient game that corresponds to a first induced level game. In the second stage the Shapley value is applied to the result of the first step again. The same procedure adapted Vidal-Puga (2012) for the value  $\zeta$ , an extension of a weighted Shapley value to coalition structures where the weights are given by the size of the coalitions. Gómez-Rúa and Vidal-Puga (2011) presented an extension to level structures and replaced the two-step approach by a multi-step mechanism as suggested also in Owen (1977). It is still not known and there is no hint in Gómez-Rúa and Vidal-Puga (2011) that their value can easily be generalized to an extension of the weighted Shapley values in general if someone knows how to deal with arbitrary exogenously given weights for coalitions. The substantial idea behind our definition, similar to the one suggested in Owen (1977) and implemented as an algorithm in Gómez-Rúa and Vidal-Puga (2011) is as follows:

In a first step we distribute the worth of the grand coalition  $v(N)$  among the components  $B^h$  of the  $h$ -th level as players using a weighted Shapley value. In the second step each payoff to a component  $B^h$  from the first step is splitted up by the weighted Shapley value to all components  $B^{h-1}$  (the new players) which are subsets of the component  $B^h$  and so on for all levels. In the last step we distribute the payoff to components  $B^1$  from the first level among the original players  $i \in N$ . Therefore we need a weight system for coalitions to have weights if a coalition is regarded as a player. For notational parsimony we give the following notation.

**Notation 3.1.** Let  $(N, v, \underline{\mathcal{B}}) \in \mathbb{VL}^N$ ,  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_h$ ,  $i \in N$  and  $T \in \Omega^{\mathcal{B}^k(i)}$ ,  $0 \leq k \leq h$ . We denote by  $\mathcal{T}_i^k := \{B \in \mathcal{B}^k : B \subseteq \mathcal{B}^{k+1}(i), \bar{B} \neq \mathcal{B}^k(i)\} \cup \{T\}$  the set that contains all components of the  $k$ -th level which are subsets of the component of the  $(k+1)$ -th level that contains the player  $i$  without the component of the  $k$ -th level containing player  $i$  and that contains in addition the coalition  $T$ .

**Definition 3.2.** Let  $(N, v, \underline{\mathcal{B}}) \in \mathbb{VL}^N$ ,  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_h$ ,  $w \in \mathcal{W}^{2^N}$ ,  $i \in N$  and let for all  $k$ ,  $0 \leq k \leq h$ ,  $T \in \Omega^{\mathcal{B}^k(i)}$ ,  $\mathcal{T}_i^k$  the set from notation 3.1,  $w^{k,T} \in W$  a weight system with coalitions  $S$  as players such that  $w_S^{k,T} := w_S$ ,  $w_S \in w$ , for all  $S \in \mathcal{T}_i^k$ ,  $\bar{v}_i^{h+1} := v$  and  $\bar{v}_i^k$  given by

$$\bar{v}_i^k(T) := Sh_T^{w^{k,T}}(\mathcal{T}_i^k, \bar{v}_i^k) \text{ for all } T \in \Omega^{\mathcal{B}^k(i)} \quad (3)$$

where  $\bar{v}_i^k$  is specified by

$$\bar{v}_i^k(Q) := \bar{v}_i^{k+1}\left(\bigcup_{S \in Q} S\right) \text{ for all } Q \subseteq \mathcal{T}_i^k.$$

Then the **weighted Shapley hierarchy levels value**  $Sh^{wHL}$  is defined by

$$Sh_i^{wHL}(N, v, \underline{\mathcal{B}}) := \bar{v}_i^0(\{i\}) \text{ for all } i \in N.$$

**Remarks 3.3.**  $Sh^{wHL}$  coincides with  $Sh^w$  if  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_0$  and  $w_{\{i\}} = w_i$  for all  $i \in N$ . Since  $Sh^w$  is efficient and additive, it is obvious, that  $Sh^{wHL}$  meets **E**, **A** and **LG**.

Similar to Calvo, Lasaga and Winter (1996), which characterized the Shapley levels value by efficiency and balanced group contributions, Gómez-Rúa and Vidal-Puga (2011) characterized their value by efficiency and balanced per capita contributions, a special case of the following axiom where the weights are the size of the relevant components.

**Weighted balanced group contributions, WBGC<sup>5</sup>.** For all  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$ ,  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_h$ ,  $w \in \mathcal{W}^{2^N}$ ,  $B_k, B_\ell \in \mathcal{B}^r$ ,  $0 \leq r \leq h$ , such that  $B_\ell \subseteq \mathcal{B}^{r+1}(B_k)$ , we have

$$\frac{\sum_{i \in B_k} \varphi_i(N, v, \underline{\mathcal{B}}) - \sum_{i \in B_k} \varphi_i(N \setminus B_\ell, v, \underline{\mathcal{B}}|_{N \setminus B_\ell})}{w_{B_k}} = \frac{\sum_{i \in B_\ell} \varphi_i(N, v, \underline{\mathcal{B}}) - \sum_{i \in B_\ell} \varphi_i(N \setminus B_k, v, \underline{\mathcal{B}}|_{N \setminus B_k})}{w_{B_\ell}}.$$

In difference to **BGC**, here the contributions of one component to the players of the other component and vice versa are proportional to the weights of the components.

**Theorem 3.4.** *Let  $w \in \mathcal{W}^{2^N}$ .  $Sh^{wHL}$  is the unique LS-value that satisfies **E** and **WBGC**.*

*Proof.* It is clear that def. 3.2 is equivalent to the algorithm given in Gómez-Rúa and Vidal-Puga (2011, section 3) if we use as weights the cardinality of the coalitions. Thus the rest of the proof of theorem 3.4 is omitted because it's only an exercise to adapt the proofs of proposition 7 and theorem 8 in Gómez-Rúa and Vidal-Puga (2011), using weighted balanced contributions (Myerson, 1980; Hart and Mas-Colell, 1989) for the weighted Shapley values in proposition 7 and replacing the given weights (size of the components) by arbitrary weights in theorem 8.  $\square$

**Remark 3.5.** *If all weights are equal, we use in fact in line (3) the Shapley value and **WBGC** equals **BGC**. So, by Calvo, Lasaga and Winter (1996, Theorem 2), def. 3.2 determines in this case the Shapley levels value  $Sh^L$ .*

**Remark 3.6.** *As shown by example 1 in Gómez-Rúa and Vidal-Puga (2011),  $Sh^{wHL}$  doesn't satisfy **N** in general.*

## 4 Level-consistency

Hart and Mas-Colell (1989) used a consistency property to characterize the weighted Shapley values. Therefore they introduced a reduced game of a TU-game.

**Definition 4.1.** *Let  $(N, v) \in \mathbb{V}^N$ ,  $R \in \Omega^N$  and  $\phi$  a TU-value. The **reduced game**  $(R, v_R^\phi) \in \mathbb{V}^R$  is given for all  $S \in \Omega^R$  by*

$$v_R^\phi(S) := v(S \cup R^c) - \sum_{j \in R^c} \phi_j(S \cup R^c, v) \quad (4)$$

where  $R^c := N \setminus R$ .

**Remark 4.2.** *If  $\phi$  is efficient eq. (4) can be simplified to*

$$v_R^\phi(S) := \sum_{j \in S} \phi_j(S \cup R^c, v).$$

The interpretation of a reduced game must be seen in context with a TU-value that must have a special property. So Hart and Mas-Colell (1989) formulated the following axiom.

**Consistency, C.** For all  $(N, v) \in \mathbb{V}^N$ ,  $R \subseteq N$ , we have  $\phi_i(N, v) = \phi_i(R, v_R^\phi)$  for all  $i \in R$ .

<sup>5</sup>This axiom extends weighted balanced contributions (Myerson, 1980).



The meaning of this axiom is as follows: A coalition of players  $R^c$  leaves the game. In the reduced game each coalition  $S$  which is a subset of the coalition  $R$  of the remaining players obtains the worth of the old coalition  $S \cup R^c$  less the payoff that the removed players obtain in a game played on  $S \cup R^c$ . Then a TU-value is consistent if the players in  $R$  obtain in the reduced and in the original game the same payoff. It turned out that the weighted Shapley values have got an intense connection to this axiom.

**Theorem 4.3.** (*Hart and Mas-Colell, 1989*) Let  $w \in W$ .  $Sh^w$  is the unique TU-value that satisfies **C** and **WS**<sup>0</sup>.

If we exchange **WS**<sup>0</sup> by **ST**<sup>0</sup> this theorem characterizes the Shapley value. By adding **LG**, Winter (1992) extended this axiomatization for the Shapley value to the Owen value (see proof of theorem 3 in Winter (1992)). Similar Huettner (2015) transferred the related axiomatization of Feldman (1999) and Ortmann (1999) for the proportional value to his proportional value for coalition structures. Our new reduced game for level structures extends the reduced game for TU-games in Hart and Mas-Colell (1989) too but it is no extension of the reduced game used in Winter (1992) and Huettner (2015) respectively.

**Definition 4.4.** Let  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$ ,  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_h$ ,  $R \in \Omega^N$  such that  $R = \bigcup_{Q \in \mathcal{B}^h} Q$  and  $\varphi$  an LS-value. The **reduced level game**  $(R, v_R^\varphi, \underline{\mathcal{B}}|_R) \in \mathbb{V}\mathbb{L}^R$  is given for all  $S \in \Omega^R$  by

$$v_R^\varphi(S) := v(S \cup R^c) - \sum_{Q \in \mathcal{B}^h, Q \not\subseteq R} \varphi_Q(\mathcal{B}^h|_{S \cup R^c}, v^h, \underline{\mathcal{B}}^h|_{S \cup R^c}) \quad (5)$$

where  $R^c := N \setminus R$  and  $(\mathcal{B}^h|_{S \cup R^c}, v^h, \underline{\mathcal{B}}^h|_{S \cup R^c})$  is the induced  $h$ -th level game restricted to  $S \cup R^c$ .

**Remark 4.5.** If  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_0$ , def. 4.4 coincides with def. 4.1.

**Remark 4.6.** If  $\varphi$  is efficient eq. (5) can be simplified to

$$v_R^\varphi(S) := \sum_{Q \in \mathcal{B}^h, Q \subseteq R} \varphi_{S \cap Q}(\mathcal{B}^h|_{S \cup R^c}, v^h, \underline{\mathcal{B}}^h|_{S \cup R^c}).$$

The following axiom extends **C**.

**Level-consistency, LC.** For all  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$ ,  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_h$ ,  $R = \bigcup_{Q \in \mathcal{B}^h} Q$ , we have

$$\varphi_i(N, v, \underline{\mathcal{B}}) = \varphi_i(R, v_R^\varphi, \underline{\mathcal{B}}|_R) \text{ for all } i \in R. \quad (6)$$

$R$  is a coalition of all players of some components of the  $h$ -th level. All players of coalition  $R^c$  of the players of the other components of the  $h$ -th level leave the game. In the reduced level game each coalition  $S \in \Omega^R$  obtains the worth of the old coalition  $S \cup R^c$  less the payoff that the removed components obtain in the induced  $h$ -th level game restricted to  $S \cup R^c$ . Then an LS-value is consistent if each player in  $R$  obtain in the reduced game the same payoff as in the original game.

**Remark 4.7.** If  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_0$ , **LC** coincides with **C**.

We adapt **WS**<sup>0</sup> to level structures.

**Weighted standardness, WS.** For all  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$  with  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_0$ ,  $N = \{i, j\}$ ,  $i \neq j$ , and  $w \in \mathcal{W}^{2^N}$ , we have

$$\varphi_i(N, v, \underline{\mathcal{B}}) = v(\{i\}) + \frac{w_{\{i\}}}{w_{\{i\}} + w_{\{j\}}} [v(\{i, j\}) - v(\{i\}) - v(\{j\})].$$



To characterize the weighted Shapley hierarchy levels values finally we need a very weak property that states that if the  $h$ -th level consists only of one component we can remove this level from the game without consequences.

**Inessential level property, IL.** For all  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$ ,  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_h$ ,  $h \geq 1$ , and  $\mathcal{B}^h = \mathcal{B}^{h+1}$  we have  $\varphi(N, v, \underline{\mathcal{B}}) = \varphi(N, v, \underline{\mathcal{B}}_{h-1})$ .

The last three properties fit best with the weighted Shapley hierarchy levels values:

**Theorem 4.8.** *Let  $w \in \mathcal{W}^{2^N}$ .  $Sh^{wHL}$  satisfies **IL**, **WS** and **LC**.*

*Proof.* Let  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$ ,  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_h$  and  $w \in \mathcal{W}^{2^N}$ :

- **IL:** If  $\mathcal{B}^h = \mathcal{B}^{h+1}$  we have in notation 3.1 for all  $i \in N$  and  $T \in \Omega^{\mathcal{B}^{h(i)}}$ ,  $\mathcal{T}_i^h = \{T\}$ . It follows in line (3),  $\bar{v}_i^h(T) = Sh_T^{w^{h,T}}(\{T\}, \tilde{v}_i^h) = v(T)$  for all  $T \in \Omega^{\mathcal{B}^{h(i)}}$ . Thus, if  $h \geq 1$ , we have no change in line (3) for all  $k$ ,  $0 \leq k \leq h-1$ , if we drop the  $h$ -th level and **IL** is shown.
- **WS:** By def. (3.2) and line (2) the claim is obvious.
- **LC:** Let  $i \in N$ ,  $R = \bigcup_{Q \in \mathcal{B}^h} Q$ ,  $R \supseteq \mathcal{B}^{h(i)}$ . By **IL** it is sufficient to show **LC** on level structures  $\underline{\mathcal{B}}_h$  with  $\mathcal{B}^h \neq \mathcal{B}^{h+1}$ . If  $R = N$  or  $|N| = 1$ , eq. (6) is trivially satisfied. Let now  $|N| \geq 2$  and  $R \neq N$ . By Hart and Mas-Colell (1989),  $Sh^w$  satisfies **C** for all  $w \in \mathcal{W}$ . For each  $T \in \Omega^{\mathcal{B}^{h(i)}}$  we have a set  $\mathcal{T}_i^h$  from notation 3.1 and a related  $\mathcal{T}_i^h|_R$  on the restricted level structure  $\underline{\mathcal{B}}|_R$ . Thus follows by eq. (3) for  $k = h$

$$\bar{v}_i^h(T) = Sh_T^{w^{h,T}}(\mathcal{T}_i^h, \tilde{v}_i^h) \stackrel{\text{rem. 4.5}}{=} Sh_T^{w^{h,T}}(\mathcal{T}_i^h|_R, (\widetilde{v}_R^\varphi)_i^h) \quad (7)$$

where  $(\widetilde{v}_R^\varphi)_i^h$  is given by  $(\widetilde{v}_R^\varphi)_i^h(\mathcal{Q}) := v_R^\varphi(\bigcup_{S \in \mathcal{Q}} S)$  for all  $\mathcal{Q} \subseteq \mathcal{T}_i^h|_R$ . Thus there is no difference in eq. (3) between both games for all  $k \neq h$  too and **LC** is shown.  $\square$

It transpires that the weighted Shapley hierarchy levels values are characterized by the following theorem.

**Theorem 4.9.** *Let  $w \in \mathcal{W}^{2^N}$ .  $Sh^{wHL}$  is the unique TU-value that satisfies **IL**, **WS** and **LC**.*

*Proof.* By theorem 4.8 it is sufficient to show uniqueness. Let  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$ ,  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_h$ ,  $w \in \mathcal{W}^{2^N}$ ,  $i \in N$  arbitrary and  $\varphi$  an LS-values that satisfies **IL**, **WS** and **LC**. By **LC**, we have

$$\varphi_j(N, v, \underline{\mathcal{B}}) = \varphi_j(\mathcal{B}^{h(i)}, v_{\mathcal{B}^{h(i)}}^\varphi, \underline{\mathcal{B}}|_{\mathcal{B}^{h(i)}}) \text{ for all } j \in \mathcal{B}^{h(i)}$$

where  $v_{\mathcal{B}^{h(i)}}^\varphi$  is unique by **WS**, **LC**, rem. 4.7, rem. 4.5 and theorem 4.3.

By **IL**, we have for each restricted  $k$ -th cut level structure  $\underline{\mathcal{B}}_k|_{\mathcal{B}^{k(i)}}$ ,  $k > 0$ , from  $\underline{\mathcal{B}}$  on  $\mathcal{B}^k(i)$ ,

$$\varphi(\mathcal{B}^k(i), \hat{v}_{\mathcal{B}^k(i)}^\varphi, \underline{\mathcal{B}}_k|_{\mathcal{B}^k(i)}) = \varphi(\mathcal{B}^k(i), \hat{v}_{\mathcal{B}^k(i)}^\varphi, \underline{\mathcal{B}}_{k-1}|_{\mathcal{B}^k(i)})$$

where  $\hat{v}_{\mathcal{B}^k(i)}^\varphi$  is recursive defined by

$$\hat{v}_{\mathcal{B}^k(i)}^\varphi := \begin{cases} v_{\mathcal{B}^k(i)}^\varphi & \text{if } k = h, \\ (\hat{v}_{\mathcal{B}^{k+1}(i)}^\varphi)_{\mathcal{B}^k(i)}^\varphi & \text{else.} \end{cases}$$

Hence by an induction on the levels  $k$ , **IL** and **LC** it follows that

$$\varphi_j(N, v, \underline{\mathcal{B}}) = \varphi_j(\mathcal{B}^1(i), \hat{v}_{\mathcal{B}^1(i)}^\varphi, \underline{\mathcal{B}}_0|_{\mathcal{B}^1(i)}) \text{ for all } j \in \mathcal{B}^1(i)$$

where  $\hat{v}_{\mathcal{B}^1(i)}^\varphi$  is unique by **WS**, **LC**, rem. 4.7, rem. 4.5 and theorem 4.3. Therefore, by **WS**, **LC**, rem. 4.7, rem. 4.5 and theorem 4.3,  $\varphi$  is well defined and the proof is completed.  $\square$

To introduce a new axiomatization of the Shapley levels value we adapt **ST**<sup>0</sup>.

**Standardness, ST.** For all  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$  with  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_0$  and  $N = \{i, j\}, i \neq j$ , we have

$$\varphi_i(N, v, \underline{\mathcal{B}}) = v(\{i\}) + \frac{1}{2}[v(\{i, j\}) - v(\{i\}) - v(\{j\})].$$

Since **ST** is a special case of **WS** we obtain the following corollary by theorem 4.9.

**Corollary 4.10.** *Sh<sup>L</sup> is the unique TU-value that satisfies **IL**, **ST** and **LC**.*

**Remark 4.11.** *The axioms in theorem 4.9 are logically independent.*

*Proof.* Let  $(N, v, \underline{\mathcal{B}}) \in \mathbb{V}\mathbb{L}^N$ ,  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_h$  and  $w \in \mathcal{W}^{2^N}$ :

- **IL:** The LS-value  $\varphi$  defined by
 
$$\varphi := \begin{cases} \frac{1}{2}Sh^{wHL}, & \text{if } h \geq 1, \\ Sh^{wHL}, & \text{else,} \end{cases}$$

satisfies **WS** and **LC** but not **IL**.

- **WS:** Let  $v \in \mathbb{V}^N, v(S) > 0$  for all  $S \in \Omega^N$ , and  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_1$ . If we regard the proportional value for cooperative games with a coalition structure in [Huettner \(2015\)](#) in the notation for a level structure with  $\underline{\mathcal{B}} = \underline{\mathcal{B}}_1$  this value satisfies **IL** and **LC** but not **WS**.
- **LC:** The LS-value  $\varphi$  defined by  $\varphi(N, v, \underline{\mathcal{B}}) := Sh^w(N, v)$  with  $w_i := w_{\{i\}}$  for all  $i \in N$  satisfies **IL** and **WS** but not **LC**.  $\square$

## 5 Conclusion

It would seem reasonable in our view to use weights if operating players are in some sense not symmetric, e. g., if some players have not the same political, financial or military power, contribute more to the result and so on. Often players act not as lone fighters. They join forces and small groups form together larger groups and so on. So the weighted Shapley hierarchy levels values meet with the multi-stage mechanism a widespread idea how a hierarchical value has to operate.

That these values don't satisfy the null player property is not necessary a disadvantage. A null player receives a compensation since she supports in an acting coalition the other members with her contribution to the weight of the coalition. So a null player is only a null in the coalition function and not necessary in the weight system for coalitions. For further information on this area see also [Gómez-Rúa and Vidal-Puga \(2010\)](#) and [Gómez-Rúa and Vidal-Puga \(2011\)](#).

An emphasis of this paper is on consistency. Consistency plays an important role in many allocation concepts in cooperative game theory. The differences in the numerous

varieties of the reduced games are decisive for the diversity of the different consistency characteristics. So our reduced game is no extension of the reduced game in Winter (1992). There the players which determine the worth in the coalition function of the reduced game are the original players of the game. In our reduced game determine acting coalitions in a level game the worths of the coalitions. So also this reduced game coincides with the reduced game suggested in Hart and Mas-Colell (1989) if the level structure is trivial since there the players are singletons.

## References

- Aumann, R.J., Drèze, J., 1974. Cooperative games with coalition structures. *International Journal of Game Theory* 3, 217–237.
- Besner, M. (2018a). The weighted Shapley support levels values. Working paper.
- Calvo, E., Lasaga, J. J., & Winter, E. (1996). The principle of balanced contributions and hierarchies of cooperation, *Mathematical Social Sciences*, 31(3), 171–182.
- Feldman, B. (1999). The proportional value of a cooperative game. *Manuscript. Chicago: Scudder Kemper Investments*.
- Gómez-Rúa, M., & Vidal-Puga, J. (2010). The axiomatic approach to three values in games with coalition structure. *European Journal of Operational Research*, 207(2), 795–806.
- Gómez-Rúa, M., & Vidal-Puga, J. (2011). Balanced per capita contributions and level structure of cooperation. *Top*, 19(1), 167–176.
- Hart, S., & Mas-Colell, A. (1989). Potential, value, and consistency. *Econometrica*, 57(3) 589–614.
- Huettner, F. (2015). A proportional value for cooperative games with a coalition structure. *Theory and Decision*, 78(2), 273–287.
- Kalai, E., & Samet, D. (1987). On weighted Shapley values. *International Journal of Game Theory* 16(3), 205–222.
- Myerson, R. B. (1980). Conference Structures and Fair Allocation Rules, *International Journal of Game Theory*, Volume 9, Issue 3, 169–182.
- Ortmann, K. M. (2000). The proportional value for positive cooperative games. *Mathematical Methods of Operations Research*, 51(2), 235–248.
- Owen, G. (1977). Values of games with a priori unions. In *Essays in Mathematical Economics and Game Theory*, Springer, Berlin Heidelberg, 76–88.
- Shapley, L. S. (1953a). Additive and non-additive set functions. *Princeton University*.
- Shapley, L. S. (1953b). A value for n-person games. H. W. Kuhn/A. W. Tucker (eds.), *Contributions to the Theory of Games*, Vol. 2, Princeton University Press, Princeton, 307–317.
- Vidal-Puga, J. (2012). The Harsanyi paradox and the "right to talk" in bargaining among coalitions. *Mathematical Social Sciences*, 64(3), 214–224.
- Winter, E. (1989). A value for cooperative games with levels structure of cooperation. *International Journal of Game Theory*, 18(2), 227–240.
- Winter, E. (1992). The consistency and potential for values of games with coalition structure. *Games and Economic Behavior*, 4(1), 132–144.