

MPRA

Munich Personal RePEc Archive

Boundary and interior equilibria: what drives convergence in a ‘beauty contest’?

Andrea, Morone and Piergiuseppe, Morone

2008

Online at <https://mpra.ub.uni-muenchen.de/9584/>
MPRA Paper No. 9584, posted 16 Jul 2008 00:45 UTC

Boundary and interior equilibria: what drives convergence in a ‘beauty contest’?

Andrea Morone

Università degli studi di Bari
Dipartimento di Scienze Economiche e Metodi
Matematici
a.morone@gmail.com

Piergiuseppe Morone

Università degli studi di Foggia
Dipartimento di Scienze Economiche Matematiche
e Statistiche
p.morone@gmail.com

Abstract

We present an experimental game in the p-beauty framework. Building on the definitions of boundary and interior equilibria, we distinguish between ‘speed of convergence towards the game-theoretic equilibrium’ and ‘deviations of the guesses from the game-theoretic equilibrium’. In contrast to earlier findings (Güth et al., 2002), we show, under a different game parameterisation, that (i) interior equilibria initially produce smaller deviation of the guesses from the game-theoretic equilibrium compared to boundary equilibria; (ii) interior and boundary equilibria do not differ in the timeframe needed for convergence; (iii) the speed of convergence is higher in the boundary equilibrium.

JEL classification: C72, C91

Keywords: Guessing game, p-beauty contest, individual behaviour

1. Introduction

The p-beauty contest game is a well known and extremely simple game (Keynes, 1936; Nagel, 1995; Duffy and Nagel, 1997; Canerer et al., 1998; Weber, 2003) where n players are asked to choose a number from a closed interval $[L, H]$. The winning player will be the one that gets closer to a target number G . Such target number is defined as the average of all guesses plus a constant (d), multiplied by a real number (p) known to all players.

Formally, we have: $G = p \left(\frac{1}{n} \sum_{i=1}^n g_i + d \right)$. In its simplest form the game parameterisation is

set as follows: $0 \leq p < 1$, d is equal to 0 and subject i 's guess is $g_i \in [0,100] \subset \mathbf{R}$.

Under such definition of G the game-theoretic solution is a unique Nash equilibrium where all players choose 0. In fact, playing 0 is the only strategy that survives the procedure of *iterated elimination of dominated strategies* (IEDS). Moreover, under such a standard parameterisation the game converges to the same value, and within the same number of iterations, if players follow Nagel's *iterative naïve best replies* (INBR) strategy.¹

The game becomes more complicated if we set $d \neq 0$; in this case the game might well exhibit an interior equilibrium (i.e. different from 0 or 100) and for specific values of p , the solution of the game obtained using the two different strategies (IEDS and INBR) involves different numbers of iterations needed to reach the equilibrium.

Güth et al. (2002) proposed a game where d was initially set equal to 0 and subsequently equal to 50. This allowed them to analyse the p-beauty contest from a different perspective, comparing, among other things, interior and boundary equilibria. They showed that the convergence toward the equilibrium is faster when the equilibrium is interior.

In this paper we shall confute Güth et al. (2002) finding showing that the speed of convergence is actually not necessarily faster for interior equilibria. We will look at both the 'speed of convergence towards the game-theoretic equilibrium' and the 'timeframe required for convergence' (i.e. the number of iterations). These two concepts are indeed different as they crucially depend on the initial distance of a system from its equilibrium level. In the p-beauty contest game, such distance is captured by the initial 'deviations of the guesses from the game-theoretic equilibrium'. We will prove that, for specific game's

¹ See Morone and Morone (2008) for a description of both strategies.

parameterisations, the speed of convergence can be slower in interior equilibria than in boundary equilibria.

The paper is structured as follows: in section 2 we present the experimental game and pose our research hypothesis. In section 3 we present the design of the experiment and in section 4 our findings. We conclude the paper in section 5.

2. Aim and setting of the experiment

The aim of our experiment is testing the validity of Güth et al. (2002) finding that interior equilibria (e.g. $d=50$) yield smaller deviations of the guesses from the game-theoretic equilibrium when compared to boundary equilibria;² this finding leads the authors to conclude that “swifter convergence to the equilibrium [is found] when the equilibrium is interior” (2002: 225).

Schematically, we can summarise Güth et al. experiment’s parameterisations and equilibria (s^*), in the following table:

	Parametrisation	Game-Theoretic equilibrium	Salient point à la Schelling	Number of iterations required for convergence	
				IEDS	INBR
Treatment 1	$p=1/2, d=0$ $g \in [0,100]$	$s^*=0$ (boundary equilibrium)	50	∞	∞
Treatment 2	$p=1/2, d=50$ $g \in [0,100]$	$s^*=50$ (interior equilibrium)	50	27	1

Table 1: Güth et al. (2002) summary of parameters and results

As we can see, the authors presented two comparable cases and showed how the treatment where the game-theoretic equilibrium is interior, requires less iterations for convergence. However, we can observe that for the interior equilibrium case there is the simultaneous coincidence, around the game-theoretic equilibrium, of two possible focal points: the d value as well as the salient point à la Schelling.³ Moreover, in the case of $d=50$, if we believe that subjects follow Nagel’s *iterative naïve best replies* strategy, they would converge towards the equilibrium after one iteration.

² A plausible explanation of this finding is that experimental subjects often try to avoid extreme choices (see, for instance, Rubinstein et al., 1997).

³ Nagel (1995) suggested that the salient point à la Schelling would be 50 (i.e. the middle of the interval). However, we believe there are other possible focal points; for instance d could be perceived as such by experimental subjects.

The picture changes when we look at the boundary equilibrium treatment. Now, the salient point *à la* Schelling and the d value differ, as the salient point is always set in the middle of the interval $[0,100]$. Moreover, both theoretical strategies (i.e. IEDS and INBR) predict an infinite number of iteration for complete convergence to the equilibrium.⁴

Departing from these observations we intend to test whether the short timeframe required for convergence is an actual property of interior equilibria or whether it is just arising from the *ad hoc* specification chosen by Güth et al. (2002). We do so considering a new set of problems' characterisation defined by different parameterisations of the game. Specifically, we shall compare the original parameterisation adopted by Güth et al. with a similar setting where we vary the value of p (set equal to $1/3$) and the value of d (set equal to 0, 33 and 50). It is worth noting that, like in the original experimental setting, these new parameterisations produce both interior and boundary equilibria.

If Güth et al.'s result is robust to different model parameterisations, we will always observe a shorter timeframe required for convergence in the games with interior equilibrium; otherwise, we shall confute the validity of their results for problems' parameterisations different from those originally selected by the authors.

3. The design of the experiment

In each treatment of the experiment there are $n = 32$ subjects divided into 8 groups, each of 4 subjects. In each group subjects have to guess a number in the real interval $[L, H]$. The closer their guess is to the target, the higher is the pay-off. The general form of the pay-off

function is:
$$u(g_i) = C - c \left| g_i - p \left(\frac{1}{n} \sum_{j=1}^n g_j + d \right) \right|.$$


The experiments were run in May 2008 at ESSE (Economia Sperimentale al Sud d'Europa) at the University of Bari. The software of the computerised experiment was developed in z-Tree (Fischbacher, 1998). Groups were formed randomly at the beginning of the experiment and were kept invariant over the whole experiment (i.e. 10 periods).

⁴ Please see table A1 in the annex where we report the converging patterns generated by IEDS and INBR.

4. Results

In this section we will analyse the results obtained in our experiments. However, before moving to our new findings we shall present the results obtained when running the experiment using exactly the same parameterisation adopted by Güth et al. (2002). This will serve to cast away any doubt on the presence of any source of difference between our experimental design and the one adopted by Güth et al.

In figure 1 we report the average values of treatments 1 and 2;⁵ the converging patterns obtained in our preliminary set of experiments replicate exactly those obtained by Güth et al., as it shows smaller deviations of the guesses from the game-theoretic equilibrium and a shorter time frame for convergence in the treatment with boundary equilibrium.⁶ In figure 2 we report two further treatments where p is kept equal to $1/2$ and d is set first equal to 33 and, subsequently, to 100. These two treatments produce very similar results to those reported in figure 1. Again, the system converges within a smaller number of iterations to the interior equilibrium, consistently displaying a smaller deviation of the guesses from the game-theoretic equilibrium.⁷



⁵ As mentioned earlier, each treatment was repeated 10 times. Hence, all values reported in figures 1 and 2 are averages of 10 rounds.

⁶ Deviation from the equilibrium are significantly smaller in treatment 2 than in treatment 1. $P < 0.01$ for rounds 1-10 and 1-5, $p < 0.1$ for rounds 6-10 (U-test, two-sided).

⁷ Deviation from the equilibrium are significantly smaller in treatment 3 than in treatment 4. $P < 0.01$ for rounds 1-10, 1-5, and 6-10 (U-test, two-sided).

Figure 1. Treatment averages ($p=1/2$; $d=0$ and $d=50$)




Figure 2. Treatment averages ($p=1/2$; $d=33$ and $d=100$)

We summarise the parameterisation and the results of these two last treatments in table 2. As we can see, in this case the boundary equilibrium does not present the simultaneous coincidence of the two focal points around the equilibrium value (i.e. d and the salient point *à la* Schelling are different).

	Parametrisation	Game-Theoretic equilibrium	Silient point <i>à la</i> Schelling	Number of iterations required for convergence	
				IEDS	INBR
Treatment 3	$p=1/2, d=33$ $g \in [0,100]$	$s^*=33$ (boundary equilibrium)	50	27	26
Treatment 4	$p=1/2, d=100$ $g \in [0,100]$	$s^*=100$ (interior equilibrium)	50	1	27

Table 2: Summary of parameters and results⁹

These first four treatments seem to confirm Güth et al. findings. We shall now move on to consider three other treatments where p is now set equal to $1/3$ and d is set equal to 0, 33 and 50, respectively.⁸ We report a summary of these new treatments' parameterisation and results in table 3.

⁸ Note that we do not consider the treatment with $p=1/3$ and $d=100$ as it converges to an interior equilibrium ($s^*=50$) and, therefore, is not comparable with $p=1/2$ and $d=100$.

	Parametrisation	Game-Theoretic equilibrium	Stient point à la Schelling	Number of iterations required for convergence	
				IEDS	INBR
Treatment 3	$p=1/3, d=0$ $g \in [0,100]$	$s^*=0$ (boundary equilibrium)	50	∞	∞
Treatment 4	$p=1/3, d=33$ $g \in [0,100]$	$s^*=16.5$ (interior equilibrium)	50	17	16
Treatment 5	$p=1/3, d=50$ $g \in [0,100]$	$s^*=25$ (interior equilibrium)	50	17	16

Table 3: Summary of parameters and results⁹

The picture emerging from these new treatments is rather different from what we obtained so far. First and foremost, we do not observe any significant difference in the timeframe of convergence towards the equilibrium across treatments: in fact, in all cases (i.e. one boundary and two internal) it takes for the system approximately the same number of rounds (about 6) to converge.¹⁰




Figure 3. Treatment averages ($p=1/3$; $d=0, d=33$ and $d=50$)

However, we can easily observe that the initial deviation of the guesses from the game-theoretic equilibrium is smaller in the two treatments that converge towards interior

⁹ Please see table A2 in the annex where we report the converging patterns generated by IEDS and INBR.

¹⁰ Comparing treatment 5 and treatment 6, treatment 5 and treatment 7, treatment 6 and treatment 7 we can always reject the hypothesis that deviation from the equilibrium is statistically significantly smaller in interior equilibrium treatments, for rounds 1–10. We can also reject the same hypothesis for rounds 5–10; however, we cannot reject it for rounds 1–5. These confirm that that initially deviation from the equilibrium is smaller in interior equilibrium treatments and that the number of rounds required for convergence is, nonetheless, similar in the both interior and boundary equilibria treatments. Hence, these tests conform the finding that speed of convergence is higher in the boundary equilibrium treatment.

equilibria. This finding suggests that the conclusions obtained by Güth et al. are only partially confirmed. While interior equilibria treatments initially produce smaller deviation, the speed of convergence is higher in the boundary equilibrium treatment. In fact, as observed above, all three treatments converge towards their theoretical equilibria in the same time frame; therefore, the treatment which starts converging from a further point (i.e. the one which displays a higher deviation from the game-theoretic equilibrium) must converge faster, as it clearly emerges from figure 3.

Hence, we can conclude that the initial deviation from the game theoretic equilibrium is always greater in boundary equilibria, independently from the initial parameterisation of the experiment. We share Güth et al.'s view that this finding might depend on the tendency to choose interior instead of extreme, boundary strategies. However, along this finding we also observe that under the new experiment parameterisation the timeframe required for convergence is the same for boundary and interior equilibria and, consequently, the speed of convergence is higher in boundary equilibria. In fact, if two runners reach the same target in the same timeframe, the one starting from further away must run faster in order to cover a larger distance in the same time.

5. Conclusions

The experiment presented in this paper follows quite closely Güth et al., 2002 as we attempt to investigate differences in the speed of convergence towards boundary and interior equilibria in the p-beauty contest game. In doing so, we consider both 'speed of convergence towards the game-theoretic equilibrium' and 'timeframe required for convergence'. These two concepts crucially depend upon the initial 'deviations of the guesses from the game-theoretic equilibrium'. In contrast to earlier findings (Güth et al., 2002), we obtain, with a different parameterisation of the game, the following results: (i) interior equilibria treatments initially produce smaller deviation compared to boundary equilibria treatments; (ii) interior and boundary equilibria treatments do not differ in the timeframe needed for convergence; (iii) the speed of convergence is higher in the boundary equilibrium treatment.

These leads us to conclude that Güth et al. findings hold only for a very specific game parameterisation and, therefore, cannot be generalised.

References

- Bosch-Domènech A., Montalvo J.G., Nagel R., Satorra A.: (2002). One, Two, (Three), Infinity, ...: Newspaper and Lab Beauty-Contest Experiments. *The American Economic Review*, 92(5):1687-1701.
- Camerer, C., Ho, T. H., and Weigelt, K.: (1998). Iterated dominance and iterated best response in experimental “p-beauty contest”. *The American Economic Review*, 88(4):947-969.
- Duffy, J., and Nagel, R.: (1997). On the robustness of behaviour in experimental “beauty contest” games. *The Economic Journal*, 107(445):1684-1700.
- Keynes, J. M. (1936). *The general theory of interest, employment and money*. London: Macmillan.
- Fischbacher, U. (2007), “Zurich toolbox for readymade economic experiments”, *Experimental Economics*, 10, 171–178.
- Güth, W., Kocher, M., and Sutter, M.: (2002). Experimental ‘beauty contest’ with homogeneous and heterogeneous players and with interior and boundary equilibria. *Economic Letters*, 74:219-228.
- Morone, A. and Morone, P. (2008). “Guessing game and people behaviours: What can we learn?”. In M. Abdellaboui and J. D. Hey (Ed.), “Advances in Decision Making under Risk and Uncertainty” Springer.
- Nagel, R.: (1995). Unraveling in guessing games: an experimental study. *The American Economic Review*, 85(5):1313-1326.
- Rubinstein, A., Tversky, A., Heller, D., 1997. Naive strategies in competitive games. In: Albers, W. et al. (Ed.), *Understanding Strategic Interaction: Essays in Honor of Reinhard Selten*. Springer, Heidelberg, pp. 394–402.
- Weber, R., A.: (2003). Learning with no feedback in a competitive guessing game. *Games and Economic Behavior*, 44:134-144.

Annex

Treatment 1		Treatment 2	
IEDS	INBR	IEDS	INBR
50	25	75	50
25	12.5	62.5	50
12.5	6.25	56.25	50
6.25	3.125	53.125	50
3.125	1.5625	51.5625	50
1.5625	0.78125	50.78125	50
0.78125	0.390625	50.390625	50
0.390625	0.1953125	50.195313	50
0.1953125	0.09765625	50.097656	50
0.09765625	0.048828125	50.048828	50
0.048828125	0.024414063	50.024414	50
0.024414063	0.012207031	50.012207	50
0.012207031	0.006103516	50.006104	50
0.006103516	0.003051758	50.003052	50
0.003051758	0.001525879	50.001526	50
0.001525879	0.000762939	50.000763	50
0.000762939	0.00038147	50.000381	50
0.00038147	0.000190735	50.000191	50
0.000190735	9.53674E-05	50.000095	50
9.53674E-05	4.76837E-05	50.000048	50
4.76837E-05	2.38419E-05	50.000024	50
2.38419E-05	1.19209E-05	50.000012	50
1.19209E-05	5.96046E-06	50.000006	50
5.96046E-06	2.98023E-06	50.000003	50
2.98023E-06	1.49012E-06	50.000001	50
1.49012E-06	7.45058E-07	50.000001	50
7.45058E-07	3.72529E-07	50	50
3.72529E-07	1.86265E-07	50	50
1.86265E-07	9.31323E-08	50	50
9.31323E-08	4.65661E-08	50	50
4.65661E-08	2.32831E-08	50	50
2.32831E-08	1.16415E-08	50	50

Table A1: IEDS and INBR game theoretical solution for treatments 1 and 2

Treatment 3		Treatment 4		Treatment 5		Treatment 6		Treatment 7	
IEDS	INBR								
66.5	41.5	100	75	33.333333	16.666667	44.333333	27.666667	50	33.333333
49.75	37.25	100	87.5	11.111111	5.555556	25.777778	20.222222	33.333333	27.777778
41.375	35.125	100	93.75	3.7037037	1.8518519	19.592593	17.740741	27.777778	25.925926
37.1875	34.0625	100	96.875	1.2345679	0.617284	17.530864	16.91358	25.925926	25.308642
35.09375	33.53125	100	98.4375	0.4115226	0.2057613	16.843621	16.63786	25.308642	25.102881
34.046875	33.265625	100	99.21875	0.1371742	0.0685871	16.61454	16.545953	25.102881	25.034294
33.523438	33.132813	100	99.609375	0.0457247	0.0228624	16.53818	16.515318	25.034294	25.011431
33.261719	33.066406	100	99.804688	0.0152416	0.0076208	16.512727	16.505106	25.011431	25.00381
33.130859	33.033203	100	99.902344	0.0050805	0.0025403	16.504242	16.501702	25.00381	25.00127
33.06543	33.016602	100	99.951172	0.0016935	0.0008468	16.501414	16.500567	25.00127	25.000423
33.032715	33.008301	100	99.975586	0.0005645	0.0002823	16.500471	16.500189	25.000423	25.000141
33.016357	33.00415	100	99.987793	0.0001882	9.408E-05	16.500157	16.500063	25.000141	25.000047
33.008179	33.002075	100	99.993896	6.272E-05	3.136E-05	16.500052	16.500021	25.000047	25.000016
33.004089	33.001038	100	99.996948	2.091E-05	1.045E-05	16.500017	16.500007	25.000016	25.000005
33.002045	33.000519	100	99.998474	6.969E-06	3.485E-06	16.500006	16.500002	25.000005	25.000002
33.001022	33.000259	100	99.999237	2.323E-06	1.162E-06	16.500002	16.500001	25.000002	25.000001
33.000511	33.00013	100	99.999619	7.744E-07	3.872E-07	16.500001	16.5	25.000001	25
33.000256	33.000065	100	99.999809	2.581E-07	1.291E-07	16.5	16.5	25	25
33.000128	33.000032	100	99.999905	8.604E-08	4.302E-08	16.5	16.5	25	25
33.000064	33.000016	100	99.999952	2.868E-08	1.434E-08	16.5	16.5	25	25
33.000032	33.000008	100	99.999976	9.56E-09	4.78E-09	16.5	16.5	25	25
33.000016	33.000004	100	99.999988	3.187E-09	1.593E-09	16.5	16.5	25	25
33.000008	33.000002	100	99.999994	1.062E-09	5.311E-10	16.5	16.5	25	25
33.000004	33.000001	100	99.999997	3.541E-10	1.77E-10	16.5	16.5	25	25
33.000002	33.000001	100	99.999999	1.18E-10	5.901E-11	16.5	16.5	25	25
33.000001	33	100	99.999999	3.934E-11	1.967E-11	16.5	16.5	25	25
33	33	100	100	1.311E-11	6.557E-12	16.5	16.5	25	25
33	33	100	100	4.371E-12	2.186E-12	16.5	16.5	25	25
33	33	100	100	1.457E-12	7.285E-13	16.5	16.5	25	25
33	33	100	100	4.857E-13	2.428E-13	16.5	16.5	25	25
33	33	100	100	1.619E-13	8.095E-14	16.5	16.5	25	25
33	33	100	100	5.397E-14	2.698E-14	16.5	16.5	25	25

Table A2: IEDS and INBR game theoretical solution for treatments 3, 4, 5, 6 and 7