



Munich Personal RePEc Archive

**The distribution of the average of  
log-normal variables and exact Pricing of  
the Arithmetic Asian Options: A Simple,  
closed-form Formula**

Alghalith, Moawia

30 November 2019

Online at <https://mpra.ub.uni-muenchen.de/99089/>  
MPRA Paper No. 99089, posted 18 Mar 2020 07:47 UTC

**The distribution of the sum of log-normal variables and exact  
pricing of the arithmetic Asian options: A simple, explicit  
formula**

Moawia Alghalith

UWI, St Augustine

malghalith@gmail.com

**Abstract:** We overcome a long-standing obstacle in statistics. In doing so, we show that the distribution of the sum of log-normal variables is log-normal. Furthermore, we offer a breakthrough result in finance. In doing so, we introduce a simple, exact and explicit formula for pricing the arithmetic Asian options. The pricing formula is as simple as the classical Black-Scholes formula.

Short title: The distribution of the sum of lognormals

Keywords: the distribution of the sum of log-normals, arithmetic Asian option, the Black-Scholes formula.

# 1 Introduction

A long-standing obstacle in statistics is the determination of the distribution of the sum of log-normal variables. This paper overcomes this obstacle and shows that the distribution of the sum of log-normal variables is log-normal.

Recent literature used orthogonal polynomial expansions to approximate the distribution of the arithmetic average. Examples include Willems (2019) and Asmussen et al (2016). Some of the literature used Edgeworth expansions to approximate the distributions (see, for example, Li and Chen (2016)). Others such as Aprahmian and Maddah (2015) used the Gamma distribution approach. Some studies relied on Monte Carlo simulations. Examples include Lapeyre et al (2001) and Fu et al (1999). Others adopted a numerical approach. Examples include Linetsky (2004), Cerny and Kyriakou (2011), and Fusai et al (2011). Curran (1994) used the geometric mean to estimate the arithmetic mean.

The literature on pricing the arithmetic Asian options has two main features in common. First, it relies on approximations. Secondly, it largely adopts (very) complex methods. Consequently, this paper overcomes these two limitations. In this paper, we use a pioneering approach to pricing the

arithmetic Asian options. In doing so, we present an exact (yet very simple) method. Particularly, we show that the price of the arithmetic Asian option is exactly equivalent to the price of the European option with an earlier (known) expiry. The pricing formula is as simple as the classical Black-Scholes formula.

## 2 The method

The arithmetic average of the price underlying asset  $S(u)$  over the time interval  $[t, T]$  is given by

$$A_t = \frac{\int_t^T S(u) du}{T - t},$$

where  $t$  is the current time and  $T$  is the expiry time. So that, using the Black-Scholes assumptions,  $EA_t = E \frac{\int_t^T S(u) du}{T - t} = \frac{e^{r(T-t)} - 1}{r(T-t)} S(t)$ , where  $r$  is the risk-free rate of return. By the mean value theorem for integrals,  $E \frac{\int_t^T S(u) du}{T - t} = ES(\hat{t})$ , where  $\hat{t}$  is a time such that  $t < \hat{t} < T$ , and  $ES(\hat{t}) = e^{r(\hat{t}-t)} S(t)$ . This implies that  $\frac{e^{r(T-t)} - 1}{r(T-t)} = e^{r(\hat{t}-t)}$ . We can solve for  $\hat{t} - t$  as follows

$$\hat{t} - t = \frac{\ln\left(\frac{e^{r(T-t)} - 1}{r(T-t)}\right)}{r}.$$

Thus  $\hat{t}$  is known. For example, if  $T - t = 1$  and  $r = .01$ ,  $\hat{t} - t = \frac{\ln\left(\frac{e^{.01} - 1}{.01}\right)}{.01} = .498$ . We also can show that  $A_t$  is log-normal<sup>1</sup> and equal to  $S(\hat{t})$ ; thus the Black-Scholes formula can be directly and exactly applied. That is, the price of the Asian option (expiring at time  $T$ ) is given by

$$C(t) = e^{-r(\hat{t}-t)} E [S(\hat{t}) - K]^+ = e^{-r(\hat{t}-t)} E [A_t - K]^+,$$

where  $K$  is the strike price. Clearly, this is the price of a European option with expiry  $\hat{t}$ . Thus, the price of the arithmetic Asian option (with expiry time  $T$ ) is equal to the price of the equivalent European option with expiry time  $\hat{t}$ . This explains why the Asian option is cheaper than its European counterpart.

Needless to say, the pricing formula for an arithmetic Asian call with expiry time  $T$  is

---

<sup>1</sup>See the appendix for the proofs.

$$C(t, s) = sN(d_1) - e^{-r(\hat{t}-t)}KN(d_2),$$

where  $s$  is the current price,  $d_1 = \frac{1}{\sqrt{\sigma^2(\hat{t}-t)}} [\ln(s/K) + (r + \sigma^2/2)(\hat{t} - t)]$ ,  $d_2 = d_1 - \sqrt{\sigma^2(\hat{t} - t)}$ , and  $\sigma$  is the volatility of the return rate of the underlying asset.

**Practical example:**

If  $r = .05$ ,  $T = 1$ ,  $\sigma = .2$ ,  $s = K = \$100$ , then  $\hat{t} = .502$  and thus the option price is  $C(t) = \$6.91$ .

**Appendix. Proofs of  $A_t$  is log-normal.**

1. Consider the stock price,  $S(T) - s = \int_0^T dS(t)$ , where  $s \equiv S(0)$ ;

squaring both sides yields

$$(S(T))^2 + s^2 = 2sS(T) + \left( \int_0^T dS(t) \right)^2 = 2sS(T) + \sigma^2 \int_0^T (S(t))^2 dt \quad (1)$$

since  $(dS(t))^2 = \sigma^2 (S(t))^2 dt$ . The left-hand-side of (1) is clearly log-normal (a lognormal plus a constant), and the right-hand-side of the equation is a sum of lognormal variables; therefore, the sum (or average) of log-normal

variables is log-normal.

We can also present the sum without the constant  $s^2$  by differentiating both sides of (1) with respect to  $r$

$$\frac{\partial (S(T))^2}{\partial r} = 2s \frac{\partial S(T)}{\partial r} + \sigma^2 \int_0^T \frac{\partial (S(t))^2}{\partial r} dt$$

clearly the left-hand-side of the above equation is log-normal, and the right-hand-side of the equation is a sum of log-normal variables.

We can also show that the integral alone is log-normal; dividing both sides of the above equation by  $S(T)$  yields

$$2TS(T) = 2Ts + \sigma^2 \int_0^T \frac{\partial (S(t))^2}{S(T) \partial r} dt,$$

differentiating twice w.r.t.  $r$

$$2T \frac{\partial^2 S(T)}{\partial r^2} = \sigma^2 \int_0^T \frac{\partial^2 X}{\partial r^2} dt,$$

where  $X \equiv \frac{\partial (S(t))^2}{S(T) \partial r}$ ; the left-hand-side of the above equation is log-normal, and the right-hand-side of the equation is a sum of log-normal variables.  $\square$

2. The simplest and intuitive proof is that the time continuity implies

that the average price  $A_t$  is a price on the interval  $[S(t), S(T)]$ . To be more precise, each (random) price at a specific time is an interval-valued (an interval of all possible outcomes of the price). Thus the elements of  $[S(t), S(T)]$  are (vertical) intervals, then the time continuity guarantees the existence of a vertical interval of outcomes on  $[S(t), S(T)]$ , but the vertical interval is a price at a specific time. So the difference between a random variable and a non-random variable is that the random variable is interval-valued, and thus the mean-value theorem can be applied in the same way to non-random variables if we view the elements of  $[S(t), S(T)]$  as interval-valued.

3. The outcomes of  $A_t$  are the averages of paths and therefore they are outcomes (realizations) of prices. That is, each outcome is in the form  $S(t) e^{(r-\frac{1}{2}\sigma^2)u+\Omega_1}$ , where  $\Omega$  is an outcome of a Brownian motion; thus it can be expressed as  $A_t = S(t) e^{(r-\frac{1}{2}\sigma^2)u+W(u)}$ ; otherwise it will not be possible, using the price probability density, to obtain  $EA_t = S(t) e^{r(\hat{t}-t)}$ .

4. Using the classical mean-value theorem, it is straightforward to show that the variance of  $A_t$  is in the form  $S(t)^2 e^{2ru} (e^{\sigma^2 u} - 1)$ . Higher moments can also be obtained by the classical mean-value theorem; and on a bounded

interval, the distribution is identified by its moments.

In sum, this paper offers two ground-breaking contributions. The first one is in mathematical statistics (the distribution of the sum of log-normal variables). The second one is in finance (an explicit, simple formula for the price of the arithmetic Asian options). The first contribution will have a great impact on statistics since it will have so many applications in the future.

## References

- [1] Aprahamian, H. and B. Maddah (2015). Pricing Asian options via compound gamma and orthogonal polynomials. *Applied Mathematics and Computation* 264, 21–43.
- [2] Asmussen, S., P.-O. Goffard, and P. J. Laub (2016). Orthonormal polynomial expansions and lognormal sum densities. arXiv preprint arXiv:1601.01763.
- [3] Cerny, A. and I. Kyriakou (2011). An improved convolution algorithm for discretely sampled Asian options. *Quantitative Finance* 11, 381–389.

- [4] Curran, M. (1994). Valuing Asian and portfolio options by conditioning on the geometric mean price. *Management Science*, 40, 1705–1711.
- [5] Fu, M. C., D. B. Madan, and T. Wang (1999). Pricing continuous Asian options: a comparison of Monte Carlo and Laplace transform inversion methods. *Journal of Computational Finance* 2, 49–74.
- [6] Fusai, G., D. Marazzina, and M. Marena (2011). Pricing discretely monitored Asian options by maturity randomization. *SIAM Journal on Financial Mathematics* 2, 383–403.
- [7] Lapeyre, B., E. Temam, et al. (2001). Competitive Monte Carlo methods for the pricing of Asian options. *Journal of Computational Finance* 5, 39–58.
- [8] Li, W. and S. Chen (2016). Pricing and hedging of arithmetic Asian options via the Edgeworth series expansion approach. *The Journal of Finance and Data Science* 2, 1–25.
- [9] Linetsky, V. (2004). Spectral expansions for Asian (average price) options. *Operations Research* 52, 856–867.

- [10] Willems, S. (2019). Asian option pricing with orthogonal polynomials, *Quantitative Finance*, 19, 605-618.