



Munich Personal RePEc Archive

# **Boltzmann Entropy in Cryptocurrencies: A Statistical Ensemble Based Approach**

Grilli, Luca and Santoro, Domenico

Università degli Studi di Foggia

11 April 2020

Online at <https://mpra.ub.uni-muenchen.de/99591/>  
MPRA Paper No. 99591, posted 17 Apr 2020 10:47 UTC

# Boltzmann Entropy in Cryptocurrencies: A Statistical Ensemble Based Approach

Luca Grilli · Domenico Santoro

## Abstract

In this paper we try to build a statistical ensemble to describe a cryptocurrency-based system, emphasizing an “affinity” between the system of agents trading in these currencies and statistical mechanics. We focus our study on the concept of entropy in the sense of Boltzmann and we try to extend such a definition to a model in which the particles are replaced by  $N$  agents completely described by their ability to buy and to sell a certain quantity of cryptocurrencies. After providing some numerical examples, we show that entropy can be used as an indicator to forecast the price trend of cryptocurrencies.

**Keywords** Cryptocurrency · Entropy · Prices Forecast · Boltzmann · Blockchain

**Mathematics Subject Classification (2000)** 91G80 · 62P05 · 28D20 · 91B80 · 62P20 · 82B30

---

Luca Grilli · Domenico Santoro

Università degli Studi di Foggia, Dipartimento di Economia, Management e Territorio, Via Da Zara, 11, I-71121  
Foggia, Italy

E-mail: luca.grilli@unifg.it

## 1 Introduction

Trying to model the price trend of financial instruments has always been the focus of finance. Traditional theories are based on the search for significant variables that can explain the trend, such as Fama and French [11], Campbell and Shiller [39] and Perasan and Timmermann [12]. The evolution of the theory has pushed towards non-linear models such as Laffont et al. [17] and Qi [25]. Subsequently, Brissaud [5] assimilated **entropy** to disorder, so as to make this tool that has always been applied the physics part of the economy.

Since the mid-19th century, entropy has been a key element linking mechanics to thermodynamics. The first to introduce this concept was Clausius [31], whose definition was applied to a thermodynamic system that performs a transformation; however, this entropy suffered from a conceptual problem which, as demonstrated by Gibbs [38], was revealed in the case of identical gases (*Gibbs Paradox*). He solved this problem by changing the count of states. On the other hand, Boltzmann [20] presented his statistical interpretation of thermodynamic entropy, managing to link the macroscopic properties of a system with the microscopic ones. Based on Gibbs, in 1949 Shannon [9] developed a theory capable of evaluating the amount of information that is lost in receiving a message from a source to a recipient. This form of entropy was generalized by Rényi [4], Tsallis [8], Adler [19] (in topology), redefined by Pincus [24] (*approximate entropy*) and - more recently - by Chen [37] as a time series regularity measure.

It was the development of entropy in information theory that made it widely used in finance, in particular the generalizations of Shannon, Rényi and Tsallis contributed to creating a new line of application for the management of financial portfolios. These new types of entropy has been used by Philippatos and Wilson [7], Usta and Kantar [13], Jana [30], Gulko [21] that introduced the *Entropy Pricing Theory*, Nawrocki [29], Dionisio [2] and Ormos [23].

In this paper, we want to demonstrate that it is possible to assimilate the system of cryptocurrencies to thermodynamic systems to be able to determine their entropy in the sense of Boltzmann so that we can make price predictions related to the possibility that they move in a more or less wide range; unlike all the recent applications concerning theories based on Shannon entropy and its derivations. Innovation is linked to the reinterpretation of the monetary system of cryp-

tocurrencies. In this sense, we can apply physical theories to a social science. It is interesting to develop this approach as we assume that the physical system described by adapting the economy can be summarized by the movements that cryptocurrencies perform the currency markets. Once the system has been described, our goal is to verify that entropy calculated in the physical sense also occurs in the economic context to allow us to make assumptions on how the process could move in the next future. This type of conjecture has been presented by Sergeev [26], Zakiras [36], Khrennikov [3] and Smith and Foley [10]. In particular McCauley [16], based on this previous theory, maintains that the illiquidity of the markets does not allow for the application of the concepts of statistical mechanics.

The paper structure is the following: in Section 2 we analyze cryptocurrencies and their key characteristics, focusing on the fact that they have a supply limit; in Section 3 we describe the evolution of a system of a particle in statistical thermodynamics and how to determine its entropy, subsequently applying these notions to our monetary system; in Section 4 we define the theoretical assumptions we can link to the system created previously to study the price evolution in these currency markets and we analytically describe the calculation of entropy; finally in Section 5 we show the entropy values and the corresponding price movements using real data.

## 2 Cryptocurrency

Cryptocurrencies represent a digital currency system with no guarantee institution and no transaction control. The main cryptocurrencies, by media coverage or by the possibility that some financial intermediaries offer to use them as a payment instrument, are: Bitcoin, Ethereum, Ripple.

Unlike traditional financial assets, their value is not based on tangible assets such as the economy of a country or a company, but it is based on the security of an algorithm that tracks transactions. Their definition is controversial since by some entities [14] they are considered intangible assets (IFRS) while according to the German financial supervisory authority (BsFin) they are officially financial instruments [6]. Just as specified by Corbet et al. [32] the literature is still immature and new empirical and theoretical evidence continues to emerge monthly. Moreover,

the same authors claim that in cryptocurrencies there are unique and specific issues that cannot be addressed using quantitative research and data mining as regulatory disorientation, cyber-criminality, and environmental sustainability.

All the cryptocurrencies have been based on the *Bitcoin*, a currency created by Nakamoto [33] who in 2009 released a software capable of implementing transactions. The currency itself is a unique alphanumeric string that represents a certain transaction, a transaction which will then be entered in a public register called *blockchain*.

The transfer of the currency takes place through a digital signature mechanism by using the value of a function (called *hash function*) which is inserted in the previous transaction and guarantees its authenticity. A hash function has some properties that guarantee its safety thanks to the resistance to the preimages and the most used functions are the SHA256 and the RIPEMD. Transactions through cryptocurrencies are encrypted through digital signature mechanisms based on public or private keys.

The blockchain is the fulcrum of these systems and is essentially a register in which the data of the owners of the currency are entered, transactions occur in an encrypted manner. The blockchain is a data structure consisting of a list of transaction blocks linked together so that each refers to the previous one in the chain. Each block in the blockchain is identified by a hash generated using the SHA256 cryptographic algorithm on the block header. A block is a data structure that aggregates transactions to include them in the public register. The block is made of a header, containing metadata, followed by a long list of transactions. A complete block, with all transactions, is, thus, 1000 times larger than the block header [1]. The first identifier of a block is the cryptographic hash generated by the SHA256 algorithm, which returns, as a result, a 32-bit hash called *block hash*; the second identifier is the position in the blockchain called *block height*.

The cryptocurrency generation process is called *mining*, which adds money to the supply. Cryptocurrencies are “minted” during the creation of each block at a fixed and decreasing rate [1]: each block generated on average every 10 minutes contains new currency. For example, if we consider Bitcoin, every 210000 blocks the currency issue rate decreases by 50% (the availability of new coins grows as a geometric series every 4 years). It is estimated that around the year 2140, the production of the last block will be reached (6930000) and the number of coins produced

will tend to its upper limit of 21 million (precisely 20999999.97690000), value introduced by Nakamoto himself and contained in the variable “MAX\_MONEY” as can be read in the source code present on GitHub.<sup>1</sup> This value represents a sanity check, especially used to avoid bugs in which it is possible to generate currency from nothing and therefore moving towards a situation in which the blockchain diverges into different potential paths (called *fork*).

The integrity of the blockchain network is guaranteed through consensus algorithms such as *Proof-of-Work* (PoW) and *Proof-of-Stake* (PoS), that solve the Byzantine Generals Problem [15] (a problem of consent in the presence of errors). A consensus algorithm is a mechanism used by the network to reach consensus, i.e. ensuring that the protocol rules are followed and that transactions occur correctly so that coins can only be spent once.

### 3 Methodology

The main assumption in this paper is that the prices of cryptocurrencies behave like a thermodynamic system, so it is possible to determine entropy by using the Boltzmann formula. In order to present the theoretical framework and the methodology, we need to briefly introduce the main physical results. In Statistical Mechanics a macroscopic system is made up of  $N$  molecules ( $N \sim 10^{24}$  is the Avogadro’s constant) whose mechanics provide the evolution of  $6N$  dynamic variables describing completely the microscopic states of this system. Motion in the phase space can be studied using the  $3N$  position components and the  $3N$  momenta components, indicated with  $\{\mathbf{q}_i\}$  and  $\{\mathbf{p}_i\}$  whose evolution is driven by Hamilton’s equations.

Mechanics, therefore, provides a very detailed description of the system contrary to thermodynamics which studies the collective variations; for this reason, the mechanical point of view can be defined *microscopic* and the thermodynamic one *macroscopic*. The study of the system from a microscopic point of view concerns experimental observation on one or a few molecules.

Everything that happens from the microscopic side can be expressed in macroscopic terms through thermodynamics, defined in this case as a large amount of microscopic variables. We consider an isolated system of  $N$  particles described by the  $3N$  coordinates and the  $3N$  momenta in

---

<sup>1</sup> Source: <https://github.com/bitcoin/bitcoin/blob/master/src/amount.h>

a  $6N$ -dimensional space at a certain time  $t$ . Particles are subject to the laws of classical mechanics and therefore  $\mathbf{X}(t)$  evolves according to Hamilton's equations. Since the Hamiltonian  $H(p, q)$  does not depend on time, the energy  $E$  is a conserved quantity during motion and develops on a fixed hypersurface. We want, for example, to measure an observable  $A(\mathbf{X})$  (a function defined in the phase space) of the system in thermodynamic equilibrium, but since the scale of macroscopic times is much larger than the microscopic one, we can consider a datum as the result of a system that has gone through a large series of microscopic states; this implies that the observable must be compared with an average performed along with the evolution of the system calculated over very long times  $\bar{A}$ . The calculation of  $\bar{A}$  would require knowledge of both the microscopic state at a certain moment and the determination of the corresponding trajectory in the phase space, which corresponds to a practically inexhaustible request. To determine the observable, the *ergodic* theory intervenes, according to which each energy surface is completely accessible to any motion with the given energy and the average residence time in a certain region is proportional to its volume. If these conditions are satisfied, the average  $\bar{A}$  can be calculated as the average of  $A(\mathbf{X})$  in which the states with the fixed energy contribute with equal weight. In applications it is convenient to consider on average all states with energy within a fixed range  $[E, E + \Delta E]$ ; furthermore, we are only interested in some macroscopic properties such as particle number  $N$  and the volume  $V$ . There is an infinite number of systems that satisfy these conditions: these form the *Gibb's ensemble* which is represented by a set of points in the phase space characterized by a density function  $\rho(p, q, t)$  defined so that  $\rho(p, q, t) d^{3N}p d^{3N}q$  corresponds to the number of representative points of the system during the instant  $t$  contained in the infinitesimal volume of the phase space  $d^{3N}p d^{3N}q$ . Furthermore, since energy, volume and number of particles are constants of motion, the total number of systems in an ensemble is conservative.

We can thus introduce the *postulate of equal a priori probability* who claims that when a macroscopic system is in thermodynamic equilibrium its state can be with equal probability each of those which satisfies the macroscopic conditions of the system. This postulate implies that the

system under consideration belongs to an ensemble called *microcanonic* with density function

$$\rho(p, q) = \begin{cases} \rho^* & \text{if } E < H(p, q) < E + \Delta \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

where  $\rho^*$  is constant and all members of the ensemble have the same number of particles and equal volume.

We can define  $\Gamma(E)$  the volume occupied by the microcanonical ensemble in the phase space as:

$$\Gamma(E) \equiv \int_{E < H(p, q) < E + \Delta E} d^{3N} p d^{3N} q \quad (2)$$

and  $\Sigma(E)$  the volume bounded by the energy surface  $E$ :

$$\Sigma(E) \equiv \int_{H(p, q) < E} d^{3N} p d^{3N} q \quad (3)$$

so that

$$\Gamma(E) = \Sigma(E + \Delta E) - \Sigma(E). \quad (4)$$

Entropy, then, can be defined as:

$$S_\Gamma = \kappa_B \ln \Gamma(E) \quad (5)$$

where  $\kappa_B \sim 1.3806 * 10^{-23}$  is the Boltzmann constant.

Let us now try to translate this physical theory into a financial dress. Viaggiu et al. [34] have developed a representation of an economic model relating to money from a thermodynamic point of view. In their description the ensemble is made up of the  $N$  interacting economic subjects, entirely described by two variables  $\{x_i, y_i\}$  which represent money and credit/debt capacity and which are not conjugated in the sense of mechanics Hamiltonian. The key characteristic is to consider a representative function of the total currency as a conservative law, to be able to exploit the ergodic hypothesis.

Our idea is to go back to their hypothesis by applying it to the case of cryptocurrencies. We consider a model in which the particles are replaced by  $N$  economic subjects (agents) who intend to trade in cryptocurrencies (compared only to a reference currency, such as the USD). These agents are completely described by 2 variables, which we can, however, identify as  $\{x_i, y_i\}$ , where  $x_i$  and  $y_i$  indicate, respectively, the ability to buy and to sell a certain quantity of cryptocurrencies

(both expressed in monetary terms). The latter hypothesis is possible according to the fact that the market to which we refer is influenced only by the supply and demand leverage. As for [34], even if the complete Hamiltonian formalism is not respected, we can consider as a conserved quantity the total number of cryptocurrencies in circulation which by their definition is constant over a suitable time interval through the function  $M(x_i, y_i)$  (as in the particular case of Bitcoins for which the supply limit is fixed at 21 million). However, since the supply limit has not yet been reached by any cryptocurrency we consider this quantity constant concerning the currency in circulation in a precise time  $t$ , therefore:

$$M = \sum_{i=1}^N x_i + y_i. \quad (6)$$

In this sense, the sum of the ability to sell and buy of the  $N$  agents fully describes the cryptocurrencies in circulation. The ergodic hypothesis allows us, given a certain function  $f(x_i, y_i)$ , to express its average with respect to the time in terms of an average over the ensemble at fixed  $M$ :

$$\bar{f} = \int_{M=const} f(x, y) \rho(x, y) dx dy \quad (7)$$

where  $\rho(x, y)$  denotes the probability distribution of the ensemble. Through these assumptions we can verify the economic transformations through thermodynamics; in particular, as in statistical mechanics, we can calculate the *volume in the phase space* [34]. If we integrate over all the available volume of the configuration space spanned by  $\{x, y\}$  with  $\bar{M} = m$  (where  $\bar{M}$  denotes the average over the whole configuration space) we have  $\int_{\bar{M}=m} d^N x d^N y = 0$ . So introducing a thick shell  $\Delta$  where  $\Delta \ll m$  we can define:

$$\Gamma(m) = \int_{m < M < m + \Delta M} \frac{d^N x d^N y}{k^{2N}} \quad (8)$$

where  $d^N x d^N y$  is understood as the phase space and  $k$  is a normalization factor such that  $\Gamma$  is dimensionless. This functional represents the number of microscopic realizations of the system under examination and allows us to calculate the entropy  $S$  as described in the equation (5).

#### 4 The model

In the previous section, we have hypothesized that we can consider financial transactions concerning cryptocurrencies as a thermodynamic system. In this section, however, we try to define, through a new type of approach, how it is possible to calculate entropy considering essentially the prices obtainable from the currency markets (FOREX).

First, we know that cryptocurrencies are used by an approximate number of economic entities equal to 44 million for which  $N \gg 1$ . We also know that every subject in our system is fully described by its ability to buy and sell  $(\{x_i, y_i\})$ . Let us consider that these two variables are summarized in the *last prices* of the cryptocurrency on the currency markets, a type of price used to keep track of changes in the value of an asset throughout a session. In this sense, the latest prices allow us to understand whether, compared to the previous session, the ability to buy or sell prevailed. We can summarize this price capability in the sentence *“prices describe the strength with which agents position themselves in the phase space”*.

The key point is that we can use the function  $M$  (described above) because in a certain time  $t$  the quantity of cryptocurrencies is constant and quantifiable, in this way we can go back to the previous economic model and determine  $\Gamma$  as described in the equation (8). Analytically, we do not consider the number of economic subjects present in the market but indirectly deduce their “position” in the phase space from the difference between the closing prices. The process that led to the definition of the results is the following:

- We take a certain reference interval (5 days) and cluster the closing price series based on this interval;
- For each cluster there is a maximum and a minimum price, we calculate the difference in terms of necessary steps to pass from one to the other obtaining a certain value of **gap G** (this assumption is based on the idea that the distance between maximum and minimum is a measure of the dispersion of agents in our phase space);
- We use combinatorial analysis considering the value used for clustering to determine the “volume” occupied by the disposition of the agents, therefore:

$$\Gamma = G^5 \tag{9}$$

Once the value of  $\Gamma$  is determined, entropy can be calculated by using the Boltzmann formula:

$$S = \kappa_B \ln \Gamma. \quad (10)$$

Finally, precisely because Boltzmann's constant is of the order of Avogadro's number, we can "*rationalize*" this entropy value obtained by multiplying it by  $10^{23}$ .

Our data analysis shows that in situations where entropy is drastically reduced, in the following phase it must grow in an "almost obligatory" way; this in terms of cryptocurrency prices indicates that in situations in which the gap between the maximum and the minimum is drastically reduced in the transition from one cluster to another "almost compulsorily" follows a situation in which it is certainly wider than the previous one. This type of price-based entropy defines how agents move in the phase space, so it allows us to understand if there is more movement towards one area rather than another.

#### 4.1 Dataset

The empirical analysis has been applied to the closing prices of the main three cryptocurrencies<sup>2</sup>, all related to the US dollar (USD), that are:

- Bitcoin, whose price with 1 decimal digit provides for a step equal to 0.1;
- Ethereum, whose price with 2 decimal places provides for a step equal to 0.01;
- Ripple, whose price with 5 decimal places provides a step equal to 0.00001.

Prices are considered with a daily time frame over 1 year, from 1/1/2019 to 31/12/2019 and they are clustered in 5 days. To make the figures more clear, the 1-year interval has been divided into 4 trimesters. Furthermore, to better test the idea, the same test was carried out also on daily prices at 1 minute of 1/4/2020 recorded from 10:56 to 11:52, instead of clustered in 5 minutes. The difference from the daily case is that these prices were collected, always from the same source, but observed on different currency markets; in particular Bitcoin on the GDAX exchange, Ethereum on Bibox exchange and Ripple on Binance exchange.

---

<sup>2</sup> Source: Investing.com

## 4.2 Numerical examples

We can start the analysis from the annual case. The first cryptocurrency analyzed is Bitcoin (BTC/USD). We distinguish the trend of entropy compared to prices in the 4 ranges previously defined:




Figure 1: Prices (blue) and entropy (orange) Bitcoin in the period 1/1 - 31/12

As can be seen graphically, when entropy reaches a point of relative minimum falling below a certain threshold (it therefore undergoes a sharp reduction) it is forced in the next cluster to grow, almost as if to rebalance itself. This hypothesis does not seem to occur in the case of entropy lowering only a few points, as seen in the figure 1(c) in the case of clusters 3, 4 and 5. In terms of prices, this implies that in the cluster in which the entropy descent occurred there was

a very small gap and, in the subsequent cluster, since entropy increases the gap also increases. The last result, translated into the economic model created previously, indicates that in clusters in which entropy drops drastically, economic subjects concentrate in a relatively small “volume”. The second cryptocurrency analyzed are Ethereum (ETH/USD) as in the previous case:




Figure 2: Prices (blue) and entropy (orange) Ethereum in the period 1/1 - 31/12

Again, especially as seen in the figure 2(c) when entropy decreased sharply after a period of standing (small ups and downs) in the following period it was “forced” to grow. A particular situation, however, occurs in the figure 2(a) and in particular in clusters 2, 3 and 4: in this case entropy continued to fall despite having suffered a sudden movement. This situation allows us to highlight 2 things: that the sharp drop comes from a situation where the gap between the max and min prices is lower than a certain threshold value and that probably the descent of entropy

should not be considered only in the passage from one cluster to another, but in the passage from groups of clusters; in fact, if we consider clusters 2 to 5 as a single group, it is easy to see how entropy has undergone a really sharp reduction followed by growth.

The third cryptocurrency is represented by Ripple (XRP/USD):




Figure 3: Prices (blue) and entropy (orange) Ripple in the period 1/1 - 31/12

The situation is not very different from the Ethereum, but the marked variations are related to the fact that this cryptocurrency moves in a price range  $[0, 1]$  for which every movement is important. As seen in the figure 3(d) in clusters 4, 5 and 6 the entropy has returned to its original level following a sharp fall. We expected A growth but the fact that it has grown so much is related to the range in which prices move.

As for the case of 1-minute prices, we can summarize the trend of the different cryptocurrencies

together as shown in figure 4 which shows how all the assumptions made in the previous case are also respected for prices of this type.




Figure 4: Prices (blue) and entropy (orange) of cryptocurrencies based on 1 minute

## 5 Effects on prediction

We can, at this point, use entropy as an indicator to make predictions on the price trend of cryptocurrencies in the currency markets (which for the thermodynamic system correspond to the position of economic subjects in the phase space). Suppose that we are in a cluster  $C$  where entropy has declined sharply. As previously defined, we expect entropy to grow in the next cluster and this leads to an increase in the price gap. The hypothesis we can make is that the value of

the gap in the cluster  $C + 1$  is at least one unit higher than the value in the cluster  $C$ : we can use this information to understand what the future price range will be.

Let's consider a series of clusters in which we know the trend of prices and entropy (in this case we have considered an extract of the last Bitcoin price in the period 1/1/2019 - 31/3/2019). We know that in cluster 5, entropy has greatly reduced so we expect it to grow in cluster 6, creating a greater gap between prices than the previous one. At this point, knowing the value of the gap in the cluster  $C$ , we can create a bifurcation that represents the possible evolution of the price in the event of a bullish or bearish trend. Suppose a gap value 4 times larger than the previous one (as happens in reality) and a first cryptocurrency price close enough to the last price of the previous cluster; what we can expect is such a situation:

- If the second closing price of the cluster  $C + 1$  is **higher** than the previous price in the same cluster and assuming an upward trend we can assume that the series of prices continues in an area that we have defined as  $Gap^-$  (indicated in red);
- If the second closing price of the cluster  $C + 1$  is **lower** than the previous price in the same cluster and assuming a bearish trend we can assume that the price series continues in an area that we have defined as  $Gap^+$  (indicated in green).




Figure 5: Forecast effect on prices

Such information can be fundamental for example for an investor who intends to choose the ideal moment to enter (or exit) the market or balance any price limits.

## 6 Conclusions

In this paper, we have shown how it is possible to apply Boltzmann's entropy to cryptocurrencies. We have defined a similarity between a thermodynamic system and a currency system based on cryptocurrencies characterized by the presence of  $N$  subjects interested in buying (or selling) this type of currency, and assuming that the quantity of money at a certain moment  $t$  is fixed and determinable it is possible to hypothesize that the position of each economic entity is summarized by the last price of the cryptocurrency itself in the currency markets. With this hypothesis, it was possible to determine the entropy using the Boltzmann formula, in particular, its calculation was made by dividing the time interval into clusters and calculating the gap between the different prices. This analysis has shown that when entropy falls sharply then it must necessarily grow shortly; which in terms of price corresponds to a situation in which the gap between maximum and minimum is wider than the previous one.

## References

1. Antonopoulos A. A. *Mastering Bitcoin: Unlocking Digital Cryptocurrencies*. O'Reilly Media, 2014.
2. Dionisio A., Menezes R, and Menezes D. A. An econophysics approach to analyse uncertainty in financial markets: an application to the portuguese stock market. *The European Physical Journal*, 2006.
3. Krennikov A. The financial heat machine: coupling with the present financial crises. *Wilmott*, 2012.
4. Rényi A. Probability theory. *North-Holland Series in Applied Mathematics and Mechanics*, 1970.
5. Brissaud J. B. The meanings of entropy. *Entropy*, 2005.
6. BaFin. Merkblatt: Hinweise zum tatbestand des kryptoverwahrgeschäfts. 2020.
7. Philippatos G. C. and Wilson C. J. Entropy, market risk and the selection of efficient portfolios. *Applied Economics*, 1972.
8. Tsallis C. Possible generalization of boltzmann-gibbs statistics. *Springer-Verlag*, 1988.
9. Shannon C. E. and Weaver W. The mathematical theory of communication. *University of Illinois Press*, 1949.
10. Smith E. and Foley D. K. Is utility theory so different from thermodynamics? *SFI Preprint*, 2002.
11. Fama E. F. and French K. R. Dividend yields and expected stock returns. *Journal of Financial Economics*, 1988.
12. Perasan M. H. and Timmermann A. Predictability of stock returns: Robustness and economic significance. *The Journal of Finance*, 1995.

13. Usta I. and Kantar Y. M. Mean-variance-skewness-entropy measures: a multi objective approach for portfolio selection. *Entropy*, 2011.
14. IFRS. Holdings of cryptocurrencies - agenda paper 12. *IFRS Staff Paper*, 2019.
15. Chen J., Duan K., Zhang R., Zeng L., and Wang W. An ai based super nodes selection algorithm in blockchain networks. *arXiv:1808.00216*, 2018.
16. McCauley J. Thermodynamic analogies in economics and finance: instability of markets. *MPRA*, 2004.
17. Laffont J.J., Ossard H., and Vuong Q. Econometrics of first-price auctions. *Econometrica*, 1995.
18. Huang K. *Statistical Mechanics*. Wiley, 2008.
19. Adler R. L., Konheim A. G., and McAndrew M. H. Topological entropy. *American Mathematical Society*, 1965.
20. Boltzmann L. Vorlesungen über gastheorie. *Leipzig: J. A. Barth.*, 1896.
21. Gulko L. The entropy theory of stock option pricing. *Int. J. Theoretical Appl. Finance*, 1999.
22. Falcioni M. and Vulpiani A. *Meccanica Statistica Elementare*. Springer-Verlag, 2015.
23. Ormos M. and Zibriczky D. Entropy-based financial asset pricing. *PLoS ONE 9(12):e115742*, 2014.
24. Pincus S. M. Approximate entropy as a measure of system complexity. *Proceedings of the National Academy of Science*, 1991.
25. Qi M. and Maddala G. S. Economic factors and the stock market: a new perspective. *Wiley*, 1999.
26. Sergeev V. M. Rationality, property rights, and thermodynamic approach to market equilibrium. *Journal of Mathematical Sciences*, 2006.
27. Sheraz M., Dedu S., and Preda V. Entropy measures for asserring volatile markets. *Procedia Economics and Finance*, 2014.
28. Szmigiera M. Number of blockchain wallet users globally 2016-2019. *Statista*, 2020.
29. Nawrocki D. N. and Harding W. H. State-value weighted entropy as a measure of investment risk. *Applied Economics*, 1985.
30. Jana P., Roy T. K., and Mazumder S. K. Multi-objective possibilistic model for portfolio selection with transaction cost. *J. Comput. Appl. Math.*, 2009.
31. Clausius R. Ueber verschiedene für die anwendung bequeme formen der hauptgleichungen der mechanischen wärmetheorie. *Ann. Phys. Chem.*, 1865.
32. Corbet S., Lucey B., Urquhart A., and Yarovaya L. Cryptocurrencies as a financial asset: A systematic analysis. *Elsevier*, 2019.
33. Nakamoto S. Bitcoin: A per-to-peer electronic cash system. *Bitcoin.org*, 2008.
34. Viaggiu S., Lionetto A., Bargigli L., and Longo M. Statistical ensembles for money and debt. *arXiv:1109.0891*, 2018.
35. Kirchner U. and Zunchel C. Measuring portfolio diversification. *arXiv:1102.4722*, 2011.
36. Zakiras V., Christopoulos A. G., and Rendoumis V. L. A thermodynamic description of the time evolution of a stock market index. *European Journal of Economics, Finance and Administrative Sciences*, 2009.

37. Chen W., Wang Z., Xie H., and Yu W. Characterization of surface emg signal based on fuzzy entropy. *IEEE Transaction*, 2007.
38. Gibbs J. W. On the equilibrium of heterogeneous substances. *Am. J. Sci.*, 1878.
39. Campbell J. Y. and Shiller R. J. Stock prices, earnings, and expected dividends. *The Journal of Finance*, 1988.